

1-1-2007

Trench's Perturbation Theorem for Dynamic Equations

Stevo Stevic

Martin Bohner

Missouri University of Science and Technology, bohner@mst.edu

Follow this and additional works at: http://scholarsmine.mst.edu/math_stat_facwork



Part of the [Mathematics Commons](#), and the [Statistics and Probability Commons](#)

Recommended Citation

S. Stevic and M. Bohner, "Trench's Perturbation Theorem for Dynamic Equations," *Discrete Dynamics in Nature and Society*, Hindawi Publishing Corporation, Jan 2007.

The definitive version is available at <https://doi.org/10.1155/2007/75672>

This Article - Journal is brought to you for free and open access by Scholars' Mine. It has been accepted for inclusion in Mathematics and Statistics Faculty Research & Creative Works by an authorized administrator of Scholars' Mine. This work is protected by U. S. Copyright Law. Unauthorized use including reproduction for redistribution requires the permission of the copyright holder. For more information, please contact scholarsmine@mst.edu.

Research Article

Trench's Perturbation Theorem for Dynamic Equations

Martin Bohner and Stevo Stević

Received 10 June 2007; Accepted 31 October 2007

We consider a nonoscillatory second-order linear dynamic equation on a time scale together with a linear perturbation of this equation and give conditions on the perturbation that guarantee that the perturbed equation is also nonoscillatory and has solutions that behave asymptotically like a recessive and dominant solutions of the unperturbed equation. As the theory of time scales unifies continuous and discrete analysis, our results contain as special cases results for corresponding differential and difference equations by William F. Trench.

Copyright © 2007 M. Bohner and S. Stević. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

We consider the second-order linear dynamic equation

$$(r(t)x^\Delta)^\Delta + p(t)x^\sigma = 0, \quad t \in \mathbb{T}, \quad (1.1)$$

together with its linear perturbation

$$(r(t)y^\Delta)^\Delta + p(t)y^\sigma = f(t)y^\sigma, \quad t \in \mathbb{T}, \quad (1.2)$$

where we assume that \mathbb{T} is a time scale, that is, a nonempty closed subset of the real numbers \mathbb{T} that is unbounded above, $p : \mathbb{T} \rightarrow \mathbb{R}$, $r : \mathbb{T} \rightarrow \mathbb{R}^+$, $f : \mathbb{T} \rightarrow \mathbb{C}$, are rd-continuous, and (1.1) is nonoscillatory, that is, $rxx^\sigma > 0$ eventually for all solutions of (1.1). For the theory of time scales, we refer the reader to [1, 2] and we mention here only that the forward shift and the derivative of a function $z : \mathbb{T} \rightarrow \mathbb{R}$ are given by $z^\sigma(t) = z(t)$ and $z^\Delta(t) = z'(t)$ if $\mathbb{T} = \mathbb{R}$ and $z^\sigma(t) = z(t+1)$ and $z^\Delta(t) = z(t+1) - z(t)$ if $\mathbb{T} = \mathbb{Z}$, and that the product rule

2 Discrete Dynamics in Nature and Society

and the quotient rule for differentiable $z_1, z_2 : \mathbb{T} \rightarrow \mathbb{R}$ read [1, Theorem 1.20]

$$(z_1 z_2)^\Delta = z_1^\Delta z_2 + z_1^\sigma z_2^\Delta, \quad \left(\frac{z_1}{z_2}\right)^\Delta = \frac{z_1^\Delta z_2 - z_1 z_2^\Delta}{z_2 z_2^\sigma}, \quad (1.3)$$

where $z^\sigma = z \circ \sigma$ for a function $z : \mathbb{T} \rightarrow \mathbb{R}$. By [1, Theorem 4.61], since (1.1) is nonoscillatory, there exists a solution x_1 of (1.1) such that $\lim_{t \rightarrow \infty} (u(t)/x_1(t)) = \infty$ for all of x_1 linearly independent solutions u of (1.1).

Let u be any solution of (1.1) that is linearly independent of x_1 , that is,

$$r(u^\Delta x_1 - u x_1^\Delta) \equiv c \neq 0, \quad (1.4)$$

and define $x_2 = u/c$ so that

$$r(x_2^\Delta x_1 - x_2 x_1^\Delta) = 1, \quad \lim_{t \rightarrow \infty} \frac{x_2(t)}{x_1(t)} = \infty, \quad (1.5)$$

which implies that

$$\varphi := \frac{x_2}{x_1} \text{ satisfies } \varphi^\Delta = \frac{1}{r x_1 x_1^\sigma} > 0, \quad \lim_{t \rightarrow \infty} \varphi(t) = \infty. \quad (1.6)$$

Note that x_1 and x_2 are called *recessive* and *dominant* solutions of the nonoscillatory dynamic equation (1.1).

The main result of this paper gives conditions on f , x_1 , and x_2 that guarantee that the perturbed equation (1.2) is also nonoscillatory and has solutions y_1 and y_2 that behave asymptotically like x_1 and x_2 . Our results unify corresponding results by William F. Trench for differential equations [3, 4] and for difference equations [5] and extend them to other dynamic equations, for example, to q -difference equations [6]. Note that the unification process forces us to give many of the calculations from [5] in a “shifted form” for the discrete case.

In the next section, we will derive some preparatory results while the main theorem is stated and proved in Section 3. The paper concludes with some corollaries given in Section 4.

2. Some auxiliary results

Suppose now and in the remainder of this paper that

$$\psi(t) := \int_t^\infty f(\tau) x_1(\sigma(\tau)) x_2(\sigma(\tau)) \Delta \tau \text{ converges for all } t \in \mathbb{T}. \quad (2.1)$$

Then

$$\phi(t) := \sup_{\tau \geq t} |\psi(\tau)| \rightarrow 0 \text{ monotonically as } t \rightarrow \infty. \quad (2.2)$$

LEMMA 2.1. Using notation (2.2) and condition (2.1),

$$G(t) := \int_t^\infty f(\tau)x_1^2(\sigma(\tau))\Delta\tau \quad \text{is well defined for all } t \in \mathbb{T} \quad (2.3)$$

and satisfies

$$|G| \leq \frac{2\phi}{\phi^\sigma} \quad \text{on } \mathbb{T}. \quad (2.4)$$

Proof. We use (1.6), the product rule, the definition of the integral, and [1, Theorems 1.75 and 1.16(iv)] to find

$$\begin{aligned} G(t) &= \int_t^\infty f(\tau)x_1(\sigma(\tau))x_2(\sigma(\tau))\frac{1}{\phi(\sigma(\tau))}\Delta\tau = - \int_t^\infty \psi^\Delta(\tau)\frac{1}{\phi(\sigma(\tau))}\Delta\tau \\ &= - \int_t^\infty \left[\left(\psi \cdot \frac{1}{\phi} \right)^\Delta(\tau) - \psi(\tau)\left(\frac{1}{\phi}\right)^\Delta(\tau) \right] \Delta\tau = \frac{\psi(t)}{\phi(t)} + \int_t^\infty \psi(\tau)\left(\frac{1}{\phi}\right)^\Delta(\tau)\Delta\tau \\ &= \frac{\psi(t)}{\phi(t)} + \int_{\sigma(t)}^\infty \psi(\tau)\left(\frac{1}{\phi}\right)^\Delta(\tau)\Delta\tau + \psi(t)\left[\left(\frac{1}{\phi}\right)(\sigma(t)) - \left(\frac{1}{\phi}\right)(t) \right] \\ &= \frac{\psi(t)}{\phi^\sigma(t)} + \int_{\sigma(t)}^\infty \psi(\tau)\left(\frac{1}{\phi}\right)^\Delta(\tau)\Delta\tau, \end{aligned} \quad (2.5)$$

and therefore

$$\begin{aligned} |G(t)| &\leq \frac{\phi(t)}{\phi^\sigma(t)} + \phi(t) \int_{\sigma(t)}^\infty \left| \left(\frac{1}{\phi}\right)^\Delta(\tau) \right| \Delta\tau \\ &= \frac{\phi(t)}{\phi^\sigma(t)} - \phi(t) \int_{\sigma(t)}^\infty \left(\frac{1}{\phi}\right)^\Delta(\tau)\Delta\tau = \frac{\phi(t)}{\phi^\sigma(t)} + \frac{\phi(t)}{\phi^\sigma(t)}, \end{aligned} \quad (2.6)$$

where we have used (1.6), (2.2), and its consequence $\phi^\sigma \leq \phi$. \square

In the sequel, we use the Landau “O” symbol defined in the standard way for asymptotic behavior as $t \rightarrow \infty$ and consider the collection of differentiable functions

$$\mathcal{B} := \left\{ z : \mathbb{T} \rightarrow \mathbb{R} : z = O(\phi), z^\Delta = O\left(\frac{\phi\phi^\Delta}{\phi^\sigma}\right) \right\}, \quad (2.7)$$

which can easily be seen to be a Banach space when equipped with the norm

$$\|z\| = \|z\|_{\mathcal{B}} := \left\| \max \left\{ \frac{|z|}{\phi}, \frac{|z^\Delta| \phi^\sigma}{\phi\phi^\Delta} \right\} \right\|_\infty \quad \text{for } z \in \mathcal{B}. \quad (2.8)$$

LEMMA 2.2. Using notations (2.2), (2.3), (2.7), (2.8) and under condition (2.1),

$$u := \psi - \phi G \quad \text{satisfies } u \in \mathcal{B}, \quad \|u\|_{\mathcal{B}} \leq 2. \quad (2.9)$$

4 Discrete Dynamics in Nature and Society

Proof. Note that the fourth equal sign of the first calculation in the proof of Lemma 2.1 implies

$$u(t) = -\varphi(t) \int_t^\infty \psi(\tau) \left(\frac{1}{\varphi}\right)^\Delta(\tau) \Delta\tau, \quad (2.10)$$

and therefore

$$|u(t)| \leq \varphi(t)\phi(t) \int_t^\infty \left|\left(\frac{1}{\varphi}\right)^\Delta(\tau)\right| \Delta\tau = -\varphi(t)\phi(t) \int_t^\infty \left(\frac{1}{\varphi}\right)^\Delta(\tau) \Delta\tau = \phi(t) \quad (2.11)$$

due to (2.2) and (1.6). By the product rule and (1.6),

$$\begin{aligned} u^\Delta(t) &= \psi^\Delta(t) - \varphi^\Delta(t)G(t) - \varphi(\sigma(t))G^\Delta(t) \\ &= -f(t)x_1(\sigma(t))x_2(\sigma(t)) - \varphi^\Delta(t)G(t) + \varphi(\sigma(t))f(t)x_1^2(\sigma(t)) = -\varphi^\Delta(t)G(t), \end{aligned} \quad (2.12)$$

and therefore

$$|u^\Delta(t)| = \varphi^\Delta(t) |G(t)| \leq \frac{2\varphi^\Delta(t)\phi(t)}{\varphi^\sigma(t)} \quad (2.13)$$

due to (2.4) from Lemma 2.1. \square

For $z \in \mathcal{B}$, let us define the operator

$$(\mathcal{L}z)(t) := \int_t^\infty [x_2(\sigma(\tau)) - x_1(\sigma(\tau))\varphi(t)]f(\tau)x_1(\sigma(\tau))z(\sigma(\tau))\Delta\tau. \quad (2.14)$$

LEMMA 2.3. *Using notations (2.2), (2.3), (2.7), (2.8), (2.14), and under the conditions (2.1) and*

$$\tilde{\phi}(t) := \int_t^\infty |G(\tau)|\phi(\tau)\varphi^\Delta(\tau)\Delta\tau < \infty, \quad \nu := \limsup_{t \rightarrow \infty} \frac{\tilde{\phi}(t)}{\phi(t)} < \frac{1}{2}, \quad (2.15)$$

we have that, on $[T, \infty)$ for sufficiently large $T \in \mathbb{T}$,

$$\mathcal{L} : \mathcal{B} \longrightarrow \mathcal{B} \text{ is a contraction.} \quad (2.16)$$

Proof. Let $z \in \mathcal{B}$. For $T \geq t$, consider

$$w(z, T)(t) := \int_t^T [x_2(\sigma(\tau)) - x_1(\sigma(\tau))\varphi(t)]f(\tau)x_1(\sigma(\tau))z(\sigma(\tau))\Delta\tau. \quad (2.17)$$

Then by (1.6), (2.3), and the product rule,

$$\begin{aligned}
w(z, T)(t) &= \int_t^T [\varphi(\sigma(\tau)) - \varphi(t)] x_1(\sigma(\tau)) f(\tau) x_1(\sigma(\tau)) z(\sigma(\tau)) \Delta\tau \\
&= - \int_t^T [\varphi(\sigma(\tau)) - \varphi(t)] z(\sigma(\tau)) G^\Delta(\tau) \Delta\tau \\
&= - \int_t^T \{ [(\varphi - \varphi(t)) z G]^\Delta(\tau) - [(\varphi - \varphi(t)) z]^\Delta(\tau) G(\tau) \} \Delta\tau \\
&= - [\varphi(T) - \varphi(t)] z(T) G(T) + \int_t^T [\varphi(\sigma(\tau)) - \varphi(t)] z^\Delta(\tau) G(\tau) \Delta\tau \\
&\quad + \int_t^T \varphi^\Delta(\tau) z(\tau) G(\tau) \Delta\tau.
\end{aligned} \tag{2.18}$$

We estimate each of these three terms separately

$$\begin{aligned}
|[\varphi(T) - \varphi(t)] z(T) G(T)| &\leq |\varphi(T) - \varphi(t)| |z(T)| \frac{2\phi(T)}{\phi(\sigma(T))} \\
&\leq 2\phi(T) \|z\| \phi(T) \frac{\phi(T)}{\phi(\sigma(T))} \leq 2\|z\| \phi^2(T)
\end{aligned} \tag{2.19}$$

due to (1.6), (2.4), and (2.8);

$$|\varphi^\Delta(\tau) z(\tau) G(\tau)| \leq \varphi^\Delta(\tau) \|z\| \phi(\tau) |G(\tau)| \tag{2.20}$$

due to (1.6) and (2.8); and

$$\begin{aligned}
|[\varphi(\sigma(\tau)) - \varphi(t)] z^\Delta(\tau) G(\tau)| &\leq |\varphi(\sigma(\tau)) - \varphi(t)| \|z\| \frac{\phi(\tau) \varphi^\Delta(\tau)}{\varphi^\sigma(\tau)} |G(\tau)| \\
&\leq \|z\| \phi(\tau) \varphi^\Delta(\tau) |G(\tau)| \quad \text{for } t \leq \sigma(\tau),
\end{aligned} \tag{2.21}$$

due to (1.6) and (2.8). Altogether,

$$|w(z, T)(t)| \leq 2\|z\| \left\{ \phi^2(T) + \int_t^T |G(\tau)| \phi(\tau) \varphi^\Delta(\tau) \Delta\tau \right\}. \tag{2.22}$$

Thus, using (2.2) and (2.15),

$$|(\mathcal{L}z)(t)| \leq 2\|z\| \tilde{\phi}(t) \tag{2.23}$$

so that with $\theta := 2\nu < 1$,

$$\left| \frac{(\mathcal{L}z)(t)}{\phi(t)} \right| \leq 2\|z\| \frac{\tilde{\phi}(t)}{\phi(t)} \leq 2\|z\| \nu = \theta \|z\|. \tag{2.24}$$

Next, from the first equal sign in the first calculation of this proof,

$$(\mathcal{L}z)(t) = \int_t^\infty [\varphi(\sigma(\tau)) - \varphi(t)] f(\tau) x_1^2(\sigma(\tau)) z(\sigma(\tau)) \Delta\tau, \tag{2.25}$$

so that by [1, Theorem 1.117(ii)] and the product rule,

$$\begin{aligned} (\mathcal{L}z)^\Delta(t) &= - \int_t^\infty \varphi^\Delta(t) f(\tau) x_1^2(\sigma(\tau)) z(\sigma(\tau)) \Delta\tau = \varphi^\Delta(t) \int_t^\infty G^\Delta(\tau) z(\sigma(\tau)) \Delta\tau \\ &= \varphi^\Delta(t) \int_t^\infty \{(zG)^\Delta(\tau) - z^\Delta(\tau)G(\tau)\} \Delta\tau = -\varphi^\Delta(t) \left\{ z(t)G(t) + \int_t^\infty G(\tau)z^\Delta(\tau)\Delta\tau \right\}. \end{aligned} \quad (2.26)$$

Thus, using (2.7), (2.8), (2.4), (1.6), (2.15),

$$\begin{aligned} |(\mathcal{L}z)^\Delta(t)| &\leq \varphi^\Delta(t) \left\{ \|z\| \phi(t) \frac{2\phi(t)}{\varphi^\sigma(t)} + \int_t^\infty |G(\tau)| \|z\| \frac{\phi(\tau)\varphi^\Delta(\tau)}{\varphi^\sigma(\tau)} \Delta\tau \right\} \\ &\leq \frac{\|z\|\varphi^\Delta(t)}{\varphi^\sigma(t)} \{2\phi^2(t) + \tilde{\phi}(t)\} \end{aligned} \quad (2.27)$$

and therefore,

$$\left| \frac{(\mathcal{L}z)^\Delta(t)}{\phi(t)\varphi^\Delta(t)/\varphi^\sigma(t)} \right| \leq \|z\| \left\{ 2\phi(t) + \frac{\tilde{\phi}(t)}{\phi(t)} \right\} \leq \|z\| \{2\phi(t) + \nu\} < \theta \|z\| \quad (2.28)$$

eventually due to (2.2) and (2.15). Hence, $\mathcal{L}z \in \mathcal{B}$ and

$$\|\mathcal{L}z\| \leq \theta \|z\| \quad \forall z \in \mathcal{B} \quad (2.29)$$

on $[T, \infty)$ for sufficiently large $T \in \mathbb{T}$. □

3. The perturbation result

In this section, we use the auxiliary results from Section 2 to prove the following main theorem of this paper.

THEOREM 3.1. *Let x_1 and x_2 be the recessive and dominant solutions of the unperturbed linear second-order dynamic equation (1.1) satisfying (1.6). Using notations (2.2) and (2.3) and under the conditions (2.1) and (2.15), there exist solutions y_1 and y_2 of the perturbed dynamic equation (1.2) such that*

$$\frac{y_1}{x_1} = 1 + O(\phi), \quad \left(\frac{y_1}{x_1} \right)^\Delta = O\left(\frac{\phi\varphi^\Delta}{\varphi^\sigma} \right), \quad (3.1)$$

$$\frac{y_2}{x_2} = 1 + O(\hat{\phi}), \quad \left(\frac{y_2}{x_2} \right)^\Delta = O\left(\frac{\hat{\phi}\varphi^\Delta}{\varphi^\sigma} \right), \quad (3.2)$$

where $\hat{\phi}$ is defined by

$$\hat{\phi}(t) := \frac{1}{\varphi(t)} \int_{t_0}^t \phi(\tau)\varphi^\Delta(\tau)\Delta\tau. \quad (3.3)$$

Proof. We use the notation (2.7), (2.8), (2.9), and (2.14) and define

$$\mathcal{T}z := u + \mathcal{L}z \quad \text{for } z \in \mathcal{B}. \quad (3.4)$$

By Lemmas 2.2 and 2.3, $\mathcal{T} : \mathcal{B} \rightarrow \mathcal{B}$ is a contraction. By Banach's fixed point theorem, \mathcal{T} has a unique fixed point $\zeta \in \mathcal{B}$. Hence, $\zeta = \mathcal{T}\zeta = u + \mathcal{L}\zeta$. Define

$$y_1 := x_1(1 + \zeta). \quad (3.5)$$

We show that y_1 solves (1.2). First, from the proofs of Lemmas 2.2 and 2.3,

$$-r\zeta^\Delta = -r[u^\Delta + (\mathcal{L}\zeta)^\Delta] = r\varphi^\Delta G + r\varphi^\Delta(\zeta G + \tilde{\zeta}) = \frac{G(1 + \zeta) + \tilde{\zeta}}{x_1 x_1^\sigma}, \quad (3.6)$$

where $\tilde{\zeta}^\Delta := -G\zeta^\Delta$. So we have

$$y_1^\Delta = x_1^\Delta(1 + \zeta) + x_1^\sigma \zeta^\Delta \quad (3.7)$$

and hence,

$$r y_1^\Delta = r x_1^\Delta(1 + \zeta) - \frac{G(1 + \zeta) + \tilde{\zeta}}{x_1} \quad (3.8)$$

so that, by using (3.6),

$$\begin{aligned} (r y_1^\Delta)^\Delta &= (r x_1^\Delta)^\Delta(1 + \zeta)^\sigma + r x_1^\Delta \zeta^\Delta - \left(\frac{G(1 + \zeta) + \tilde{\zeta}}{x_1} \right)^\Delta \\ &= -p y_1^\sigma + r \zeta^\Delta x_1^\Delta - \frac{(G^\Delta(1 + \zeta)^\sigma + G\zeta^\Delta + \tilde{\zeta}^\Delta)x_1 - [G(1 + \zeta) + \tilde{\zeta}]x_1^\Delta}{x_1 x_1^\sigma} \\ &= -p y_1^\sigma + r \zeta^\Delta x_1^\Delta + f y_1^\sigma + \frac{G(1 + \zeta) + \tilde{\zeta}}{x_1 x_1^\sigma} x_1^\Delta = -p y_1^\sigma + f y_1^\sigma, \end{aligned} \quad (3.9)$$

and y_1 indeed solves (1.2). From (2.7) and since $\zeta \in \mathcal{B}$, (3.1) holds true. Also, by (2.2) and since $\zeta \in \mathcal{B}$, $r y_1 y_1^\sigma = r x_1 x_1^\sigma(1 + \zeta)(1 + \zeta^\sigma) > 0$ eventually, say without loss of generality on $[t_0, \infty)$. So we may define

$$y_2 := y_1(c + \xi) \quad \text{with } c = \frac{x_2(t_0)}{x_1(t_0)}, \quad \xi(t) = \int_{t_0}^t \frac{\Delta\tau}{r(\tau)y_1(\tau)y_1(\sigma(\tau))}. \quad (3.10)$$

We have

$$y_2^\Delta = y_1^\Delta(c + \xi) + y_1^\sigma \xi^\Delta = y_1^\Delta(c + \xi) + \frac{1}{r y_1} \quad (3.11)$$

and hence,

$$(r y_2^\Delta)^\Delta = (r y_1^\Delta)^\Delta(c + \xi)^\sigma + r y_1^\Delta \xi^\Delta - \frac{y_1^\Delta}{y_1 y_1^\sigma} = (-p y_1^\sigma + f y_1^\sigma)(c + \xi)^\sigma = -p y_2^\sigma + f y_2^\sigma, \quad (3.12)$$

so that y_2 is another (nonoscillatory) solution of (1.2). Next, by (1.6),

$$x_2 = x_1\varphi, \quad \varphi(t) = c + \int_{t_0}^t \frac{\Delta\tau}{r(\tau)x_1(\tau)x_1(\sigma(\tau))} \quad (3.13)$$

and therefore,

$$\frac{y_2}{x_2} = \frac{y_1}{x_1} \frac{c + \xi}{\varphi} = \frac{y_1}{x_1} (1 + \gamma), \quad (3.14)$$

where

$$\begin{aligned} \gamma(t) &:= \frac{c + \xi(t) - \varphi(t)}{\varphi(t)} = \frac{\xi(t) - \int_{t_0}^t \Delta\tau/r(\tau)x_1(\tau)x_1(\sigma(\tau))}{\varphi(t)} \\ &= -\frac{1}{\varphi(t)} \int_{t_0}^t \left[1 - \frac{x_1(\tau)x_1(\sigma(\tau))}{y_1(\tau)y_1(\sigma(\tau))} \right] \frac{\Delta\tau}{r(\tau)x_1(\tau)x_1(\sigma(\tau))} \\ &= -\frac{1}{\varphi(t)} \int_{t_0}^t \left(1 - \frac{x_1x_1^\sigma}{y_1y_1^\sigma} \right) (\tau) \varphi^\Delta(\tau) \Delta\tau. \end{aligned} \quad (3.15)$$

Now the first equation in (3.1) and (2.2) imply (see (3.3))

$$1 - \frac{x_1x_1^\sigma}{y_1y_1^\sigma} = O(\phi) \quad \text{and thus } \gamma = O(\hat{\phi}). \quad (3.16)$$

Then (2.2) together with (3.3) imply $\phi = O(\hat{\phi})$ so that

$$\frac{y_2}{x_2} = \frac{y_1}{x_1} (1 + \gamma) = (1 + O(\phi))(1 + O(\hat{\phi})) = 1 + O(\hat{\phi}), \quad (3.17)$$

which proves the first equation in (3.2). Finally,

$$\left(\frac{y_2}{x_2} \right)^\Delta = \left(\frac{y_1}{x_1} \right)^\Delta (1 + \gamma) + \left(\frac{y_1}{x_1} \right)^\sigma \gamma^\Delta = O\left(\frac{\phi\varphi^\Delta}{\varphi^\sigma} \right) (1 + O(\hat{\phi})) + (1 + O(\phi)) O\left(\frac{\hat{\phi}\varphi^\Delta}{\varphi^\sigma} \right), \quad (3.18)$$

because of (2.2), (3.1), and

$$\begin{aligned} \gamma^\Delta &= \frac{(c + \xi - \varphi)^\Delta \varphi - (c + \xi - \varphi) \varphi^\Delta}{\varphi\varphi^\sigma} = \frac{\xi^\Delta - \varphi^\Delta}{\varphi^\sigma} - \gamma \frac{\varphi^\Delta}{\varphi^\sigma} \\ &= -\frac{\varphi^\Delta}{\varphi^\sigma} \left(\gamma + 1 - \frac{\xi^\Delta}{\varphi^\Delta} \right) = O\left(\frac{\varphi^\Delta}{\varphi^\sigma} \right) [O(\hat{\phi}) + O(\phi)] = O\left(\frac{\hat{\phi}\varphi^\Delta}{\varphi^\sigma} \right). \end{aligned} \quad (3.19)$$

So (3.2) is established and this completes the proof. \square

4. Applications

In this last section, we state and prove some simple consequences of Theorem 3.1.

COROLLARY 4.1. *The conclusions of Theorem 3.1 hold if*

$$\int_t^\infty |f(\tau)|x_1(\sigma(\tau))x_2(\sigma(\tau))\Delta\tau < \infty. \quad (4.1)$$

Proof. Clearly, (4.1) implies (2.1). Define

$$\hat{\psi}(t) := \int_t^\infty |f(\tau)|x_1(\sigma(\tau))x_2(\sigma(\tau))\Delta\tau, \quad \hat{G} := \int_t^\infty |f(\tau)|x_1^2(\sigma(\tau))\Delta\tau. \quad (4.2)$$

Then

$$|G(t)| \leq \hat{G}(t) = \int_t^\infty \frac{|f(\tau)|x_1(\sigma(\tau))x_2(\sigma(\tau))}{\varphi(\sigma(\tau))}\Delta\tau \leq \frac{\hat{\psi}(t)}{\varphi(\sigma(t))}. \quad (4.3)$$

For $T \geq t$, we have by the product rule that

$$\begin{aligned} & \int_t^T |f(\tau)|x_1(\sigma(\tau))x_2(\sigma(\tau))\Delta\tau \\ &= \int_t^T |f(\tau)|x_1^2(\sigma(\tau))\varphi(\sigma(\tau))\Delta\tau = - \int_t^T \hat{G}^\Delta(\tau)\varphi(\sigma(\tau))\Delta\tau \\ &= - \int_t^T [(\hat{G}\varphi)^\Delta(\tau) - \hat{G}(\tau)\varphi^\Delta(\tau)]\Delta\tau = -(\hat{G}\varphi)(T) + (\hat{G}\varphi)(t) + \int_t^T \hat{G}(\tau)\varphi^\Delta(\tau)\Delta\tau. \end{aligned} \quad (4.4)$$

Note now that $(\hat{G}\varphi)(T) \rightarrow 0$ as $T \rightarrow \infty$ since

$$0 \leq \hat{G}(T)\varphi(T) \leq \frac{\varphi(T)\hat{\psi}(T)}{\varphi(\sigma(T))} \leq \hat{\psi}(T) \rightarrow 0 \quad \text{as } T \rightarrow \infty \quad (4.5)$$

due to (4.3) and (4.1). Note also that the quantity on the right of the last equal sign in the above calculation must have a finite limit as $T \rightarrow \infty$ since the quantity on the left of the first equal sign converges as $T \rightarrow \infty$ due to (4.1). Therefore,

$$\int_t^\infty \hat{G}(\tau)\varphi^\Delta(\tau)\Delta\tau < \infty, \quad \text{and thus by (4.3),} \quad \int_t^\infty |G(\tau)|\varphi^\Delta(\tau)\Delta\tau < \infty. \quad (4.6)$$

Hence, using (2.2),

$$\frac{\tilde{\phi}(t)}{\phi(t)} = \int_t^\infty |G(\tau)| \frac{\phi(\tau)}{\phi(t)} \varphi^\Delta(\tau)\Delta\tau \leq \int_t^\infty |G(\tau)|\varphi^\Delta(\tau)\Delta\tau \rightarrow 0 \quad \text{as } t \rightarrow \infty \quad (4.7)$$

so that (2.15) holds with $\nu = 0$. □

COROLLARY 4.2. *Assume (2.1). If ϕ satisfies (2.2) and*

$$\limsup_{t \rightarrow \infty} \left\{ \frac{1}{\phi(t)} \int_t^\infty \frac{\phi^2(\tau)\varphi^\Delta(\tau)}{\varphi(\sigma(\tau))}\Delta\tau \right\} =: \tilde{\gamma} < \frac{1}{4}, \quad (4.8)$$

then the conclusions of Theorem 3.1 hold.

Proof. From (2.4), we have

$$\frac{\tilde{\phi}(t)}{\phi(t)} = \int_t^\infty |G(\tau)| \frac{\phi(\tau)}{\phi(t)} \varphi^\Delta(\tau) \Delta\tau \leq \int_t^\infty \frac{2\phi^2(\tau)}{\varphi(\sigma(\tau))\phi(t)} \varphi^\Delta(\tau) \Delta\tau \quad (4.9)$$

so that (4.8) implies

$$\limsup_{t \rightarrow \infty} \frac{\tilde{\phi}(t)}{\phi(t)} \leq 2\tilde{\gamma} < \frac{1}{2}, \quad (4.10)$$

and therefore (2.15) is satisfied. \square

COROLLARY 4.3. *Assume that α is a positive function tending monotonically to ∞ and define*

$$S(t) := \int_{t_0}^t \alpha(\sigma(\tau)) f(\tau) x_1(\sigma(\tau)) x_2(\sigma(\tau)) \Delta\tau. \quad (4.11)$$

If $|S(t)| \leq A < \infty$ and

$$\limsup_{t \rightarrow \infty} \left\{ \alpha(\sigma(t)) \int_{t_0}^\infty \frac{\varphi^\Delta(\tau)}{(\alpha^2\varphi)(\sigma(\tau))} \Delta\tau \right\} =: \hat{\gamma} < \frac{1}{4A}, \quad (4.12)$$

then the conclusions of Theorem 3.1 hold with

$$\phi(t) = \frac{1}{\alpha(\sigma(t))}, \quad \hat{\phi}(t) = \frac{1}{\varphi(t)} \int_{t_0}^t \frac{\varphi^\Delta(\tau)}{\alpha(\sigma(\tau))} \Delta\tau. \quad (4.13)$$

Proof. Using the definition of S , we find that

$$\psi(t) = \int_t^\infty \frac{S^\Delta(\tau)}{\alpha(\sigma(\tau))} \Delta\tau, \quad G(t) = \int_t^\infty \frac{S^\Delta(\tau)}{(\alpha\varphi)(\sigma(\tau))} \Delta\tau. \quad (4.14)$$

Similar to the proof of Lemma 2.1, using integration by parts, we get

$$|\psi(t)| \leq \frac{2A}{\alpha(\sigma(\tau))}, \quad |G(t)| \leq \frac{2A}{(\alpha\varphi)(\sigma(\tau))}. \quad (4.15)$$

Hence, we can choose ϕ as in the statement of the corollary in place of (2.2) and all results as presented in this paper still hold with this ϕ . Now

$$\frac{\tilde{\phi}(t)}{\phi(t)} \leq \frac{1}{\phi(t)} \int_t^\infty \frac{2A\phi(\tau)\varphi^\Delta(\tau)}{(\alpha\varphi)(\sigma(\tau))} \Delta\tau = \alpha(\sigma(t)) \int_t^\infty \frac{2A\varphi^\Delta(\tau)}{(\alpha^2\varphi)(\sigma(\tau))} \Delta\tau, \quad (4.16)$$

so that (4.12) implies

$$\limsup_{t \rightarrow \infty} \frac{\tilde{\phi}(t)}{\phi(t)} \leq 2A\hat{\gamma} < \frac{1}{2}, \quad (4.17)$$

and therefore, (2.15) is satisfied. \square

References

- [1] M. Bohner and A. Peterson, *Dynamic Equations on Time Scales: An Introduction with Applications*, Birkhäuser, Boston, Mass, USA, 2001.
- [2] M. Bohner and A. Peterson, Eds., *Advances in Dynamic Equations on Time Scales*, Birkhäuser, Boston, Mass, USA, 2003.
- [3] W. F. Trench, “Linear perturbations of a nonoscillatory second order equation,” *Proceedings of the American Mathematical Society*, vol. 97, no. 3, pp. 423–428, 1986.
- [4] W. F. Trench, “Linear perturbations of a nonoscillatory second order differential equation. II,” *Proceedings of the American Mathematical Society*, vol. 131, no. 5, pp. 1415–1422, 2003.
- [5] W. F. Trench, “Linear perturbations of a nonoscillatory second order difference equation,” *Journal of Mathematical Analysis and Applications*, vol. 255, no. 2, pp. 627–635, 2001.
- [6] V. Kac and P. Cheung, *Quantum Calculus*, Universitext, Springer, New York, NY, USA, 2002.

Martin Bohner: Department of Mathematics and Statistics,
Missouri University of Science and Technology, MO 65401, USA
Email address: bohner@mst.edu

Stevo Stević: Mathematical Institute of the Serbian Academy of Science, Knez Mihailova 36/III,
11000 Beograd, Serbia
Email addresses: sstevic@ptt.yu; sstevo@matf.bg.ac.yu

Special Issue on Intelligent Computational Methods for Financial Engineering

Call for Papers

As a multidisciplinary field, financial engineering is becoming increasingly important in today's economic and financial world, especially in areas such as portfolio management, asset valuation and prediction, fraud detection, and credit risk management. For example, in a credit risk context, the recently approved Basel II guidelines advise financial institutions to build comprehensible credit risk models in order to optimize their capital allocation policy. Computational methods are being intensively studied and applied to improve the quality of the financial decisions that need to be made. Until now, computational methods and models are central to the analysis of economic and financial decisions.

However, more and more researchers have found that the financial environment is not ruled by mathematical distributions or statistical models. In such situations, some attempts have also been made to develop financial engineering models using intelligent computing approaches. For example, an artificial neural network (ANN) is a nonparametric estimation technique which does not make any distributional assumptions regarding the underlying asset. Instead, ANN approach develops a model using sets of unknown parameters and lets the optimization routine seek the best fitting parameters to obtain the desired results. The main aim of this special issue is not to merely illustrate the superior performance of a new intelligent computational method, but also to demonstrate how it can be used effectively in a financial engineering environment to improve and facilitate financial decision making. In this sense, the submissions should especially address how the results of estimated computational models (e.g., ANN, support vector machines, evolutionary algorithm, and fuzzy models) can be used to develop intelligent, easy-to-use, and/or comprehensible computational systems (e.g., decision support systems, agent-based system, and web-based systems)

This special issue will include (but not be limited to) the following topics:

- **Computational methods:** artificial intelligence, neural networks, evolutionary algorithms, fuzzy inference, hybrid learning, ensemble learning, cooperative learning, multiagent learning

- **Application fields:** asset valuation and prediction, asset allocation and portfolio selection, bankruptcy prediction, fraud detection, credit risk management
- **Implementation aspects:** decision support systems, expert systems, information systems, intelligent agents, web service, monitoring, deployment, implementation

Authors should follow the Journal of Applied Mathematics and Decision Sciences manuscript format described at the journal site <http://www.hindawi.com/journals/jamds/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/>, according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

Guest Editors

Lean Yu, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; Department of Management Sciences, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong; yulean@amss.ac.cn

Shouyang Wang, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; sywang@amss.ac.cn

K. K. Lai, Department of Management Sciences, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong; mssklai@cityu.edu.hk