

01 Jan 2010

## Gronwall-Oulang-type Integral Inequalities on Time Scales

Ailian Liu

Martin Bohner

Missouri University of Science and Technology, bohner@mst.edu

Follow this and additional works at: [https://scholarsmine.mst.edu/math\\_stat\\_facwork](https://scholarsmine.mst.edu/math_stat_facwork)



Part of the [Mathematics Commons](#), and the [Statistics and Probability Commons](#)

---

### Recommended Citation

A. Liu and M. Bohner, "Gronwall-Oulang-type Integral Inequalities on Time Scales," *Journal of Inequalities and Applications*, Springer Open, Jan 2010.

The definitive version is available at <https://doi.org/10.1155/2010/275826>

This Article - Journal is brought to you for free and open access by Scholars' Mine. It has been accepted for inclusion in Mathematics and Statistics Faculty Research & Creative Works by an authorized administrator of Scholars' Mine. This work is protected by U. S. Copyright Law. Unauthorized use including reproduction for redistribution requires the permission of the copyright holder. For more information, please contact [scholarsmine@mst.edu](mailto:scholarsmine@mst.edu).

## Research Article

# Gronwall-OuIang-Type Integral Inequalities on Time Scales

**Ailian Liu<sup>1,2</sup> and Martin Bohner<sup>2</sup>**

<sup>1</sup> School of Statistics and Mathematics, Shandong Economic University, Jinan 250014, China

<sup>2</sup> Department of Mathematics and Statistics, Missouri University of Science and Technology, Rolla, MO 65409-0020, USA

Correspondence should be addressed to Martin Bohner, bohner@mst.edu

Received 20 April 2010; Accepted 3 August 2010

Academic Editor: Wing-Sum Cheung

Copyright © 2010 A. Liu and M. Bohner. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We present several Gronwall-OuIang-type integral inequalities on time scales. Firstly, an OuIang inequality on time scales is discussed. Then we extend the Gronwall-type inequalities to multiple integrals. Some special cases of our results contain continuous Gronwall-type inequalities and their discrete analogues. Several examples are included to illustrate our results at the end.

## 1. Introduction

OuIang inequalities and their generalizations have proved to be useful tools in oscillation theory, boundedness theory, stability theory, and other applications of differential and difference equations. A nice introduction to continuous and discrete OuIang inequalities can be found in [1, 2], and studies in [3–5] give some of their generalizations to multiple integrals and higher-dimensional spaces. Like Gronwall's inequality, OuIang's inequality is also used to obtain a priori bounds on unknown functions. Therefore, integral inequalities of this type are usually known as Gronwall-OuIang-type inequalities [6].

The calculus on time scales has been introduced by Hilger [7] in order to unify discrete and continuous analysis. For the general basic ideas and background, we refer to [8, 9]. In this paper, we are concerned with Gronwall-OuIang-type integral inequalities on time scales, which unify and extend the corresponding continuous inequalities and their discrete analogues. We also provide a more useful and explicit bound than that in [10–12].

## 2. OuIang Inequality

We first give Gronwall's inequality on time scales which could be found in [8, Corollary 6.7]. Throughout this section, we fix  $t_0 \in \mathbb{T}$  and let  $\mathbb{T}_{t_0}^+ = \{t \in \mathbb{T} : t \geq t_0\}$ .

**Lemma 2.1.** Let  $y \in C_{rd}$ ,  $p \in \mathcal{R}^+$ ,  $p(t) \geq 0$ , for all  $t \in \mathbb{T}_{t_0}^+$ , and  $\alpha \in \mathbb{R}$ . Then

$$y(t) \leq \alpha + \int_{t_0}^t y(\tau)p(\tau)\Delta\tau \quad \forall t \in \mathbb{T}_{t_0}^+ \quad (2.1)$$

implies that

$$y(t) \leq \alpha e_p(t, t_0) \quad \forall t \in \mathbb{T}_{t_0}^+. \quad (2.2)$$

Above,  $\mathcal{R}$  is defined as the set of all regressive and rd-continuous functions,  $\mathcal{R}^+$  is the positive regressive part of  $\mathcal{R}$ , the “circle minus” subtraction  $\ominus$  on  $\mathcal{R}$  is defined by

$$(p \ominus q)(t) := \frac{p(t) - q(t)}{1 + \mu(t)q(t)} \quad \text{with } p, q \in \mathcal{R}, \quad (2.3)$$

and  $e_p(t, t_0)$  is the exponential function on time scales; for more details on time scales, see [8, 9].

Now we will give the Oulang inequality on time scales.

**Theorem 2.2.** Let  $u$  and  $v$  be real-valued nonnegative rd-continuous functions defined on  $\mathbb{T}_{t_0}^+$ . If

$$u^2(t) \leq c + \int_{t_0}^t u(\tau)v(\tau)\Delta\tau \quad \forall t \in \mathbb{T}_{t_0}^+, \quad (2.4)$$

where  $c$  is a positive constant, then

$$u(t) \leq \sqrt{c} + \frac{1}{2} \int_{t_0}^t v(\tau)\Delta\tau \quad \forall t \in \mathbb{T}_{t_0}^+. \quad (2.5)$$

*Proof.* Let

$$w(t) = \int_{t_0}^t u(\tau)v(\tau)\Delta\tau. \quad (2.6)$$

From (2.4), we have

$$u^2(t)v^2(t) \leq v^2(t)(c + w(t)). \quad (2.7)$$

The definition of  $w$  gives

$$w^\Delta(t) \leq v(t)\sqrt{c + w(t)}, \quad (2.8)$$

Dividing both sides of (2.8) by  $\sqrt{c+w(t)}$  and integrating from  $t_0$  to  $t \in \mathbb{T}_{t_0}^+$ , we have

$$\int_{t_0}^t \frac{w^\Delta(\tau)}{\sqrt{c+w(\tau)}} \Delta\tau \leq \int_{t_0}^t v(\tau) \Delta\tau. \quad (2.9)$$

According to the chain rule [8, Theorem 1.93] and since  $w$  is increasing,

$$\begin{aligned} 2 \int_{t_0}^t (\sqrt{c+w})^\Delta(\tau) \Delta\tau &= \int_{t_0}^t \frac{2w^\Delta(\tau)}{\sqrt{c+w(\tau)} + \sqrt{c+w(\sigma(\tau))}} \Delta\tau \\ &\leq \int_{t_0}^t \frac{w^\Delta(\tau)}{\sqrt{c+w(\tau)}} \Delta\tau \\ &\leq \int_{t_0}^t v(\tau) \Delta\tau, \end{aligned} \quad (2.10)$$

so

$$\sqrt{c+w(t)} - \sqrt{c} \leq \frac{1}{2} \int_{t_0}^t v(\tau) \Delta\tau. \quad (2.11)$$

Combining (2.4) and (2.11) yields (2.5) and completes the proof.  $\square$

In 1979, Dafermos [13] published a so-called Gronwall-type inequality (see also [3]). In the same way as Theorem 2.2, we now extend this result to general time scales.

**Theorem 2.3.** *Let  $y$  and  $g$  be nonnegative rd-continuous functions on  $\mathbb{T}_{t_0}^+$ . Let  $\alpha$ ,  $M$ ,  $N$  be nonnegative constants and  $-\alpha \in \mathcal{R}^+$ . If*

$$y^2(t) \leq M^2 y^2(t_0) + 2 \int_{t_0}^t [\alpha y^2(\tau) + N g(\tau) y(\tau)] \Delta\tau \quad \forall t \in \mathbb{T}_{t_0}^+, \quad (2.12)$$

then

$$y(t) \leq M y(t_0) e_{\ominus(\alpha)}(t, t_0) + \int_{t_0}^t N g(\tau) e_{\ominus(\alpha)}(t, \tau) \Delta\tau \quad \forall t \in \mathbb{T}_{t_0}^+. \quad (2.13)$$

*Proof.* Let

$$z(t) = M^2 y^2(t_0) + 2 \int_{t_0}^t [\alpha y^2(\tau) + N g(\tau) y(\tau)] \Delta\tau. \quad (2.14)$$

Then,

$$\begin{aligned} z^\Delta(t) &= 2\alpha y^2(t) + 2Ng(t)y(t) \leq 2\alpha z(t) + 2Ng(t)\sqrt{z(t)} \\ &\leq \alpha\sqrt{z(\sigma(t))}\left(\sqrt{z(t)} + \sqrt{z(\sigma(t))}\right) + Ng(t)\left(\sqrt{z(t)} + \sqrt{z(\sigma(t))}\right). \end{aligned} \quad (2.15)$$

Hence,

$$\frac{z^\Delta(t)}{\sqrt{z(t)} + \sqrt{z(\sigma(t))}} - \alpha\sqrt{z(\sigma(t))} \leq Ng(t). \quad (2.16)$$

Multiplying both sides of (2.16) by  $e_{-\alpha}(t, t_0)$ , we have

$$(\sqrt{z}e_{-\alpha}(\cdot, t_0))^\Delta(t) \leq Ng(t)e_{-\alpha}(t, t_0). \quad (2.17)$$

Integrating (2.17) from  $t_0$  to  $t$ , we obtain that

$$\sqrt{z(t)}e_{-\alpha}(t, t_0) \leq My(t_0) + \int_{t_0}^t Ng(\tau)e_{-\alpha}(\tau, t_0)\Delta\tau. \quad (2.18)$$

Combining (2.12) and (2.18), and using [8, Theorems 2.36 and 2.48] yields (2.13) and completes the proof.  $\square$

*Remark 2.4.* If  $\alpha = 0$  and  $N = 1/2$ , then Theorem 2.3 reduces to Theorem 2.2.

*Remark 2.5.* If we multiply inequality (2.16) by another exponential function on time scales, for example,  $e_{\ominus(2\alpha)}(t, t_0)$ , we could get another kind of inequality, which is a special case of Theorem 3.4.

### 3. Gronwall-OuIang-Type Inequality

Pachpatte discussed several integral inequalities arising in the theory of differential equations and difference equations [3, 4]. Now, we extend some of these results to time scales. First, we give some notations and definitions which are used in our subsequent discussion.

To simplify the expression, we let  $0 \in \mathbb{T}$ , choose rd-continuous functions  $r_i$  ( $1 \leq i \leq n$ ) such that

$$r_i(t) > 0, \quad 1 \leq i \leq n-1, \quad r_n(t) = 1 \quad \forall t \in \mathbb{T}_0^+, \quad (3.1)$$

and define the differential operators  $L_i$ ,  $0 \leq i \leq n$ , by

$$L_0x = x, \quad L_ix = \frac{1}{r_i}(L_{i-1}x)^\Delta, \quad 1 \leq i \leq n. \quad (3.2)$$

For  $t \in \mathbb{T}_0^+$  and a nonnegative function  $r$  defined on  $\mathbb{T}_0^+$ , we set

$$A[t, r_1, \dots, r_{n-1}, r] = \int_0^t r_1(t_1) \cdots \int_0^{t_{n-2}} r_{n-1}(t_{n-1}) \int_0^{t_{n-1}} r(t_n) \Delta t_n \Delta t_{n-1} \cdots \Delta t_1. \quad (3.3)$$

**Theorem 3.1.** *Let  $F$  and  $r$  be real-valued nonnegative rd-continuous functions on  $\mathbb{T}_0^+$ , and let  $q > 1$  be a constant. If*

$$F^q(t) \leq c + A[t, r_1, \dots, r_{n-1}, rF] \quad \forall t \in \mathbb{T}_0^+, \quad (3.4)$$

where  $c > 0$  is a constant, then

$$F(t) \leq \left\{ c^{(q-1)/q} + \frac{q-1}{q} A[t, r_1, \dots, r_{n-1}, r] \right\}^{1/(q-1)} \quad \forall t \in \mathbb{T}_0^+. \quad (3.5)$$

*Proof.* Let

$$z(t) = c + A[t, r_1, \dots, r_{n-1}, rF]. \quad (3.6)$$

From (3.6), it is easy to observe that

$$L_n z = rF \leq r z^{1/q}. \quad (3.7)$$

From (3.7) and using the facts that  $z$  and  $z^\Delta$  are nonnegative, and

$$\left( z^{1/q} \right)^\Delta = \frac{1}{q} z^\Delta \int_0^1 \left( z + \mu z^\Delta h \right)^{1/q-1} dh \geq 0, \quad (3.8)$$

we have

$$\frac{L_n z}{\left( z^{1/q} \right)^\sigma} \leq \frac{L_n z}{z^{1/q}} \leq r \leq r + \frac{L_{n-1} z \left( z^{1/q} \right)^\Delta}{z^{1/q} \left( z^{1/q} \right)^\sigma}, \quad (3.9)$$

that is,

$$\left( \frac{L_{n-1} z}{z^{1/q}} \right)^\Delta \leq r. \quad (3.10)$$

Integrating (3.10) with respect to  $t_n$  from 0 to  $t$  and using the fact that  $L_{n-1} z(0) = 0$ , we obtain that

$$\frac{L_{n-1} z(t)}{z^{1/q}(t)} \leq \int_0^t r(t_n) \Delta t_n, \quad (3.11)$$

which implies that

$$\frac{(L_{n-2}z)^\Delta(t)}{r_{n-1}(t)z^{1/q}(t)} \leq \int_0^t r(t_n)\Delta t_n. \quad (3.12)$$

Again as above, from (3.12), we observe that

$$\frac{(L_{n-2}z)^\Delta(t)}{(z^{1/q})^\sigma(t)} \leq \frac{(L_{n-2}z)^\Delta(t)}{z^{1/q}(t)} \leq r_{n-1}(t) \int_0^t r(t_n)\Delta t_n \leq r_{n-1}(t) \int_0^t r(t_n)\Delta t_n + \frac{L_{n-2}z(t)(z^{1/q})^\Delta(t)}{z^{1/q}(t)(z^{1/q})^\sigma(t)}, \quad (3.13)$$

that is,

$$\left(\frac{L_{n-2}z}{z^{1/q}}\right)^\Delta(t) \leq r_{n-1}(t) \int_0^t r(t_n)\Delta t_n. \quad (3.14)$$

By setting  $t = t_{n-1}$  in (3.14) and integrating with respect to  $t_{n-1}$  from 0 to  $t$  and using the fact that  $L_{n-2}z(0) = 0$ , we get

$$\frac{L_{n-2}z(t)}{z^{1/q}(t)} \leq \int_0^t r_{n-1}(t_{n-1}) \int_0^{t_{n-1}} r(t_n)\Delta t_n \Delta t_{n-1}. \quad (3.15)$$

Continuing this way, we obtain that

$$\frac{L_1z(t)}{z^{1/q}(t)} \leq \int_0^t r_2(t_2) \cdots \int_0^{t_{n-2}} r_{n-1}(t_{n-1}) \int_0^{t_{n-1}} r(t_n)\Delta t_n \Delta t_{n-1} \cdots \Delta t_2, \quad (3.16)$$

that is,

$$\frac{z^\Delta(t)}{z^{1/q}(t)} \leq r_1(t) \int_0^t r_2(t_2) \cdots \int_0^{t_{n-2}} r_{n-1}(t_{n-1}) \int_0^{t_{n-1}} r(t_n)\Delta t_n \Delta t_{n-1} \cdots \Delta t_2. \quad (3.17)$$

For  $z^\Delta(t) \geq 0$ , from the chain rule in [8, Theorem 1.90],

$$\begin{aligned} \left(\frac{1}{1-1/q}z^{-1/q+1}\right)^\Delta &= z^\Delta \int_0^1 \left(z + h\mu z^\Delta\right)^{-1/q} dh \\ &= z^{-1/q}z^\Delta \int_0^1 \left(1 + h\mu \frac{z^\Delta}{z}\right)^{-1/q} dh \\ &\leq z^{-1/q}z^\Delta. \end{aligned} \quad (3.18)$$

Letting  $t = t_1$  in (3.17) and integrating with respect to  $t_1$  from 0 to  $t$ , we have

$$\begin{aligned} & \frac{q}{q-1} \left( (z(t))^{(q-1)/q} - (z(0))^{(q-1)/q} \right) \\ & \leq \int_0^t \frac{z^\Delta(t_1)}{z^{1/q}(t_1)} \Delta t_1 \\ & \leq \int_0^t r_1(t_1) \int_0^{t_1} r_2(t_2) \cdots \int_0^{t_{n-2}} r_{n-1}(t_{n-1}) \int_0^{t_{n-1}} r(t_n) \Delta t_n \Delta t_{n-1} \cdots \Delta t_2 \Delta t_1, \end{aligned} \tag{3.19}$$

which means that

$$F(t) \leq z^{1/q}(t) \leq \left\{ c^{(q-1)/q} + \frac{q-1}{q} A[t, r_1, r_2, \dots, r_{n-1}, r] \right\}^{1/(q-1)}. \tag{3.20}$$

This completes the proof. □

*Remark 3.2.* Theorem 3.1 also holds for  $c = 0$ . To show this, assume (3.4) holds for  $c = 0$ , that is,

$$F^q(t) \leq A[t, r_1, \dots, r_{n-1}, rF] \quad \forall t \in \mathbb{T}_0^+. \tag{3.21}$$

Now, let  $d > 0$  be arbitrary. Then

$$F^q(t) \leq d + A[t, r_1, \dots, r_{n-1}, rF] \quad \forall t \in \mathbb{T}_0^+, \tag{3.22}$$

that is, (3.4) holds for  $c = d$ . By Theorem 3.1, (3.5) also holds for  $c = d$ , that is,

$$F(t) \leq \left\{ d^{(q-1)/q} + \frac{q-1}{q} A[t, r_1, \dots, r_{n-1}, r] \right\}^{1/(q-1)} \quad \forall t \in \mathbb{T}_0^+. \tag{3.23}$$

Since (3.23) holds for arbitrary  $d > 0$ , we may let  $d \rightarrow 0^+$  in (3.23) to arrive at

$$F(t) \leq \left\{ \frac{q-1}{q} A[t, r_1, \dots, r_{n-1}, r] \right\}^{1/(q-1)} \quad \forall t \in \mathbb{T}_0^+, \tag{3.24}$$

that is, (3.5) holds for  $c = 0$ .

**Theorem 3.3.** Let  $u, v$ , and  $h_j$  for  $j = 1, 2, 3, 4$  be real-valued nonnegative rd-continuous functions on  $t \in \mathbb{T}_0^+$  and let  $q > 1$  be a constant. If  $c_1, c_2$ , and  $\alpha$  are nonnegative constants such that

$$u^q(t) \leq c_1 + A[t, r_1, \dots, r_{n-1}, h_1 u] + A[t, r_1, \dots, r_{n-1}, h_2 \bar{v}] \quad \forall t \in \mathbb{T}_0^+, \tag{3.25}$$

$$v^q(t) \leq c_2 + A[t, r_1, \dots, r_{n-1}, h_3 \bar{u}] + A[t, r_1, \dots, r_{n-1}, h_4 v] \quad \forall t \in \mathbb{T}_0^+, \tag{3.26}$$



where  $\bar{u} = e_{\ominus\alpha}^q(\cdot, 0)u$  and  $\bar{v} = e_{\ominus\alpha}^q(\cdot, 0)v$ , then for all  $t \in \mathbb{T}_0^+$ ,

$$\begin{aligned} u(t) &\leq e_{\alpha}(t, 0) \left\{ [2^{q-1}(c_1 + c_2)]^{(q-1)/q} + \frac{q-1}{q} A[t, r_1, \dots, r_{n-1}, 2^{q-1}h] \right\}^{1/(q-1)}, \\ v(t) &\leq \left\{ [2^{q-1}(c_1 + c_2)]^{(q-1)/q} + \frac{q-1}{q} A[t, r_1, \dots, r_{n-1}, 2^{q-1}h] \right\}^{1/(q-1)}, \end{aligned} \quad (3.27)$$

where  $h(t) = \max\{h_1(t) + h_3(t), h_2(t) + h_4(t)\}$ .

*Proof.* Multiplying (3.25) by  $e_{\ominus\alpha}^q(t, 0)$  yields

$$\begin{aligned} e_{\ominus\alpha}^q(t, 0)u^q(t) &\leq c_1 e_{\ominus\alpha}^q(t, 0) + A[t, r_1, \dots, r_{n-1}, h_1 u] e_{\ominus\alpha}^q(t, 0) + A[t, r_1, \dots, r_{n-1}, h_2 \bar{v}] e_{\ominus\alpha}^q(t, 0) \\ &\leq c_1 + A[t, r_1, \dots, r_{n-1}, h_1 \bar{u}] + A[t, r_1, \dots, r_{n-1}, h_2 v]. \end{aligned} \quad (3.28)$$

Define

$$F(t) = e_{\ominus\alpha}(t, 0)u(t) + v(t). \quad (3.29)$$

By taking the  $q$ th power on both sides of (3.29) and using the elementary inequality  $(d_1 + d_2)^q \leq 2^{q-1}(d_1^q + d_2^q)$ , where  $d_1, d_2$  are nonnegative reals, and also noticing (3.26) and  $e_{\ominus\alpha}(t, 0) \leq 1$ , we get

$$\begin{aligned} F^q(t) &\leq 2^{q-1} \left[ e_{\ominus\alpha}^q(t, 0)u^q(t) + v^q(t) \right] \\ &\leq 2^{q-1} \{ c_1 + A[t, r_1, \dots, r_{n-1}, h_1 \bar{u}] + A[t, r_1, \dots, r_{n-1}, h_2 v] \\ &\quad + c_2 + A[t, r_1, \dots, r_{n-1}, h_3 \bar{u}] + A[t, r_1, \dots, r_{n-1}, h_4 v] \} \\ &= 2^{q-1} \{ (c_1 + c_2) + A[t, r_1, \dots, r_{n-1}, (h_1 + h_3) \bar{u}] + A[t, r_1, \dots, r_{n-1}, (h_2 + h_4) v] \} \\ &\leq 2^{q-1} (c_1 + c_2) + A[t, r_1, \dots, r_{n-1}, 2^{q-1}hF]. \end{aligned} \quad (3.30)$$

Now, Theorem 3.1 yields

$$F(t) \leq \left\{ [2^{q-1}(c_1 + c_2)]^{(q-1)/q} + \frac{q-1}{q} A[t, r_1, \dots, r_{n-1}, 2^{q-1}h] \right\}^{1/(q-1)}. \quad (3.31)$$

Noticing that (3.29) implies  $v \leq F$  and  $u \leq e_{\alpha}(\cdot, 0)F$ , the bounds in (3.27) follow, which concludes the proof.  $\square$

**Theorem 3.4.** Let  $q > 1$  and  $B$  be the set of all nonnegative real-valued rd-continuous functions defined on  $[0, t] \cap \mathbb{T}$ . Let  $K$  and  $L$  be monotone increasing linear operators on  $B$ . If there exists a

positive constant  $c$  such that, for  $y \in B$ ,

$$y^q(t) \leq c + \int_0^t \{qL[y^q](\tau) + K[y](\tau)\} \Delta\tau \quad \forall t \in \mathbb{T}_0^+, \quad (3.32)$$

then, for all  $t \in \mathbb{T}_0^+$ ,

$$y(t) \leq e_{\frac{1}{q\bar{L}}}(t, 0) \left\{ c^{(q-1)/q} + \frac{q-1}{q} \int_0^t (1 + \mu(\tau)(\ominus(q\bar{L}))(\tau)) \bar{K}(\tau) e_{\frac{1}{q\bar{L}}}^{1/q-1}(\tau, 0) \Delta\tau \right\}^{1/(q-1)}, \quad (3.33)$$

where  $\bar{L} = L[id]$ ,  $\bar{K} = K[id]$  with  $id(s) \equiv 1$  for all  $s \in \mathbb{T}$ .

*Proof.* Let

$$z(t) = c + \int_0^t \{qL[y^q](\tau) + K[y](\tau)\} \Delta\tau. \quad (3.34)$$

Hence,  $z(s) \leq z(t)$  for all  $0 \leq s \leq t$ , so that  $z \leq z(t)id$  on  $[0, t]$ , and thus

$$L[z] \leq L[z(t)id] = z(t)L[id] = z(t)\bar{L}. \quad (3.35)$$

Hence  $L[z](t) \leq z(t)\bar{L}(t)$ , and therefore  $L[z] \leq z\bar{L}$ . Similarly,  $K[z^{1/q}] \leq z^{1/q}\bar{K}$ . Using this and (3.32), we obtain that

$$z^\Delta = qL[y^q] + K[y] \leq qL[z] + K[z^{1/q}] \leq q\bar{L}z + \bar{K}z^{1/q}. \quad (3.36)$$

By the product rule [8, Theorem 1.20], we have

$$\begin{aligned} (e_{\ominus(q\bar{L})}(\cdot, 0)z)^\Delta &= \ominus(q\bar{L})e_{\ominus(q\bar{L})}(\cdot, 0)z + e_{\ominus(q\bar{L})}^\sigma(\cdot, 0)z^\Delta \\ &= \ominus(q\bar{L})e_{\ominus(q\bar{L})}(\cdot, 0)z + (1 + \mu(\ominus(q\bar{L})))e_{\ominus(q\bar{L})}(\cdot, 0)z^\Delta \\ &= e_{\ominus(q\bar{L})}(\cdot, 0)(\ominus(q\bar{L})z + (1 + \mu(\ominus(q\bar{L})))z^\Delta) \\ &\leq e_{\ominus(q\bar{L})}(\cdot, 0)(\ominus(q\bar{L})z + (1 + \mu(\ominus(q\bar{L}))))(q\bar{L}z + \bar{K}z^{1/q}) \\ &= e_{\ominus(q\bar{L})}(\cdot, 0) \left( \frac{-q\bar{L}}{1 + \mu q\bar{L}} z + \left( 1 + \mu \frac{-q\bar{L}}{1 + \mu q\bar{L}} \right) (q\bar{L}z + \bar{K}z^{1/q}) \right) \\ &= e_{\ominus(q\bar{L})}(\cdot, 0) \left( \frac{-q\bar{L}}{1 + \mu q\bar{L}} z + \frac{q\bar{L}}{1 + \mu q\bar{L}} z + (1 + \mu(\ominus(q\bar{L})))\bar{K}z^{1/q} \right) \\ &= e_{\ominus(q\bar{L})}(\cdot, 0)(1 + \mu(\ominus(q\bar{L})))\bar{K}z^{1/q}. \end{aligned} \quad (3.37)$$

In summary,

$$w^\Delta \leq \left(1 + \mu\left(\ominus(q\bar{L})\right)\right) \bar{K} w^{1/q} e_{q\bar{L}}^{1/q-1}(\cdot, 0), \quad \text{where } w = \frac{z}{e_{q\bar{L}}(\cdot, 0)}. \quad (3.38)$$

Obviously

$$ww^\sigma > 0, \quad \text{which implies } \frac{w^\Delta}{w} \in \mathcal{R}^+, \quad (3.39)$$

so that the chain rule [9, Theorem 2.37] yields

$$\left(\frac{1}{1-1/q} w^{-1/q+1}\right)^\Delta = w^{-1/q} w^\Delta \int_0^1 \left(1 + h\mu \frac{w^\Delta}{w}\right)^{-1/q} dh \leq w^{-1/q} w^\Delta. \quad (3.40)$$

Dividing both sides of (3.38) by  $w^{1/q}$  provides that

$$w^{-1/q} w^\Delta \leq \left(1 + \mu\left(\ominus(q\bar{L})\right)\right) \bar{K} e_{q\bar{L}}^{1/q-1}(\cdot, 0). \quad (3.41)$$

Integrating both sides of (3.41) from 0 to  $t$  and noticing (3.40), we find that

$$\frac{q}{q-1} \left(w^{1-1/q}(t) - w^{1-1/q}(0)\right) \leq \int_0^t \left(1 + \mu(\tau)\left(\ominus(q\bar{L})\right)(\tau)\right) \bar{K} e_{q\bar{L}}^{1/q-1}(\tau, 0) \Delta\tau. \quad (3.42)$$

Substitute the expression of  $w(t)$ , we have

$$\frac{z(t)}{e_{q\bar{L}}(t, 0)} \leq \left\{ c^{(q-1)/q} + \frac{q-1}{q} \int_0^t \left(1 + \mu(\tau)\left(\ominus(q\bar{L})\right)(\tau)\right) \bar{K} e_{q\bar{L}}^{1/q-1}(\tau, 0) \Delta\tau \right\}^{q/(q-1)}, \quad (3.43)$$

which gives the desired inequality (3.32). This concludes the proof.  $\square$

*Remark 3.5.* As in the discussion in Remark 3.2, Theorem 3.4 also holds true for  $c = 0$ .

#### 4. Some Applications

In this section, we indicate some applications of our results to obtain the estimates of the solutions of certain integral equations for which inequalities obtained in the literature thus far do not apply directly. As an application of Theorem 2.2, we consider the second-order dynamic equation

$$y^{\Delta\Delta} + p^\sigma(t)(y + y^\sigma) = 0. \quad (4.1)$$

**Theorem 4.1.** Assume that  $p$  is a differentiable positive function such that  $p^\Delta$  is rd-continuous. If there exist  $t_0 \in \mathbb{T}$  and  $M > 0$  such that

$$\frac{1}{\sqrt{p(t)}} e_{|p^\Delta|/2p}(t, t_0) \leq M \quad \forall t \in \mathbb{T}_{t_0}^+, \quad (4.2)$$

then all nonoscillatory solutions of (4.1) are bounded.

*Proof.* Let  $y$  be a nonoscillatory solution of (4.1). Without loss of generality, we assume there exists  $t_0 \in \mathbb{T}$  such that

$$y(t) > 0 \quad \forall t \in \mathbb{T}_{t_0}^+. \quad (4.3)$$

Then

$$y^{\Delta\Delta}(t) = -p^\sigma(t)(y(t) + y^\sigma(t)) < 0 \quad \forall t \in \mathbb{T}_{t_0}^+. \quad (4.4)$$

Hence,  $y^\Delta$  is strictly decreasing on  $\mathbb{T}_{t_0}^+$ . Thus, either

$$y^\Delta(t) > 0 \quad \forall t \in \mathbb{T}_{t_0}^+ \quad (4.5)$$

or there exists  $t_1 \in \mathbb{T}_{t_0}^+$  such that

$$y^\Delta(t) < 0 \quad \forall t \in \mathbb{T}_{t_1}^+. \quad (4.6)$$

We now claim that (4.6) is impossible to hold. To show this, let us assume that (4.6) is true. Then  $y$  is strictly decreasing on  $\mathbb{T}_{t_1}^+$  and

$$y(t) = y(t_1) + \int_{t_1}^t y^\Delta(\tau) \Delta\tau \leq y(t_1) + y^\Delta(t_1)(t - t_1) \quad \forall t \in \mathbb{T}_{t_1}^+. \quad (4.7)$$

Hence, there exists  $t_2 \in \mathbb{T}_{t_1}^+$  such that

$$y(t) < 0 \quad \forall t \in \mathbb{T}_{t_2}^+, \quad (4.8)$$

contradicting  $y(t) > 0$  for all  $t \in \mathbb{T}_{t_0}^+$ . Similarly, we can prove that if  $y(t) < 0$ , then  $y^{\Delta\Delta}(t) > 0$  and  $y^\Delta(t) \leq 0$  for  $t \in \mathbb{T}_{t_1}^+$ .

Multiplying (4.1) on both sides by  $y^\Delta$  and taking integral from  $t_1$  to  $t$ , we have

$$\int_{t_1}^t y^\Delta(\tau) y^{\Delta\Delta}(\tau) \Delta\tau + \int_{t_1}^t p^\sigma(\tau)(y(\tau) + y^\sigma(\tau)) y^\Delta(\tau) \Delta\tau = 0. \quad (4.9)$$

From the integration by parts in [8, Theorem 1.77],

$$\left(y^\Delta(t)\right)^2 - \left(y^\Delta(t_1)\right)^2 - \int_{t_1}^t y^{\Delta\Delta}(\tau)y^{\Delta\sigma}(\tau)\Delta\tau + p(t)y^2(t) - p(t_1)y^2(t_1) - \int_{t_1}^t p^\Delta(\tau)y^2(\tau)\Delta\tau = 0, \quad (4.10)$$

Thus, with  $c_1 = p(t_1)y^2(t_1) + (y^\Delta(t_1))^2 > 0$ , we have

$$\left(\sqrt{p(t)}y(t)\right)^2 \leq c_1 + \int_{t_1}^t \frac{|p^\Delta(\tau)||y(\tau)|}{\sqrt{p(\tau)}}\sqrt{p(\tau)}|y(\tau)|\Delta\tau \quad \forall t \in \mathbb{T}_{t_1}^+. \quad (4.11)$$

Theorem 2.2 gives that

$$\left|\sqrt{p(t)}y(t)\right| \leq \sqrt{c_1} + \frac{1}{2} \int_{t_1}^t \frac{|p^\Delta(\tau)||y(\tau)|}{\sqrt{p(\tau)}}\Delta\tau = \sqrt{c_1} + \int_{t_1}^t \frac{|p^\Delta(\tau)|}{2p(\tau)}\left|\sqrt{p(\tau)}y(\tau)\right|\Delta\tau \quad \forall t \in \mathbb{T}_{t_1}^+. \quad (4.12)$$

Applying Gronwall's inequality from Lemma 2.1 yields

$$\left|\sqrt{p(t)}y(t)\right| \leq \sqrt{c_1}e_{|p^\Delta|/2p}(t, t_1) \quad \forall t \in \mathbb{T}_{t_1}^+. \quad (4.13)$$

Hence,

$$|y(t)| \leq \sqrt{c_1} \frac{1}{\sqrt{p(t)}}e_{|p^\Delta|/2p}(t, t_1) \leq \sqrt{c_1}M \quad \forall t \in \mathbb{T}_{t_1}^+, \quad (4.14)$$

which completes the proof.  $\square$

The proof in Theorem 4.1 corrects an inaccuracy in the proof of [1, Theorem 1]. We can also obtain the following results.

**Corollary 4.2.** *Let  $\mathbb{T} = \mathbb{R}$ . If  $p$  is a continuously differentiable positive function such that  $p'$  is nonnegative, then all nonoscillatory solutions of (4.1) are bounded.*

*Proof.* For  $\mathbb{T} = \mathbb{R}$ , we have

$$\begin{aligned} \frac{1}{\sqrt{p(t)}} e_{|p^\Delta|/2p}(t, 0) &= \frac{1}{\sqrt{p(t)}} e^{\int_0^t (p'(\tau)/2p(\tau)) d\tau} \\ &= \frac{1}{\sqrt{p(t)}} e^{(1/2) \ln(p(t)/p(0))} \\ &= \frac{1}{\sqrt{p(t)}} \left( \frac{p(t)}{p(0)} \right)^{1/2} \\ &= \frac{1}{\sqrt{p(0)}}, \end{aligned} \quad (4.15)$$

and hence the statement follows from Theorem 4.1.  $\square$

*Example 4.3.* Consider the nonlinear one-dimensional integral equation of the form

$$u^q(t) = f(t) + \int_0^t k(t, s) g(s, u(s)) \Delta s, \quad (4.16)$$

where  $f : \mathbb{T}_0^+ \rightarrow \mathbb{R}$ ,  $k : \mathbb{T}_0^+ \times \mathbb{T}_0^+ \rightarrow \mathbb{R}$ ,  $g : \mathbb{T}_0^+ \times \mathbb{R} \rightarrow \mathbb{R}$  are rd-continuous functions, and  $q > 1$  is a constant. When  $\mathbb{T} = \mathbb{R}$ , its physical meaning is to model the water percolation phenomena, and Okrański has studied the existence and uniqueness of solutions [14].

Here, we assume that every solution  $u$  of (4.16) exists on the interval  $\mathbb{T}_0^+$ . We suppose that the functions  $f, k, g$  in (4.16) satisfy the conditions

$$|f(t)| \leq c_1, \quad |k(t, s)| \leq c_2, \quad |g(t, u)| \leq r(t)|u|, \quad (4.17)$$

where  $c_1, c_2$  are nonnegative constants and  $r : [0, \infty) \cap \mathbb{T} \rightarrow \mathbb{R}_+$  is an rd-continuous function. From (4.16) and using (4.17), it is easy to observe that

$$|u(t)|^q \leq c_1 + \int_0^t c_2 r(s) |u(s)| \Delta s. \quad (4.18)$$

Now an application of Theorem 3.1 with  $n = 1$  gives

$$|u(t)| \leq \left\{ c_1^{(q-1)/q} + \frac{q-1}{q} \int_0^t c_2 r(s) \Delta s \right\}^{1/(q-1)}, \quad (4.19)$$

which gives the bound on  $u$ .

Now, we consider (4.16) under the conditions

$$|f(t)| \leq c_1 e_{\ominus\alpha}^q(t, 0), \quad |k(t, s)| \leq h(s) e_{\ominus\alpha}^q(t, 0), \quad |g(t, u)| \leq r(t)|u|, \quad (4.20)$$

where  $c_1$  and  $r$  are as above,  $\alpha > 0$  is a constant,  $h : \mathbb{T}_0^+ \rightarrow \mathbb{R}_+$  is an rd-continuous function, and

$$\int_0^\infty h(s)r(s)e_{\ominus\alpha}(s,0)\Delta s < \infty. \quad (4.21)$$

From (4.16) and (4.20), it is easy to observe that

$$|e_\alpha(t,0)u(t)|^q \leq c_1 + \int_0^t h(s)r(s)e_{\ominus\alpha}(s,0)|e_\alpha(s,0)u(s)|\Delta s. \quad (4.22)$$

Applying Theorem 3.1 with  $n = 1$  yields

$$e_\alpha(t,0)|u(t)| \leq \left\{ c_1^{(q-1)/q} + \frac{q-1}{q} \int_0^t h(s)r(s)e_{\ominus\alpha}(s,0)\Delta s \right\}^{1/(q-1)}. \quad (4.23)$$

So,

$$|u(t)| \leq c^* e_{\ominus\alpha}(t,0), \quad \text{where } c^* = c_1^{(q-1)/q} + \frac{q-1}{q} \int_0^\infty h(s)r(s)e_{\ominus\alpha}(s,0)\Delta s > 0. \quad (4.24)$$

From (4.24), we see that the solution  $u(t)$  of (4.16) approaches zero as  $t \rightarrow \infty$ .

## Acknowledgments

This work is supported by Grants 60673151 and 10571183 from NNSF of China, and by Grant 08JA910003 from Humanities and Social Sciences in Chinese Universities.

## References

- [1] O. Yang-Liang, "The boundedness of solutions of linear differential equations  $y' + A(t)y = 0$ ," *Advances in Mathematics*, vol. 3, pp. 409–415, 1957.
- [2] E. Yang, "On some nonlinear integral and discrete inequalities related to Ou-Iang's inequality," *Acta Mathematica Sinica*, vol. 14, no. 3, pp. 353–360, 1998.
- [3] B. G. Pachpatte, "On a certain inequality arising in the theory of differential equations," *Journal of Mathematical Analysis and Applications*, vol. 182, no. 1, pp. 143–157, 1994.
- [4] P. Y. H. Pang and R. P. Agarwal, "On an integral inequality and its discrete analogue," *Journal of Mathematical Analysis and Applications*, vol. 194, no. 2, pp. 569–577, 1995.
- [5] Y. J. Cho, Y.-H. Kim, and J. Pečarić, "New Gronwall-Ou-Iang type integral inequalities and their applications," *The ANZIAM Journal*, vol. 50, no. 1, pp. 111–127, 2008.
- [6] W.-S. Cheung and Q.-H. Ma, "On certain new Gronwall-Ou-Iang type integral inequalities in two variables and their applications," *Journal of Inequalities and Applications*, no. 4, pp. 347–361, 2005.
- [7] S. Hilger, "Analysis on measure chains—a unified approach to continuous and discrete calculus," *Results in Mathematics*, vol. 18, no. 1-2, pp. 18–56, 1990.
- [8] M. Bohner and A. Peterson, *Dynamic Equations on Time Scales: An Introduction with Application*, Birkhäuser, Boston, Mass, USA, 2001.
- [9] M. Bohner and A. Peterson, Eds., *Advances in Dynamic Equations on Time Scales*, Birkhäuser, Boston, Mass, USA, 2003.

- [10] W. N. Li and W. Sheng, "Some nonlinear dynamic inequalities on time scales," *Proceedings of Indian Academy of Sciences. Mathematical Sciences*, vol. 117, no. 4, pp. 545–554, 2007.
- [11] E. Akin-Bohner, M. Bohner, and F. Akin, "Pachpatte inequalities on time scales," *Journal of Inequalities in Pure and Applied Mathematics*, vol. 6, no. 1, article 6, 23 pages, 2005.
- [12] W. N. Li, "Bounds for certain new integral inequalities on time scales," *Advances in Difference Equations*, vol. 2009, Article ID 484185, 16 pages, 2009.
- [13] C. M. Dafermos, "The second law of thermodynamics and stability," *Archive for Rational Mechanics and Analysis*, vol. 70, no. 2, pp. 167–179, 1979.
- [14] W. Okrasinski, "On a nonlinear convolution equation occurring in the theory of water percolation," *Annales Polonici Mathematici*, vol. 37, no. 3, pp. 223–229, 1980.