

01 Jan 2007

## On a Sub-supersolution Method for the Prescribed Mean Curvature Problem

Vy Khoi Le

Missouri University of Science and Technology, [vy@mst.edu](mailto:vy@mst.edu)

Follow this and additional works at: [https://scholarsmine.mst.edu/math\\_stat\\_facwork](https://scholarsmine.mst.edu/math_stat_facwork)



Part of the [Mathematics Commons](#), and the [Statistics and Probability Commons](#)

---

### Recommended Citation

V. K. Le, "On a Sub-supersolution Method for the Prescribed Mean Curvature Problem," *Czechoslovak Mathematical Journal*, Springer Verlag, Jan 2007.

The definitive version is available at <https://doi.org/10.1007/s10587-008-0034-7>

This Article - Journal is brought to you for free and open access by Scholars' Mine. It has been accepted for inclusion in Mathematics and Statistics Faculty Research & Creative Works by an authorized administrator of Scholars' Mine. This work is protected by U. S. Copyright Law. Unauthorized use including reproduction for redistribution requires the permission of the copyright holder. For more information, please contact [scholarsmine@mst.edu](mailto:scholarsmine@mst.edu).

## SUB-SUPERSOLUTION THEOREMS FOR QUASILINEAR ELLIPTIC PROBLEMS: A VARIATIONAL APPROACH

VY KHOI LE, KLAUS SCHMITT

*Dedicated to Hans Knobloch with much admiration and appreciation*

ABSTRACT. This paper presents a variational approach to obtain sub - supersolution theorems for a certain type of boundary value problem for a class of quasilinear elliptic partial differential equations. In the case of semilinear ordinary differential equations results of this type were first proved by Hans Knobloch in the early sixties using methods developed by Cesari.

### 1. INTRODUCTION - PROBLEM SETTING

We are interested here in a variational sub-supersolution approach to a quasilinear elliptic boundary value problem which, in the one-space dimensional and semilinear case, is a boundary value problem for a second order scalar ordinary differential equation subject to periodic boundary conditions. The latter problem was first studied by Hans Knobloch [5] and later by many other authors using various kinds of nonlinear analysis methods (see, e.g., [10, 9, 6, 2]). The present paper is a continuation of and extends the results of the recent note [11]. In the case of boundary value problems subject to Dirichlet boundary conditions, sub-supersolution results, in a variational setting, were first obtained by Peter Hess [3]. We here follow closely the approach used in [7], where Dirichlet boundary value problems for degenerate elliptic equations were studied.

Let  $\Omega \subset \mathbb{R}^N$  be a bounded domain with smooth boundary. We consider the following boundary value problem:

$$-\operatorname{div}[A(x, \nabla u)] + f(x, u) = 0, \quad x \in \Omega, \quad (1.1)$$

$$u(x) = \text{constant}, \quad x \in \partial\Omega, \quad (1.2)$$

$$\int_{\partial\Omega} A(x, \nabla u) \cdot n, S = 0. \quad (1.3)$$

(Note that in condition (1.2) it is understood that the trace of  $u$  is a constant function, with the constant not being fixed.) Here,  $A : \Omega \times \mathbb{R}^N \rightarrow \mathbb{R}^N$  is a Carathéodory function satisfying the following conditions:

---

2000 *Mathematics Subject Classification.* 35B45, 35J65, 35J60.

*Key words and phrases.* Sub and supersolutions; periodic solutions; variational approach.

©2004 Texas State University - San Marcos.

Submitted July 28, 2004. Published October 7, 2004.

- There exist  $p \in (1, \infty)$ ,  $a_1 \in L^{p'}(\Omega)$  ( $p'$  is the conjugate of  $p$ ), and  $b_1 > 0$  such that

$$|A(x, \xi)| \leq a_1(x) + b_1|\xi|^{p-1}, \quad (1.4)$$

for a.e.  $x \in \Omega$ , all  $\xi \in \mathbb{R}^N$ .

- $A(x, \xi)$  is monotone in  $\xi$ , that is

$$[A(x, \xi) - A(x, \xi')] \cdot (\xi - \xi') \geq 0, \quad \text{for a.e. } x \in \Omega, \text{ all } \xi, \xi' \in \mathbb{R}^N. \quad (1.5)$$

- $A$  has the following coercivity property: There exist  $a_2 \in L^1(\Omega)$  and  $b_2 > 0$  such that

$$A(x, \xi) \cdot \xi \geq b_2|\xi|^p - a_2(x), \quad \text{for a.e. } x \in \Omega, \text{ all } \xi \in \mathbb{R}^N. \quad (1.6)$$

**Remark 1.1.** (a) When  $N = 1$  and  $\Omega = (a, b)$ , the boundary condition (1.2)-(1.3) becomes the boundary condition on  $(a, b)$ :

$$u(a) = u(b), \quad A(a, u'(a)) = A(b, u'(b)),$$

which, when  $A(x, v) = v$  is the usual set of periodic boundary conditions

$$u(a) = u(b), \quad u'(a) = u'(b).$$

(b) An example of the operator  $A$  above is the  $p$ -Laplacian, i.e.,

$$A(x, \nabla u) = |\nabla u|^{p-2} \nabla u, \quad p > 1.$$

It is easy to check that  $A$  satisfies conditions (1.4), (1.5), and (1.6) above. In this case, the boundary condition (1.3) becomes

$$\int_{\partial\Omega} |\nabla u|^{p-2} \frac{\partial u}{\partial n} dS = 0.$$

Assume that  $f : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$  is a Carathéodory function with some appropriate growth condition to be specified later. We denote by  $W^{1,p}(\Omega)$  the usual Sobolev space, equipped with the norm

$$\|u\| = \|u\|_{W^{1,p}(\Omega)} = \left( \|u\|_{L^p(\Omega)}^p + \|\nabla u\|_{L^p(\Omega)}^p \right)^{1/p}. \quad (1.7)$$

Let  $\mathcal{A}, F : W^{1,p}(\Omega) \rightarrow [W^{1,p}(\Omega)]^*$  be defined by

$$\langle Fu, v \rangle = \int_{\Omega} f(x, u)v dx,$$

and

$$\langle \mathcal{A}u, v \rangle = \int_{\Omega} A(x, \nabla u) \cdot \nabla v dx, \quad \forall u, v \in W^{1,p}(\Omega).$$

From (1.4)-(1.6), we see that  $\mathcal{A}$  is continuous, bounded, monotone, and coercive in the following sense:

$$\langle \mathcal{A}u, u \rangle \geq b_2 \|\nabla u\|_{L^p(\Omega)}^p - \|a_2\|_{L^1(\Omega)}, \quad \forall u \in W^{1,p}(\Omega). \quad (1.8)$$

Let

$$V_c = \{u \in W^{1,p}(\Omega) : u|_{\partial\Omega} = \text{constant}\}.$$

Then  $V_c$  is a closed subspace of  $W^{1,p}(\Omega)$  and thus a reflexive Banach space with the restricted norm of (1.7). The weak (variational) formulation of the boundary value problem (1.1)-(1.3) is the following variational equality:

$$\int_{\Omega} A(x, \nabla u) \cdot \nabla v dx + \int_{\Omega} f(x, u)v dx = 0, \quad \forall v \in V_c \quad (1.9)$$

$$u \in V_c.$$

To check this, note that if  $u$  satisfies (1.1)-(1.3) and  $v \in V_c$  then

$$\begin{aligned} 0 &= - \int_{\Omega} \operatorname{div} A(x, \nabla u) v \, dx + \int_{\Omega} f(x, u) v \, dx \\ &= \int_{\Omega} A(x, \nabla u) \cdot \nabla v \, dx - (v|_{\partial\Omega}) \int_{\partial\Omega} A(x, \nabla u) \cdot n \, dS + \int_{\Omega} f(x, u) v \, dx \\ &= \int_{\Omega} A(x, \nabla u) \cdot \nabla v \, dx + \int_{\Omega} f(x, u) v \, dx. \end{aligned}$$

Hence, we have (1.9). Conversely, if  $u \in V_c$  is a solution of (1.9) then by choosing  $v \in C_0^\infty(\Omega) \subset V_c$  in (1.9) and applying the divergence theorem as above, we see that (1.1) holds. Choosing  $v = 1$  in (1.9), we have  $\int_{\Omega} f(x, u) \, dx = 0$ . On the other hand, integrating (1.1) over  $\Omega$  and using once more the Divergence theorem yield

$$0 = - \int_{\Omega} \operatorname{div} A(x, \nabla u) \, dx + \int_{\Omega} f(x, u) \, dx = - \int_{\partial\Omega} A(x, \nabla u) \cdot n \, dS.$$

Hence, we have the boundary condition (1.3).

## 2. SUB-SUPERSOLUTIONS

We shall study the existence of solutions of (1.9) by first defining appropriate concepts of sub- and supersolutions.

**Definition 2.1.** A function  $\underline{u}$  (resp.  $\bar{u}$ ) in  $V_c$  is called a subsolution (resp. supersolution) of (1.9) if

$$\int_{\Omega} A(x, \nabla \underline{u}) \cdot \nabla v \, dx + \int_{\Omega} f(x, \underline{u}) v \, dx \leq 0 \quad (\text{resp. } \geq 0), \quad (2.1)$$

for all  $v \in V_c$ ,  $v \geq 0$  a.e. in  $\Omega$ .

**Remark 2.2.** When  $A$  is the Laplacian, i.e.,  $A(x, \nabla u) = \nabla u$  and  $p = 2$ , or when  $N = 1$  (ODE case), the above definition of sub- and supersolutions is the variational form of that given in [11], without imposing additional smoothness assumptions.

As is the case with solutions satisfying additional smoothness conditions, sub- and supersolutions, when smooth enough, satisfy additional boundary conditions. Let us see this in the case of the  $p$ -Laplacian. For assume that  $\alpha \in V_c \cap W^{2,p}(\Omega)$  satisfies (cf. (17) of [11]):

$$\int_{\Omega} |\nabla \alpha|^{p-2} \nabla \alpha \cdot \nabla \phi \, dx + \int_{\Omega} f(x, \alpha) \phi \, dx \leq 0, \quad \forall \phi \in C_0^\infty(\Omega), \phi \geq 0, \quad (2.2)$$

and

$$\int_{\partial\Omega} |\nabla \alpha|^{p-2} \nabla \alpha \cdot n \, dS \leq 0. \quad (2.3)$$

Since  $\alpha \in W^{2,p}(\Omega)$ , Green's theorem (or the Divergence theorem) implies that

$$\int_{\Omega} [-\operatorname{div} (|\nabla \alpha|^{p-2} \nabla \alpha) + f(x, \alpha)] \phi \, dx \leq 0, \quad \forall \phi \in C_0^\infty(\Omega), \phi \geq 0,$$

i.e., (in the sense of distributions),

$$-\operatorname{div} (|\nabla \alpha|^{p-2} \nabla \alpha) + f(x, \alpha) \leq 0 \quad \text{a.e. on } \Omega. \quad (2.4)$$

Let  $v \in V_c$ ,  $v \geq 0$  a.e. on  $\Omega$ . It follows from (2.4) that

$$\begin{aligned} 0 &\geq \int_{\Omega} [-\operatorname{div}(|\nabla\alpha|^{p-2}\nabla\alpha) + f(x, \alpha)]v \, dx \\ &= \int_{\Omega} |\nabla\alpha|^{p-2}\nabla\alpha \cdot \nabla v \, dx - \int_{\partial\Omega} |\nabla\alpha|^{p-2}\frac{\partial\alpha}{\partial\nu}v \, dS + \int_{\Omega} f(x, \alpha)v \, dx. \end{aligned}$$

Hence,

$$\int_{\Omega} |\nabla\alpha|^{p-2}\nabla\alpha \cdot \nabla v \, dx + \int_{\Omega} f(x, \alpha)v \, dx \leq (v|_{\partial\Omega}) \int_{\partial\Omega} |\nabla\alpha|^{p-2}\frac{\partial\alpha}{\partial\nu} \, dS \leq 0,$$

that is,  $\alpha$  satisfies (2.1). Conversely, assume  $\alpha \in V_c \cap W^{2,p}(\Omega)$  satisfies (2.1). Since  $C_0^\infty(\Omega) \subset V_c$ , we have (2.2). To prove that  $\alpha$  satisfies (2.3), we choose a sequence  $\{\Omega_n\}$  of subdomains of  $\Omega$  such that

$$\overline{\Omega_n} \subset \Omega_{n+1}, \quad \forall n, \quad \text{and} \quad \Omega = \bigcup_{n=1}^{\infty} \Omega_n. \quad (2.5)$$

For each  $n \in \mathbb{N}$ , choose  $\phi_n \in C_0^\infty(\Omega)$  such that  $0 \leq \phi_n(x) \leq 1$ ,  $\forall x \in \Omega$ , and  $\phi_n(x) = 1$ ,  $\forall x \in \Omega_n$ . Let  $v_n = 1 - \phi_n$  ( $n \in \mathbb{N}$ ). Then  $v_n \in V_c$ ,  $v_n = 1$  on  $\partial\Omega$ , and  $0 \leq v_n \leq 1$  on  $\Omega$ . Letting  $v = v_n$  in (2.1), one gets

$$\begin{aligned} 0 &\geq \int_{\Omega} |\nabla\alpha|^{p-2}\nabla\alpha \cdot \nabla v_n \, dx + \int_{\Omega} f(x, \alpha)v_n \, dx \\ &= \int_{\Omega} [-\operatorname{div}(|\nabla\alpha|^{p-2}\nabla\alpha) + f(x, \alpha)]v_n \, dx + \int_{\partial\Omega} |\nabla\alpha|^{p-2}\frac{\partial\alpha}{\partial\nu}v_n \, dS \\ &= \int_{\Omega} [-\operatorname{div}(|\nabla\alpha|^{p-2}\nabla\alpha) + f(x, \alpha)]v_n \, dx + \int_{\partial\Omega} |\nabla\alpha|^{p-2}\frac{\partial\alpha}{\partial\nu} \, dS. \end{aligned} \quad (2.6)$$

Because  $v_n = 0$  on  $\Omega_n$ , from (2.5) and the Dominated convergence theorem, one obtains

$$\lim_{n \rightarrow \infty} \int_{\Omega} [-\operatorname{div}(|\nabla\alpha|^{p-2}\nabla\alpha) + f(x, \alpha)]v_n \, dx = 0.$$

Letting  $n \rightarrow \infty$  in (2.6), we obtain  $\int_{\partial\Omega} |\nabla\alpha|^{p-2}\frac{\partial\alpha}{\partial\nu} \, dS \leq 0$ .

### 3. EXISTENCE RESULTS

Our main existence result is the following theorem.

**Theorem 3.1.** *Assume there exists a pair of sub- and supersolution  $\underline{u}$  and  $\bar{u}$  of (1.9) such that  $\underline{u} \leq \bar{u}$  and that  $f$  satisfies the following growth condition:*

$$|f(x, u)| \leq a_3(x), \quad (3.1)$$

for a.e.  $x \in \Omega$ , all  $u \in [\underline{u}(x), \bar{u}(x)]$ , with  $a_3 \in L^p(\Omega)$ . Then, (1.9) has a solution  $u \in V_c$  such that  $\underline{u} \leq u \leq \bar{u}$ .

*Proof.* We define

$$b(x, u) = \begin{cases} [u - \bar{u}(x)]^{p-1} & \text{if } u > \bar{u}(x) \\ 0 & \text{if } \underline{u}(x) \leq u \leq \bar{u}(x) \\ -[\underline{u}(x) - u]^{p-1} & \text{if } u < \underline{u}(x), \end{cases} \quad (3.2)$$

for  $x \in \Omega$ ,  $u \in \mathbb{R}$ , and

$$(Tu)(x) = \begin{cases} \bar{u}(x) & \text{if } u(x) > \bar{u}(x) \\ u(x) & \text{if } \underline{u}(x) \leq u(x) \leq \bar{u}(x) \\ \underline{u}(x) & \text{if } u(x) < \underline{u}(x), \end{cases} \quad (3.3)$$

for  $x \in \Omega$  and  $u \in W^{1,p}(\Omega)$ . Straightforward calculations show that

$$|b(x, u)| \leq a_4(x) + b_4|u|^{p-1},$$

for a.e.  $x \in \Omega$ , all  $u \in \mathbb{R}$ , where  $b_4 > 0$  and  $a_4 \in L^{p'}(\Omega)$ . Therefore, the operator  $B : W^{1,p}(\Omega) \rightarrow [W^{1,p}(\Omega)]^*$  given by

$$\langle Bu, v \rangle = \int_{\Omega} b(x, u)v \, dx \quad (u, v \in W^{1,p}(\Omega))$$

is well defined, completely continuous, and bounded. Moreover, there are  $a_5, b_5 > 0$  such that

$$\langle Bu, u \rangle \geq b_5 \|u\|_{L^p(\Omega)}^p - a_5, \quad \forall u \in W^{1,p}(\Omega). \quad (3.4)$$

Let us consider the following variational equality in  $V_c$ :

$$\begin{aligned} \langle \mathcal{A}u + Bu + F(Tu), v \rangle &= 0, \quad \forall v \in V_c \\ u &\in V_c. \end{aligned} \quad (3.5)$$

It follows from (3.1) that  $F \circ T$  is well defined and completely continuous from  $W^{1,p}(\Omega)$  to its dual space. Because  $\mathcal{A}$  is monotone,  $\mathcal{A} + B + F \circ T$  is pseudo-monotone. Next, let us show that  $\mathcal{A} + B + F \circ T$  is coercive on  $W^{1,p}(\Omega)$  in the following sense:

$$\lim_{\|u\| \rightarrow \infty} \frac{\langle \mathcal{A}u + Bu + F(Tu), u \rangle}{\|u\|} = \infty. \quad (3.6)$$

In fact, from (3.3) and (3.1),

$$|\langle F(Tu), u \rangle| = \left| \int_{\Omega} f(x, Tu)u \, dx \right| \leq \int_{\Omega} a_3|u| \, dx \leq \|a_3\|_{L^{p'}(\Omega)} \|u\|_{L^p(\Omega)}. \quad (3.7)$$

Combining (3.7) with (3.4) and (1.8), we get

$$\begin{aligned} &\langle \mathcal{A}u + Bu + F(Tu), u \rangle \\ &\geq b_2 \|\nabla u\|_{L^p(\Omega)}^p - \|a_2\|_{L^1(\Omega)} + b_5 \|u\|_{L^p(\Omega)}^p - a_5 - \|a_3\|_{L^{p'}(\Omega)} \|u\|_{L^p(\Omega)} \\ &\geq \min\{b_2, b_5\} (\|u\|_{L^p(\Omega)}^p + \|\nabla u\|_{L^p(\Omega)}^p) - \|a_3\|_{L^{p'}(\Omega)} \|u\| - \|a_2\|_{L^1(\Omega)} - a_5 \\ &= b_6 \|u\|^p - a_6 \|u\| - a_7, \quad \forall u \in W^{1,p}(\Omega), \end{aligned}$$

with  $a_6, a_7, b_6 > 0$ . Because  $p > 1$ , this estimate implies (3.6).

Since  $V_c$  is a closed subspace of  $W^{1,p}(\Omega)$ , the existence of solutions of (3.5) follows from classical existence theorems for elliptic variational inequalities (cf. e.g. [8]). Assume that  $u$  is any solution of (3.5). We prove that

$$\underline{u} \leq u \leq \bar{u} \quad \text{a.e. in } \Omega, \quad (3.8)$$

and thus  $u$  is also a solution of (1.9). Let us verify the first inequality in (3.8). Since  $u, \underline{u} \in W^{1,p}(\Omega)$ , we have  $(\underline{u} - u)^+ \in W^{1,p}(\Omega)$ . Moreover, since  $u|_{\partial\Omega}$  and  $\underline{u}|_{\partial\Omega}$  are constants,

$$[(\underline{u} - u)^+]|_{\partial\Omega} = (\underline{u}|_{\partial\Omega} - u|_{\partial\Omega})^+ = \text{constant},$$

i.e.,

$$(\underline{u} - u)^+ \in V_c. \quad (3.9)$$

Choosing  $v = (\underline{u} - u)^+$  in (3.5), one obtains

$$\int_{\Omega} A(x, \nabla u) \cdot \nabla[(\underline{u} - u)^+] dx + \int_{\Omega} [b(x, u) + f(Tu)](\underline{u} - u)^+ dx = 0. \quad (3.10)$$

On the other hand, letting  $v = (\underline{u} - u)^+(\geq 0)$  in (2.1) gives us

$$\int_{\Omega} A(x, \nabla \underline{u}) \cdot \nabla[(\underline{u} - u)^+] dx + \int_{\Omega} f(\underline{u})(\underline{u} - u)^+ dx \leq 0. \quad (3.11)$$

Subtracting (3.10) from (3.11) yields

$$\begin{aligned} & \int_{\Omega} [A(x, \nabla \underline{u}) - A(x, \nabla u)] \cdot \nabla[(\underline{u} - u)^+] dx + \int_{\Omega} [f(\underline{u}) - f(Tu)](\underline{u} - u)^+ dx \\ & \leq \int_{\Omega} b(x, u)(\underline{u} - u)^+ dx. \end{aligned} \quad (3.12)$$

Note that from (1.5) and Stampacchia's theorem (cf. e.g. [4, 1]), we have

$$\begin{aligned} & \int_{\Omega} [A(x, \nabla \underline{u}) - A(x, \nabla u)] \cdot \nabla[(\underline{u} - u)^+] dx \\ & = \int_{\{x \in \Omega: \underline{u}(x) > u(x)\}} [A(x, \nabla \underline{u}) - A(x, \nabla u)] \cdot (\nabla \underline{u} - \nabla u) dx \\ & \geq 0. \end{aligned} \quad (3.13)$$

From the definition of  $Tu$  in (3.3), we have  $Tu(x) = \underline{u}(x)$  on  $\{x \in \Omega : \underline{u}(x) > u(x)\}$  and thus

$$\int_{\Omega} [f(\underline{u}) - f(Tu)](\underline{u} - u)^+ dx = \int_{\{x \in \Omega: \underline{u}(x) > u(x)\}} [f(\underline{u}) - f(Tu)](\underline{u} - u) dx = 0. \quad (3.14)$$

Using (3.13) and (3.14) in (3.12), one obtains

$$0 \leq \int_{\Omega} b(x, u)(\underline{u} - u)^+ dx = - \int_{\{x \in \Omega: \underline{u}(x) > u(x)\}} (\underline{u} - u)^p dx \leq 0.$$

This implies that

$$\int_{\{x \in \Omega: \underline{u}(x) > u(x)\}} (\underline{u} - u)^p dx = 0,$$

i.e.,  $\underline{u} - u = 0$  a.e. on  $\{x \in \Omega : \underline{u}(x) > u(x)\}$ , or,  $\{x \in \Omega : \underline{u}(x) > u(x)\}$  has measure 0. This shows the first inequality in (3.8). The other inequality there is established in the same way. From (3.8) and (3.2)-(3.3), we immediately have  $b(x, u(x)) = 0$  and  $Tu(x) = u(x)$  for a.e.  $x \in \Omega$ . (3.5) thus becomes (1.9).  $\square$

**Remark 3.2.** By modifying the proof of Theorem 3.1 appropriately, we can extend that theorem to the existence of solutions of (1.9) between a finite number of sub- and supersolutions. In fact, we can show that if  $\underline{u}_1, \dots, \underline{u}_k$  (resp.  $\bar{u}_1, \dots, \bar{u}_m$ ) are subsolutions (resp. supersolutions) of (1.9) such that

$$\max\{\underline{u}_1, \dots, \underline{u}_k\} \leq \min\{\bar{u}_1, \dots, \bar{u}_m\},$$

and that  $f$  satisfies an appropriate growth condition between these sub- and supersolutions, then there exists a solution  $u$  of (1.9) such that  $\max\{\underline{u}_1, \dots, \underline{u}_k\} \leq u \leq \min\{\bar{u}_1, \dots, \bar{u}_m\}$  ( see, for example, [7] for more details).

**Remark 3.3.** We note that in order for our method of proof of Theorem 3.1 to work the important property of the subspace  $V_c$  that was needed was that  $u^+ \in V_c$  for any  $u \in V_c$ . We therefore see that Theorem 3.1 remains valid, if  $V_c$  is replaced by any subspace  $V$  which has this property (and, of course, the definitions of sub- and supersolutions are appropriately modified). This, more general theorem, for example, contains the sub-supersolution existence result for boundary-value problems subject to Neumann boundary conditions.

## REFERENCES

- [1] D. Gilbarg and N. Trudinger, *Elliptic Partial Differential Equations of Second Order*, Springer, Berlin, 1983.
- [2] P. Habets and K. Schmitt, *Nonlinear boundary value problems for systems of differential equations*, Arch. Math. **40**(1983), 441–446.
- [3] P. Hess, *On the solvability of nonlinear elliptic boundary value problems*, Ind. Univ. Math. J. **25**(1976), 461–466.
- [4] D. Kinderlehrer and G. Stampacchia, *An Introduction to Variational Inequalities and their Applications*, Academic Press, New York, 1980.
- [5] H. W. Knobloch, *Eine neue Methode zur Approximation periodischer Lösungen nicht linearer Differentialgleichungen zweiter Ordnung*, Mat. Z. **82**(1963), 177–197.
- [6] H. W. Knobloch and K. Schmitt, *Non-linear boundary value problems for systems of differential equations*, Proc. Roy. Soc. Edinburgh **78A**(1977), 139–159.
- [7] V. K. Le and K. Schmitt, *On boundary value problems for degenerate quasilinear elliptic equations and inequalities*, J. Diff. Eq. **144** (1998), 170–218.
- [8] J. L. Lions, *Quelques méthodes de résolution des problèmes aux limites non linéaires*, Dunod, Paris, 1969.
- [9] J. Mawhin, *Nonlinear functional analysis and periodic solutions of ordinary differential equations*, Tatra Summer School on ODE's, Difford 74 (1974), 37–60.
- [10] K. Schmitt, *Periodic solutions of nonlinear second order differential equations*, Math. Z. **98**(1967), 200–207.
- [11] K. Schmitt, *Periodic solutions of second order equations - a variational approach*, pp. 213–220 in *The First 60 Years of Nonlinear Analysis of Jean Mawhin*; edited by M. Delgado, A. Suárez, J. López-Gómez, and R. Ortega, World Scientific, Singapore, 2004.

VY KHOI LE

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF MISSOURI-ROLLA, ROLLA, MO 65401, USA

*E-mail address:* [vy@umr.edu](mailto:vy@umr.edu)

KLAUS SCHMITT

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF UTAH, 155 SOUTH 1400 EAST, SALT LAKE CITY, UT 84112, USA

*E-mail address:* [schmitt@math.utah.edu](mailto:schmitt@math.utah.edu)