

01 Jun 1986

Application of a Scaled Homogeneous Nucleation-Rate Formalism to Experimental Data at $T \ll T_c$

Barbara N. Hale

Missouri University of Science and Technology, bhale@mst.edu

Follow this and additional works at: https://scholarsmine.mst.edu/phys_facwork



Part of the [Physics Commons](#)

Recommended Citation

B. N. Hale, "Application of a Scaled Homogeneous Nucleation-Rate Formalism to Experimental Data at $T \ll T_c$," *Physical Review A*, vol. 33, no. 6, pp. 4156-4163, American Physical Society (APS), Jun 1986.

The definitive version is available at <https://doi.org/10.1103/PhysRevA.33.4156>

This Article - Journal is brought to you for free and open access by Scholars' Mine. It has been accepted for inclusion in Physics Faculty Research & Creative Works by an authorized administrator of Scholars' Mine. This work is protected by U. S. Copyright Law. Unauthorized use including reproduction for redistribution requires the permission of the copyright holder. For more information, please contact scholarsmine@mst.edu.

Application of a scaled homogeneous nucleation-rate formalism to experimental data at $T \ll T_c$

Barbara N. Hale

Department of Physics and Graduate Center for Cloud Physics Research, University of Missouri—Rolla, Rolla, Missouri 65401

(Received 24 December 1985)

It is pointed out that for temperatures $T < 0.5T_c$, where T_c is the critical temperature, the classical steady-state nucleation-rate formalism of Becker and Doring predicts an approximate critical supersaturation ratio S_{cr} (for the onset of nucleation) given by $\ln S_{cr}/\Omega^{3/2} \sim 0.53(T_c/T - 1)^{3/2}$. Ω is a material-dependent quantity approximately equal to the excess surface entropy per molecule. For most substances $\Omega \sim 2.0$ and for associated liquids $\Omega \sim 1.5$. The experimental data (for nucleation from vapor to liquid) from diffusion chamber and nozzle beam studies are found to be consistent with the above expression. The classical theory also predicts that for a supersaturation ratio S corresponding to constant J , $\ln S/\ln S_{cr} - 1 \sim \ln J/2\ln J_c$, where $\ln J_c$ is a quantity evaluated at the critical point and is ~ 72 for most materials. Expansion cloud-chamber data for nonane, toluene, and water are also found to be consistent with these approximate scaling laws.

I. INTRODUCTION

In this paper we present an expression for the classical Becker-Doring nucleation rate¹ J in terms of a slightly modified temperature dependence, $(T_c/T - 1)$, and a relatively material-independent constant $\ln J_c$, which is evaluated at the critical point. The motivation has been to develop an expression for J which can be simply compared to experimental data at temperatures far below the critical point. Of particular interest is a numerical estimate of the "critical" supersaturation S_{cr} corresponding to the onset of nucleation at a given reduced temperature T/T_c . The scaling of the nucleation rate has been considered extensively¹⁻⁸ and, in particular, Binder² has given a scaled form for the (slightly modified) classical nucleation rate near the critical point. The expression presented here is not applicable near T_c . It assumes a (classical) treatment of the prefactor, and concentrates rather on the material-dependent terms in the prefactor and on the use of the above reduced-temperature function to examine S_{cr} and factors in the free-energy barrier for $T \ll T_c$. The intent has been to cast the classical J into a form useful for quick estimates of required supersaturations, to point out factors which can be used to transform data into relatively material-independent form, and to suggest the $(T_c/T - 1)$ temperature dependence for data analysis far below T_c .

Some time ago, Wu, Wegener, and Stein⁹ demonstrated the approximate linear relationship between $\ln S_{cr}$ and $T^{-3/2}$ using experimental data for SF₆. However, the exploitation of a scaled or explicit temperature dependence of $\ln S_{cr}$ for $T \ll T_c$ was apparently not further pursued until McGraw¹⁰ examined a corresponding-states formalism and demonstrated that the data for $\ln S_{cr}$ fell into identifiable groups of substances when plotted versus T/T_c . Motivated primarily by McGraw's results, we derived for $T \ll T_c$ the approximation $\ln S_{cr} \sim 0.05$

$[A_0(T_c/T - 1)]^{3/2}$ and noted that the experimental data for $\ln S_{cr}$ when plotted versus $(T_c/T - 1)^{3/2}$ fell roughly into two groups with slopes (proportional to $A_0^{3/2}$) in the ratio of 3 to 2.¹¹ $A = A_0(T_c/T - 1)$ is the coefficient of $n^{2/3}$ in the classical free energy of formation, $An^{2/3} - n \ln S$, for the n molecule or atom cluster. This surface-tension term is the origin of the $(T_c/T - 1)$ dependence for $\ln S$ — which we note is different from the usual $\epsilon = (1 - T/T_c)$ dependence in critical-point formalisms. Rasmussen and Babu^{12,13} made use of our above temperature-dependent form for $\ln S_{cr}$ and showed that the slope of the experimental data for $\ln S_{cr}$, when plotted versus $(T_c/T - 1)^{3/2}$, is correlated with the Eotvos constant.¹⁴ The explicit relationship between the slope and the Eotvos constant was not given.

The "scaled supersaturation," $x = \ln S/A^{3/2}$, from which we in part extracted the above form for $\ln S_{cr}$, is identical in form to that of Binder and Stauffer,³ $\ln S/(b\epsilon)^\beta$, where β and δ are the standard critical-point exponents.¹⁵ Near the critical point $(T_c/T - 1) \sim \epsilon$, and $\beta\delta \sim 1.54$.³ (For classical three-dimensional fluids $\beta\delta = \frac{3}{2}$.) In a field-theoretic model for near-critical-point nucleation, Langer and Turski⁴ use a closely related quantity, the "scaled supercooling" $\tau = \delta T/(\epsilon T_c)$. The scaled supersaturation (or the scaled supercooling) influences the nucleation rate primarily via the so-called "energy of formation" of the critically sized cluster:⁷ $(x_0/x)^2 = (\tau_0/\tau)^2$. The x_0 and τ_0 are constants dependent on critical-point amplitudes.²⁻⁸ In the classical theory, $x_0 = 2/(3)^{3/2}$. However, τ_0 is less well defined for $T \ll T_c$ — and not independent of T . The classical nucleation rate J is proportional to $\exp[-(x_0/x)^2]$ and in the case that the kinetic prefactor has only slight S and T dependence, the $(x/x_0)^2$ is nearly constant for fixed J and leads to the approximate scaling law for $\ln S$.

Classical nucleation theory and theories applicable near the critical point differ primarily in the approximation of

the kinetic prefactor which describes the growth of the clusters subsequent to the nucleation event and before observation of macroscopic effects.^{2-8,16} In predicting critical-point phenomena the prefactor must account for diffusion-controlled growth and the vanishing of the diffusion constant as T approaches T_c .³ For $T \ll T_c$ diffusion-controlled growth is not in general applicable and the rate of formation of the new phase is primarily dictated by the birth (nucleation) of new phase embryos.^{2,8} However, in this low-temperature region the (classical) kinetic prefactor for vapor-to-liquid nucleation is proportional to the equilibrium vapor pressure squared and appears to be highly temperature and material dependent.

It is the temperature dependence of the classical kinetic prefactor which prompted Rassmussen and Babu¹² to comment that a theoretical explanation for the observed experimental agreement with the scaling law for $\ln S_{cr}$ (Ref. 11) was lacking. We here present a method of casting the kinetic prefactor into an approximately material-independent (and T/T_c -dependent) form and show that the classical theory predicts $\ln S_{cr}/\Omega^{3/2} \sim 0.53 (T_c/T - 1)^{3/2}$ where Ω is a material-dependent quantity approximately equal to the excess surface entropy per molecule. For most substances $\Omega \sim 2$, while for associated liquids $\Omega \sim 1.5$ and reflects the reduced entropy of the surface molecules. A modified scaling law for $\ln S$ corresponding to a constant J/J_c follows and we finally comment on the consequences of assuming a term in the energy of formation which takes into account the translation of the center of mass of the embryonic cluster.¹⁷ For simplicity we consider no replacement factors.¹⁸

The formalism for vapor-to-liquid nucleation is presented in Sec. II and in Sec. III the results are compared with experimental $\ln S_{cr}$ for $J \sim 1$, 10^4 , and $10^8 \text{ cm}^{-3} \text{ sec}^{-1}$. Comments and conclusions are in Sec. IV.

II. FORMALISM FOR J IN TERMS OF CRITICAL-POINT QUANTITIES

The classical Becker-Doring theory¹ for the steady-state homogeneous nucleation rate^{19,20} (including the so-called Zeldovitch factor²¹) can be written as follows:

$$J = J_c I (P/P_c)^\alpha \exp[-(x_0/x)^2], \quad (1)$$

where $\alpha = 2$,

$$I = [(4/3)/\pi^{1/2}]^{1/3} A^{1/2} (T_c/T)^{3/2} (\rho_c/\rho)^{2/3}, \quad (2)$$

and

$$J_c = (P_c/h)(\lambda_c/\rho_c)^{2/3} (P_c/kT_c\lambda_c). \quad (3)$$

The ρ , P , h , k , S , and λ are the number density, pressure, Planck's constant, Boltzmann's constant, supersaturation ratio, and inverse thermal wavelength cubed, $(2\pi mkT/h^2)^{3/2}$, respectively. The subscript c denotes critical-point quantities. The form for the exponent, $(x_0/x)^2$, follows from the classical free energy of formation for the n atom or molecule cluster, $g(n) = An^{2/3} - nB$, where $B = \ln S$.²² Classically, A is equal to the surface tension (divided by kT) times the area per surface molecule. From $dg(n^*)/dn = 0$, one readily obtains the number of molecules in the critical cluster,

$$n^* = (2A/3B)^3, \quad \text{and} \quad g(n^*) = 0.5n^*B = (x_0/x)^2. \quad \text{The} \quad x_0 = 2/(3)^{3/2}.$$

If one assumes the form $\sigma = \sigma_0(T_c - T)$ for the surface tension (where σ_0 is a material-dependent constant),²³ $A = (36\pi)^{1/3} \Omega (T_c/T - 1)$. The $\Omega = \sigma_0/k\rho^{2/3}$ can be interpreted as an effective excess surface entropy per molecule (in units of k) in the embryonic cluster. The bulk liquid value for Ω (the Eotvos constant¹⁴) is approximately 2 for most liquids. For associated liquids Ω is smaller (~ 1.5) and reflects the reduced excess entropy for surface molecules.²⁴ The corresponding values of A_0 are about 10 for ordinary substances and 7 for associated liquids. This approximate material independence of A_0 was noted when calculating thermodynamic properties of microscopic clusters using Monte Carlo methods and effective pair potentials.²⁵ In some preliminary work it was found that $A_0 \sim 10$ for Lennard-Jones argon clusters²⁵ and $A_0 \sim 7.5$ for Rahman-Stillinger²⁶ central-force (rigid-molecule) water clusters.²⁷ These values of A_0 correspond to $\Omega \sim 2.1$ and 1.6 for (Lennard-Jones) argon and (rigid-molecule central-force) water, respectively.

Using Eqs. (1) and (2), the scaled supersaturation $\ln S/A^{3/2}$ becomes

$$\ln S/A^{3/2} = x_0 (\ln J_c)^{-1/2} \delta_0, \quad (4)$$

where

$$\delta_0 = \{1 - \alpha[\ln(P_c/P_0) - \ln S]/\ln(J_c/J) + \ln I/\ln(J_c/J)\}^{-1/2} \quad (5a)$$

$$\sim 1 + 0.7\alpha W_0 (T_c/T - 1)^\eta / [2 \ln(J_c/J)] \quad (5b)$$

for $0.3 < T/T_c < 0.5$. For $\ln J = 0$, $\delta_0 \sim 1.13 \pm 0.04(T_c/T - 1)^\eta$. In obtaining this approximation, $\rho_c/\rho \sim \frac{1}{3}$ and

$$\ln(P_c/P_0) \sim W_0 (T_c/T - 1)^\eta \quad (6)$$

are used. W_0 and η are approximately 7 ± 2 and 1, respectively, for most substances, and W_0 can be roughly represented by L/kT_c , where L is the latent heat of vaporization near the boiling point. While there is some cancellation of the $\ln(P_c/P_0)$ term by $\ln S$ and $\ln I$ in Eq. (5a), $\ln I$ contributes less than 1.5% to δ_0 and the major contribution to the approximation in Eq. (5b) comes from $\ln(P_c/P_0)$. For substances (such as toluene) which have relatively small values of W_0 and large values of Ω (nonassociated liquids) there is a considerable cancellation of $\ln(P_c/P_0)$ by $\ln S$ and the temperature dependence of δ_0 is weak. Finally, one can show that $\ln J_c \sim 72$ to within 3% for most substances. For example, the values are 72.8, 74.7, 71.8, 70.8, and 73.7 for the substances ethanol, water, toluene, nonane, and argon, respectively. Since the square root of $\ln J_c$ enters into the expression for $\ln S_{cr}$, a 3% error in $\ln J_c$ produces a 1.5% error in $\ln S_{cr}$. Using $\ln J_c = 72$ the following approximate scaling laws result:

$$\ln S_{cr}/\Omega^{3/2} \sim 0.53 (T_c/T - 1)^{3/2} \quad \text{for} \quad J \sim 1 \text{ cm}^{-3} \text{ sec}^{-1} \quad (7)$$

and

$$\ln S \sim \ln S_{cr} [1 + \ln J / (2 \ln J_c)] \quad (8)$$

for larger J of physical interest. The major deviations from these approximate scaling laws occur at low temperatures where $(T_c/T - 1)$ is large (> 1.5). One can show also that the critical cluster size (for onset of nucleation) takes the form

$$n^* \sim (3/4\pi)^{1/2} (\ln J_c / \Omega)^{3/2} (T_c/T - 1)^{-3/2} \\ \sim 106(2/\Omega)^{3/2} (T_c/T - 1)^{-3/2}. \quad (9)$$

In expansion chamber experiments it is often more convenient to use the supercooling. Equation (6) (with the approximation $\eta = 1$) and Eq. (7) yield

$$\delta T' W_0 / \Omega^{3/2} \sim 0.53 (T_c/T - 1)^{3/2}, \quad (10)$$

where

$$\delta T' = (T_c/T_{\text{final}} - T_c/T_{\text{initial}}). \quad (11)$$

It is interesting to consider the modification to Eq. (7) when one includes the free energy associated with the

translation of the center of mass of the cluster.¹⁷ For simplicity, replacement factors¹⁸ are not considered. In this case the right-hand side of Eq. (1) is multiplied by $(\lambda kT/P)(n^*)^{3/2} \exp(-\frac{9}{4})$ and one considers the n^* to be modified by the small ($\sim 3\%$) correction factor, $[1 + 3/(2n^* \ln S)]^{-3} \sim [1 + (\frac{3}{4})(x/x_0)^2]^{-3}$. The resulting $J = J_t$ is given by Eq. (1) with $\alpha = 1$, I replaced by

$$I' = [(4/3)/\pi^{1/2}]^{1/3} 3^{9/4} (x_0/x)^{9/2} (T/T_c) \\ \times (\rho_c/\rho)^{2/3} A^{(-7/4)}, \quad (12)$$

J_c replaced by

$$J'_c = P_c (\lambda_c / \rho_c) / h, \quad (13)$$

and δ_0 replaced by δ_0 ($\alpha = 1$, $I = I'$, $J_c = J'_c$). Inclusion of the center-of-mass translational free energy reduces by one the power of P/P_c in the prefactor of J_t . This softens the temperature dependence of the prefactor and predicts a $\ln S_{cr}$ which is more nearly linear in




FIG. 1. Natural logarithm of the threshold ($J = 1 \text{ cm}^{-3} \text{ sec}^{-1}$) supersaturation ratio S_{cr} , divided by $\Omega^{3/2}$ from diffusion chamber and nozzle beam experimental data. The data points are for toluene (Ref. 33) (Δ), nonane (Ref. 32) (\times), water (Ref. 28) (\square), *n*-butylbenzene (Ref. 33) (\circ), sulfur hexafluoride (Ref. 9) ($+$), carbon tetrachloride (Ref. 34) (∇), chloroform (Ref. 34) (\diamond), ethanol (Ref. 31) (\diamond), octane (Ref. 33) ($*$), argon (Ref. 30) [taken from McGraw (Ref. 10, Fig. 1)] (\blacktriangledown), and acetic acid (Ref. 29) (\blacktriangle). The dashed line is $0.53 (T_c/T - 1)^{3/2}$ from Eq. (7). The values used for Ω are (Ref. 12) 2.35 for nonane, octane, and *n*-butylbenzene; 1.5 for water and ethanol; and 2.0 for SF_6 . For the remaining substances the ideal gas value 2.12 is used (Ref. 14).

$(T_c/T-1)^{3/2}$. In addition, $\ln J'_c$ is almost universally 86 and the prefactor for J_t becomes less material dependent. For example, for water, ethanol, toluene, nonane, and xenon the values of $\ln J'_c$ are 86.2, 86.1, 86.7, and 86.5, respectively. The value for argon (84.4) is notably smaller. The resulting scaling law (without replacement factors) is

$$\ln S'_{cr}/A^{3/2} = x_0 (\ln J'_c)^{-1/2} \delta'_0 \quad (14a)$$

$$\sim 0.44 (T_c/T-1)^{3/2} \quad (14b)$$

For $J_t \gg 1$, $\ln S'$ is related to $\ln S'_{cr}$ via Eq. (8) with $J_c = J'_c$.

III. COMPARISON WITH EXPERIMENTAL DATA

The approximations in Eqs. (7) and (8) serve as good predictors for $\ln S$ over a range of nucleation rates. Figure 1 shows experimental homogeneous vapor-to-liquid data for $\ln S_{cr}/\Omega^{3/2}$ for a number of substances,²⁸⁻³⁴ using bulk values²⁴ for Ω . The data for $\ln S_{cr}$ conform to the approximate scaling law in Eq. (7) rather well in spite of the scatter in data and the approximation of Ω by the

bulk value. In fact, the $\ln S_{cr}$ data appear to be more linear in $(T_c/T-1)^{3/2}$ than the corrections to Eq. (7) [via δ_0 , Eq. (5b)] would indicate. The linearity of the data of Katz *et al.*³³ for toluene is particularly striking, and it is noteworthy that almost all of the Katz data³¹⁻³⁴ fit this linear dependence extremely well. In Fig. 2 is plotted the $J \sim 10^4$ cm⁻³/sec expansion cloud-chamber data of Miller, Anderson, and Kassner for water^{35,36} and of Schmitt, Adams, and Zalabsky for toluene³⁷ and nonane.³⁸ The expansion chamber data appear to be consistent with the $(T_c/T-1)^{3/2}$ temperature dependence for $\ln S/\Omega^{3/2}$ as predicted by Eqs. (7) and (8).

It is interesting to compare the experimental nonane data for $J \sim 1$ (Katz *et al.*³²), $J \sim 10^4$ (Schmitt *et al.*³⁸), and $J \sim 10^8$ (Wagner and Strey³⁹) in a way which emphasizes the role of prefactor and exponent for J . The exponent

$$(x_0/x)^2 = (16\pi/3)(T_c/T-1)^3/(\ln S)^2 \Omega^3,$$

and if one uses $\Omega = \sigma/[kT\rho^{2/3}(T_c/T-1)]$ the standard classical model obtains. For most nonassociated liquids the Ω so calculated is stable with respect to T (in the




FIG. 2. $\ln S/\Omega^{3/2}$ for $J \sim 10^4$ cm⁻³sec⁻¹ from the expansion chamber data for water (Refs. 35 and 36) (\square), nonane (Ref. 38) (\times), and toluene (Ref. 37) (\triangle). The dashed line is the prediction from Eq. (8).

range of T/T_c of interest) to about 1%.²⁴ The stability of these values will depend somewhat on the choice of extrapolation for σ and ρ at low temperatures. In Fig. 3 is plotted the log of the prefactor versus $(x_0/x)^2$ for these data in the classical model. The three sets of data fall not too far from the straight line — which is the prediction of the classical model. The dashed lines above (below) the solid line indicate errors in J of 10^{-3} (10^{+3}). While the Wagner and Strey data and the Katz *et al.* data appear to be closer in magnitude to the classical model prediction, the expansion chamber data of Schmitt *et al.* show a more nearly linear relationship. The low-temperature data correspond in general to smaller values of $(x_0/x)^2$. Thus in Fig. 3 the high-temperature (expansion chamber) data lie furthest from the solid line and the classical model prediction.

In Fig. 4 is a comparison of the temperature dependence of the log of the prefactor and $(x_0/x)^2$ for the three sets of nonane data. For the diffusion chamber data it is assumed that $J=1 \text{ cm}^{-3} \text{ sec}^{-1}$. The prefactor has a smooth linear dependence on $(T_c/T-1)$. However, the temperature dependence of $(x_0/x)^2$ differs for the three

sets of data and none of the data appear to have quite the "classical model" form. This could be in part the result of the sensitivity of $(x_0/x)^2$ to the assumed temperature dependence of $\ln P_0$ and σ . Both Schmitt *et al.*³⁸ and Wagner and Strey³⁹ found major discrepancies in comparing their data with the classical model at low temperatures. For example, Schmitt *et al.* (using an equilibrium vapor pressure⁴⁰ different from the vapor pressure⁴¹ used by Katz and Wagner and Strey) found that their data disagreed with the classical model by factors of 10^9 in J at low temperatures. On the other hand, Wagner and Strey (using an expression for the surface tension⁴¹ different from that used by Schmitt *et al.* and Katz⁴²) found that their data disagreed with the classical model by factors of as much as 10^6 at low temperatures. As can be seen in Fig. 3, the vapor pressure⁴¹ and surface tension⁴² formulas used by Katz bring all the data into approximate mutual agreement with the classical model. We note that this does not imply that these particular formulas are without problems. This dilemma emphasizes the need to sort out competing temperature dependences of terms in $(x_0/x)^2$, and to assume a valid equilibrium vapor pressure at low




FIG. 3. $\ln[J_c/J(P/P_c)^2 I]$ vs the classical energy of formation, $(x_0/x)^2$, for the nonane diffusion chamber data of Katz *et al.* (Ref. 32) (\times); the nonane expansion chamber data of Schmitt *et al.* (Ref. 38) where $J \sim 10^4$ (\cdot); and the nonane expansion chamber data of Wagner and Strey (Ref. 39) (Δ), where $J \sim 10^8$. The $x = A^{3/2}/\ln S$ is the scaled supersaturation. The solid line is the prediction of the classical theory for J , and the lower (upper) dashed line indicates the range of data corresponding to $J \times 10^3$ ($J \times 10^{-3}$). Ω is calculated from $\sigma/[kT\rho^{2/3}(T_c/T-1)]$, as prescribed by the classical model.




FIG. 4. Classical model prefactor of J (open symbols) and $(x_0/x)^2$ (closed symbols) vs $(T_c/T - 1)$ for (a) the nonane $J = 10^4$ data of Schmitt *et al.* (Ref. 38) (\square ■) and the $J = 10^8$ data of Wagner and Strey (Ref. 39) (\triangle ▲) and (b) the nonane $J = 1$ data of Katz *et al.* (Ref. 32) (\diamond ◆). The equilibrium vapor pressure and surface tension formulas as given by Katz (Ref. 32) are used for all the data. The dashed lines guide the eye through the prefactor data points.

temperature. Finally, it seems appropriate to note that the expansion chamber data of Kassner, Miller, and Anderson^{35,36} and of Schmitt *et al.*^{37,38} offer stringent tests for the temperature dependence of the theory, and it is unfortunate that these data have been so long overlooked. It is noted, however, that the nonane and toluene expansion chamber data appear to give values of $\ln S$ which are slightly smaller than the classical model would predict at high temperatures. An approximate value of $\Omega = 2.35$ can be extracted from the Schmitt data³⁸ by plotting the $\ln(J/S^2)$ versus $(16\pi/3)(T_c/T-1)^3/(\ln S)^2$. It is interesting that this is the value of the Eotvos constant for nonane cited by Rasmussen¹² and that if one uses $\Omega = 2.35$ the data of Schmitt *et al.* agree with the classical model to less than a factor of 10 at all temperatures.

Some comments on the scaling law in Eq. (14) are relevant. The softened temperature dependence of the kinetic prefactor for J_t [which predicts a more linear dependence of $\ln S_{cr}$ on $(T_c/T-1)^{3/2}$] and the more nearly material independence of $\ln J'_c$ merit some consideration. However, it is well known that J_t/J (without replacement factors) is $\sim 10^{17}$ and unless the corresponding Ω could be increased by 15% the scaling law in Eq. (14) does not agree with experiment.

IV. COMMENTS AND CONCLUSIONS

The expression for the classical Becker-Doring nucleation rate presented here emphasizes the $(T_c/T-1)$ dependence, the relatively material-independent quantity $\ln J'_c$, and the overall weak temperature dependence of the prefactor for J when predicting $\ln S$ at $T \ll T_c$. The scaling law in Eq. (7) appears to describe the experimental $\ln S_{cr}$ for onset of nucleation rather well, and points out the usefulness of Ω in characterizing critical supersaturation values. An interpretation of Ω is the effective excess

entropy per surface molecule in the embryonic clusters. The fact that the bulk value for this quantity is nearly 2 for most substances and reduced to about 1.5 for associated liquids provides a convenient rule of thumb for estimating critical supersaturations for a wide variety of materials. The linearity of the data for $\ln S_{cr}$ [when plotted versus $(T_c/T-1)^{3/2}$] underscores the role of the $\frac{2}{3}$ exponent for the $An^{2/3}$ surface energy contribution in the classical energy of formation far below T_c . Katz's³³ toluene data and the cloud-chamber data^{35-38,43} for larger nucleation rates ($J \sim 10^4$) yield exponents for $(T_c/T-1)$ surprisingly close to 1.5. A comparison of the diffusion chamber and expansion chamber nonane data (with J ranging from 1 to 10^8) indicates that the classical model does a credible job of predicting J . There does appear to be, however, some anomalous temperature dependence in the $(x_0/x)^2$ data. Related to this anomaly are uncertainties in the low temperature dependence of the equilibrium vapor pressure, which can generate apparent discrepancies as large as 10^6 between data and the classical model. A careful consideration of the competing temperature factors in $(x_0/x)^2$ appears to be in order. Finally, we note the temperature functions considered here in predicting a general form for $\ln S$ and suggest that $(T_c/T-1)$ might be useful in analyzing phenomena far below the critical temperature. A modified supercooling, $(T_c/T_f - T_c/T_i)$, and the equilibrium vapor pressure in the form $\ln(P_c/P_0) = W_0(T_c/T-1)^7$ appear to be interesting examples.

ACKNOWLEDGMENTS

We would like to acknowledge helpful discussions with P. Wagner, R. Strey, and J. Taylor, and to thank J. Schmitt and G. Adams for making the details of their data available to us. This work is supported in part by the National Science Foundation under Grant No. ATM83-10854.

¹R. Becker and W. Doring, *Ann. Phys. (N.Y.)* **24**, 719 (1935).

²K. Binder, *J. Phys. C* **4**, 51 (1980).

³K. Binder and D. Stauffer, *Adv. Phys.* **25**, 343 (1976).

⁴J. S. Langer and L. A. Turski, *Phys. Rev. A* **8**, 3230 (1973); **22**, 2189 (1980).

⁵J. S. Langer and A. J. Schwartz, *Phys. Rev. A* **21**, 948 (1980).

⁶H. Furukawa and K. Binder, *Phys. Rev. A* **26**, 556 (1982).

⁷K. Binder, *Phys. Rev. A* **25**, 1699 (1982).

⁸J. D. Gunton, in *Phase Transitions and Critical Phenomena*, edited by C. Domb and J. L. Lebowitz (Academic, New York, 1983), Vol. 8.

⁹B. J. C. Wu, P. P. Wegener, and G. D. Stein, *J. Chem. Phys.* **68**, 308 (1978).

¹⁰R. McGraw, *J. Chem. Phys.* **75**, 5514 (1981).

¹¹B. N. Hale in *Proceedings of the 56th Colloid and Surface Science Symposium*, Virginia Polytechnic Institute, Blacksburg, Virginia (1982) (unpublished). A derivation of $\ln S_{cr}/A^{3/2} \sim 0.05$ was presented and the relation compared with experimental data for $\ln S_{cr}$ vs $(T_c/T-1)^{3/2}$. The figure presented was essentially identical to Fig. 1 of Rasmussen and Babu (Ref. 12). Mention of the Symposium paper can be

found in Ref. 19 of Rasmussen *et al.* (Ref. 13).

¹²D. H. Rasmussen and S. V. Babu, *Chem. Phys. Lett.* **108**, 449 (1984).

¹³D. H. Rasmussen, M. R. Appleby, G. L. Leedom, S. V. Babu, and R. J. Naumann, *J. Cryst. Growth* **64**, 229 (1983).

¹⁴In 1886 Eotvos found that the quantity $\sigma(M/d)^{2/3}$, where M is the molecular weight and d is the density, was a linear function of temperature. In cgs units, the proportionality constant for the ideal liquid is ~ 2.12 . For associated liquids, the constant is reduced. F. H. MacDougall, *Physical Chemistry* (Macmillan, New York, 1936), p. 96. The Eotvos constant defined in this way is actually Ω times $k(N_a)^{2/3}$, where N_a is Avogadro's number. This slight correction factor is ~ 0.984 .

¹⁵See, for example, H. E. Stanley, *Introduction to Phase Transitions and Critical Phenomena* (Oxford University Press, New York, 1971).

¹⁶R. G. Howland, N. C. Wong, and C. M. Knobler, *J. Chem. Phys.* **73**, 522 (1980).

¹⁷J. Lothe and G. M. Pound, *J. Chem. Phys.* **36**, 2080 (1962).

¹⁸H. Reiss, J. L. Katz, and E. R. Cohen, *J. Chem. Phys.* **48**, 5553 (1968); *Nucleation Phenomena*, edited by A. C.

- Zettlemoyer (Elsevier, New York, 1977).
- ¹⁹N. H. Fletcher, *The Physics of Rainclouds* (Cambridge University Press, Cambridge, England, 1969), Chap. 3.
- ²⁰F. F. Abraham, *Homogeneous Nucleation Theory* (Academic, New York, 1974).
- ²¹Y. B. Zeldovitch, *Acta Physicochim.* **18**, 11 (1943).
- ²²The $Ai^{2/3} \cdot iB$ notation appears to have been first used by D. Turnbull and J. C. Fisher, *J. Chem. Phys.* **17**, 71 (1948).
- ²³A linear fit to the surface tension data for liquids is quite successful over a wide range of temperatures, including a region not too far from the critical temperature (but outside the region of interest for critical phenomena). Some liquids (such as water) show anomalies in the slope near the freezing point. These anomalies are apparently related to the properties of associated liquids.
- ²⁴Eotvos constants for a variety of materials are given in Ref. 12. In comparing the results of the scaling laws in this paper with the data, the ideal value of 2.12 is used for the majority of substances. Exceptions are 1.5 for water and ethanol and 2.35 for nonane, octane, and *n*-butylbenzene. For SF₆ the value 2.0 from Ref. 12 is used. The latter values agree approximately with $\sigma / [(T_c/T - 1)kT\rho^{2/3}]$ for these substances for $0.35 < T/T_c < 0.50$. For example, this classical formula gives 2.44 ± 0.01 for nonane, 2.35 ± 0.02 for octane and *n*-butylbenzene, 2.17 ± 0.03 for toluene and carbon tetrachloride, 2.10 ± 0.01 for chloroform, 1.45 ± 0.05 for water, and 1.40 ± 0.20 for ethanol.
- ²⁵B. N. Hale and R. C. Ward, *J. Stat. Phys.* **28**, 487 (1982).
- ²⁶F. Stillinger and A. Rahman, *J. Chem. Phys.* **68**, 666 (1978).
- ²⁷P. Kemper and B. N. Hale (unpublished).
- ²⁸R. H. Heist and H. Reiss, *J. Chem. Phys.* **59**, 665 (1973).
- ²⁹R. H. Heist, K. M. Colling, and C. S. Dupuis, *J. Chem. Phys.* **65**, 382 (1976).
- ³⁰B. J. C. Wu, P. P. Wegener, and G. D. Stein, *J. Chem. Phys.* **69**, 1776 (1978).
- ³¹J. L. Katz and B. J. Ostermier, *J. Chem. Phys.* **47**, 478 (1967).
- ³²J. L. Katz, *J. Chem. Phys.* **52**, 4733 (1970).
- ³³J. L. Katz, C. J. Scoppa, N. G. Kumar, and P. Mirabel, *J. Chem. Phys.* **62**, 448 (1975).
- ³⁴J. L. Katz, P. Mirabel, C. J. Scoppa, and T. L. Virkler, *J. Chem. Phys.* **65**, 382 (1976).
- ³⁵R. J. Anderson, R. C. Miller, J. L. Kassner, and D. E. Hagen, *J. Atmos. Sci.* **37**, 2508 (1980).
- ³⁶R. C. Miller, R. J. Anderson, J. L. Kassner, and D. E. Hagen, *J. Chem. Phys.* **78**, 3204 (1983).
- ³⁷J. L. Schmitt, R. A. Zalabsky, and G. W. Adams, *J. Chem. Phys.* **79**, 4496 (1983).
- ³⁸G. W. Adams, J. L. Schmitt, and R. A. Zalabsky, *J. Chem. Phys.* **81**, 5074 (1984).
- ³⁹P. E. Wagner and R. Strey, *J. Chem. Phys.* **80**, 5266 (1984).
- ⁴⁰G. F. Carruth and R. Kobayashi, *J. Chem. Eng. Data* **18**, 115 (1973).
- ⁴¹Selected Values of Properties of Hydrocarbons and Related Compounds, American Petroleum Institute Research Project 44, Thermodynamics Research Center, Texas A & M University, College Station, Texas, 1965 (unpublished).
- ⁴²J. L. Jasper, *J. Phys. Chem. Ref. Data* **1**, 841 (1972).
- ⁴³J. L. Schmitt, G. W. Adams, and R. A. Zalabsky, *J. Chem. Phys.* **77**, 2089 (1981).