

## OPENNESS AND MONOTONEITY OF INDUCED MAPPINGS

WŁODZIMIERZ J. CHARATONIK

(Communicated by Alan Dow)

**ABSTRACT.** It is shown that for locally connected continuum  $X$  if the induced mapping  $C(f) : C(X) \rightarrow C(Y)$  is open, then  $f$  is monotone. As a corollary it follows that if the continuum  $X$  is hereditarily locally connected and  $C(f)$  is open, then  $f$  is a homeomorphism. An example is given to show that local connectedness is essential in the result.

All spaces considered in this paper are assumed to be metric. A *mapping* means a continuous function. We denote by  $\mathbb{N}$  the set of all positive integers, and by  $\mathbb{C}$  the complex plane. Given a space  $S$ , a point  $c \in S$  and a number  $\varepsilon > 0$ , we denote by  $B_S(c, \varepsilon)$  the open ball in  $S$  with center  $c$  and radius  $\varepsilon$ .

A *continuum* means a compact connected space. Given a continuum  $X$  with a metric  $d$ , we let  $2^X$  denote the hyperspace of all nonempty closed subsets of  $X$  equipped with the Hausdorff metric  $H$  defined by

$$H(A, B) = \max\{\sup\{d(a, B) : a \in A\}, \sup\{d(b, A) : b \in B\}\}$$

(see, e.g., [5, (0.1), p. 1 and (0.12), p. 10]). Further, we denote by  $C(X)$  the hyperspace of all subcontinua of  $X$ , i.e., of all connected elements of  $2^X$ , and by  $F_1(X)$  the hyperspace of singletons. The reader is referred to Nadler's book [5] for needed information on the structure of hyperspaces.

Given a mapping  $f : X \rightarrow Y$  between continua  $X$  and  $Y$ , we consider mappings (called the *induced* ones)

$$2^f : 2^X \rightarrow 2^Y \quad \text{and} \quad C(f) : C(X) \rightarrow C(Y)$$

defined by

$$2^f(A) = f(A) \quad \text{for every } A \in 2^X \quad \text{and} \quad C(f)(A) = f(A) \quad \text{for every } A \in C(X).$$

A mapping between continua is said to be:

- *open* provided the image of an open subset of the domain is open in the range;
- *monotone* provided the point-inverses are connected;
- *light* provided the point-inverses are zero-dimensional.

The following theorem is the main result of this paper.

**1. Theorem.** *Let a continuum  $X$  be locally connected, and a mapping  $f : X \rightarrow Y$  be such that the induced mapping  $C(f) : C(X) \rightarrow C(Y)$  is open. Then  $f$  is monotone.*

---

Received by the editors June 16, 1997 and, in revised form, January 4, 1998.

1991 *Mathematics Subject Classification.* Primary 54B20, 54F15, 54E40.

*Key words and phrases.* Continuum, hyperspace, induced mapping, open, monotone.

*Proof.* Assume  $f$  satisfies the assumptions of the theorem and that it is not monotone. Let  $p$  and  $q$  be two points of  $X$  such that  $f(p) = f(q)$  that belong to different components of  $f^{-1}(f(p))$ . By continuity of  $f$  there is a positive  $\varepsilon$  such that for every continuum  $L \subset Y$  such that  $f(p) \in L$  and  $H(L, \{f(p)\}) < \varepsilon$  the components of  $f^{-1}(L)$  containing  $p$  and  $q$  respectively are distinct. By local connectedness of  $Y$  there is a continuum  $V$  such that  $f(p) \in \text{int } V$  and  $H(V, \{f(p)\}) < \varepsilon$ , i.e.,  $V \subset B_Y(f(p), \varepsilon)$ . Let  $U_p$  and  $U_q$  be components of  $f^{-1}(V)$  containing  $p$  and  $q$  respectively. Since in locally connected continua components of open sets are open [4, §49, II, Theorem 4, p. 230], we conclude that  $p \in \text{int } U_p$  and  $q \in \text{int } U_q$ . Let  $\delta > 0$  be such that  $B_X(p, \delta) \subset U_p$  and  $B_X(q, \delta) \subset U_q$ .

Let  $\mathcal{B}$  be an order arc in  $C(Y)$  from  $\{f(p)\}$  to  $Y$  through  $V$ . Define  $\mathcal{A}$  as a subset of  $\mathcal{B}$  composed of all elements  $L \in \mathcal{B}$  such that the component of  $f^{-1}(L)$  containing  $p$  is distinct from the component of  $f^{-1}(L)$  containing  $q$ . Note that  $V \in \mathcal{A}$  and that if  $L, L' \in \mathcal{B}$ ,  $L \in \mathcal{A}$  and  $L' \subset L$ , then  $L' \in \mathcal{A}$ . Thus  $\mathcal{A}$  is a connected subset of  $\mathcal{B}$  containing  $\{f(p)\}$  and  $V$ . Since  $\mathcal{B} \setminus \mathcal{A}$  is closed, we see that  $\mathcal{A}$  is an open subset of  $\mathcal{B}$ . Let  $Q = \sup \mathcal{A} = \inf(\mathcal{B} \setminus \mathcal{A})$ . Then  $Q \in \text{cl } \mathcal{A} \setminus \mathcal{A}$ . Denote by  $P$  the component of  $f^{-1}(Q)$  containing both  $p$  and  $q$ . Openness of  $C(f)$  implies that  $f$  is open (see [3, Theorem 4.3, p. 243]; compare also [2, Theorem 3.2]), so  $f(P) = Q$  [6, (7.5), p. 148]. We will show that  $C(f)(B_{C(X)}(P, \delta))$  is not open in  $C(Y)$ . So, assume the contrary. Then there is a continuum  $K \in B_{C(X)}(P, \delta)$  with  $f(K) \in \mathcal{A}$ . Since  $p, q \in P$  and  $H(P, K) < \delta$ , we have  $K \cap U_p \neq \emptyset \neq K \cap U_q$ . Then  $U_p \cup K \cup U_q$  is a continuum containing both  $p$  and  $q$ , whose image  $f(U_p \cup K \cup U_q) = f(K)$  is in  $\mathcal{A}$ , contrary to the definition of  $\mathcal{A}$ . The proof is finished.  $\square$

**2. Corollary.** *Let a continuum  $X$  be hereditarily locally connected, and a mapping  $f : X \rightarrow Y$  be such that the induced mapping  $C(f) : C(X) \rightarrow C(Y)$  is open. Then  $f$  is a homeomorphism.*

*Proof.* It is enough to show that monotone open mappings on hereditarily locally connected continua are homeomorphisms. Assume the contrary, and let  $y \in Y$  be such that  $f^{-1}(y)$  is a nondegenerate continuum in  $X$ . Let  $\{y_n\}$  be an arbitrary sequence converging to  $y$ . Then continua  $f^{-1}(y_n)$  tend to  $f^{-1}(y)$ , so  $f^{-1}(y)$  is a nondegenerate continuum of convergence, contrary to hereditary local connectedness of  $X$ .  $\square$

**3. Example.** There are a continuum  $X$  and a mapping  $f : X \rightarrow X$  such that  $C(f) : C(X) \rightarrow C(X)$  is light and open, but not monotone.

*Proof.* Let  $S = \{z \in \mathbb{C} : |z| = 1\}$  be the unit circle. For  $n \in \mathbb{N}$  put  $X_n = S$ , and let  $\varphi_n : X_{n+1} \rightarrow X_n$  be defined by  $\varphi_n(z) = z^3$ . Then  $X = \varprojlim (X_n, \varphi_n)$  is the triodic solenoid. Define  $f : X \rightarrow X$  by  $f(\{z_1, z_2, \dots\}) = \{z_1^2, z_2^2, \dots\}$ , and note that  $f$  is well-defined. It has been proved in [1, Example 4.5] that the restriction  $C(f)|(C(X) \setminus \{X\})$  is two-to-one and  $C(f)^{-1}(X)$  is a singleton. Thus  $C(f)$  is light and it is not a homeomorphism. We will prove that  $C(f)$  is open. To this aim it is enough to show that the mapping is interior at each point of its domain [6, p. 149], i.e., that for each  $P \in C(X)$  and for each open neighborhood  $\mathcal{U}$  of  $P$  in  $C(X)$  we have  $C(f)(P) \in \text{int } C(f)(\mathcal{U})$ . For each  $n \in \mathbb{N}$  let  $f_n : X_n \rightarrow X_n$  be defined by  $f_n(z) = z^2$  (and thus  $f = \varprojlim f_n$ ), and let  $\pi_n : X \rightarrow X_n$  be the projection. Let  $P \in C(X)$  be a proper subcontinuum of  $X$ . Then there exists an index  $n \in \mathbb{N}$  such that  $\pi_{n-1}(P)$  is a proper subcontinuum of  $X_{n-1}$ , so  $\pi_n(P)$  is an arc of length less than  $2\pi/3$ . Let  $U_n$  be an open arc in  $X_n$  containing  $\pi_n(P)$  and having its length still less

than  $2\pi/3$ . Then the set  $\mathcal{V} = \{A \in C(X) : \pi_n(A) \in U_n\}$  is an open neighborhood of  $P$  in  $X$  such that the restriction  $C(f)|_{\mathcal{V}} : \mathcal{V} \rightarrow C(f)(\mathcal{V})$  is a homeomorphism onto the open set  $C(f)(\mathcal{U}) = \{A \in C(X) : \pi_n(A) \in f_n(U_n)\}$  containing  $C(f)(P)$ . So interiority of  $C(f)$  at  $P$  is shown in the case  $P \neq X$ . To prove that  $C(f)$  is interior at  $X$  consider, for  $n \in \mathbb{N}$ , the sets  $\mathcal{V}_n = \{A \in C(X) : \pi_n(A) = X_n\}$  and note that the family  $\{\mathcal{V}_n : n \in \mathbb{N}\}$  is a local base of (closed) neighborhoods of  $X$  on  $C(X)$ . So, it is enough to prove that  $C(f)(\mathcal{V}_n) \supset \mathcal{V}_{n+1}$ . To this end take  $A \in \mathcal{V}_{n+1}$ , and let  $B \in X$  be such that  $f(B) = A$ . Since

$$f_{n+1}(\pi_{n+1}(B)) = \pi_{n+1}(f(B)) = \pi_{n+1}(A) = X_{n+1},$$

we see that  $\pi_{n+1}(B)$  is an arc in  $X_{n+1}$  of length at least  $\pi$ . Thus  $\pi_n(B) = \varphi_n(\pi_{n+1}(B)) = X_n$ , i.e.,  $B \in \mathcal{V}_n$ , whence it follows that  $A = f(B) \in C(f)(\mathcal{V}_n)$ . The proof is then complete.  $\square$

In connection with Theorem 1 and Example 3 it would be interesting to know if a stronger result is true, namely whether or not the conclusion of Theorem 1 can be deduced from local connectedness of  $Y$  only (without assuming local connectedness of  $X$ ). In other words we have the following question.

**4. Question.** Can the assumption of local connectedness of the domain continuum  $X$  be relaxed to that of the range continuum  $Y$  in Theorem 1?

#### REFERENCES

1. J. J. Charatonik and W. J. Charatonik, *Lightness of induced mappings*, Tsukuba J. Math. **22** (1998), 179–192. CMP 98:16
2. H. Hosokawa, *Induced mappings between hyperspaces II*, Bull. Tokyo Gakugei Univ. (4) **44** (1992), 1–7. MR **94a**:54036
3. H. Hosokawa, *Induced mappings on hyperspaces*, Tsukuba J. Math. **21** (1997), 239–250. MR **98f**:54011
4. K. Kuratowski, *Topology*, vol. 2, Academic Press and PWN, New York, London and Warszawa, 1968. MR **41**:4467
5. S. B. Nadler, Jr., *Hyperspaces of sets*, M. Dekker, 1978. MR **58**:18330
6. G. T. Whyburn, *Analytic topology*, Amer. Math. Soc. Colloq. Publ. 28, Providence, 1942, reprinted with corrections 1971. MR **32**:425

MATHEMATICAL INSTITUTE, UNIVERSITY OF WROCLAW, PL. GRUNWALDZKI 2/4, 50-384 WROCLAW, POLAND

*E-mail address:* wjcharat@hera.math.uni.wroc.pl

DEPARTAMENTO DE MATEMÁTICAS, FACULTAD DE CIENCIAS, UNIVERSIDAD NACIONAL AUTÓNOMA DE MÉXICO, CIRCUITO EXTERIOR, CIUDAD UNIVERSITARIA, 04510 MÉXICO, D. F., MÉXICO

*E-mail address:* wjcharat@lya.fciencias.unam.mx