


01 Jan 1995

Point-Valued Mappings of Sets

Matt Insall

Missouri University of Science and Technology, insall@mst.edu

Follow this and additional works at: https://scholarsmine.mst.edu/math_stat_facwork

 Part of the [Mathematics Commons](#), and the [Statistics and Probability Commons](#)

Recommended Citation

M. Insall, "Point-Valued Mappings of Sets," *International Journal of Mathematics and Mathematical Sciences*, Hindawi Publishing Corporation, Jan 1995.

The definitive version is available at <https://doi.org/10.1155/S0161171295001049>

This Article - Journal is brought to you for free and open access by Scholars' Mine. It has been accepted for inclusion in Mathematics and Statistics Faculty Research & Creative Works by an authorized administrator of Scholars' Mine. This work is protected by U. S. Copyright Law. Unauthorized use including reproduction for redistribution requires the permission of the copyright holder. For more information, please contact scholarsmine@mst.edu.

POINT-VALUED MAPPINGS OF SETS

MATT INSALL

Department of Mathematics and Statistics
University of Missouri-Rolla
Rolla, MO 65401

(Received December 28, 1994 and in revised form January 18, 1995)

ABSTRACT. Let X be a metric space and let $CB(X)$ denote the closed bounded subsets of X with the Hausdorff metric. Given a complete subspace Y of $CB(X)$, two fixed point theorems, analogues of results in [1], are proved, and examples are given to suggest their applicability in practice.

KEY WORDS AND PHRASES. Fixed Point Theorems

1980 AMS SUBJECT CLASSIFICATION CODE. 47H10; 54H25

Let X be a metric space with metric d and let Y be a complete subspace of the space $CB(X)$ of all closed and bounded subsets of X , with the Hausdorff metric ρ :

$$\rho(A, B) = \max\{\sup_{x \in B} d(x, A), \sup_{x \in A} d(x, B)\}. \quad (1)$$

In Hicks [1], fixed point theorems for set-valued maps $T : X \rightarrow CB(X)$ were proved; and illustrated with examples. We show that similar results for maps $T : Y \rightarrow X$ can be obtained, using essentially the same techniques as in Hicks [1].

THEOREM 1. Let $T : Y \rightarrow X$ be continuous. Then there is an $A \in Y$ such that $T(A) \in A$ iff there exists a sequence $\{A_n\}_{n=0}^{\infty}$ in Y with $T(A_n) \in A_{n+1}$ (or $T(A_{n+1}) \in A_n$) and

$$\sum_{n=0}^{\infty} \rho(A_n, A_{n+1}) < \infty. \quad (2)$$

In this case, $A_n \rightarrow A$ as $n \rightarrow \infty$. (In fact, we may let $A_{n+1} = A_n \cup \{T(A_n)\}$, for each n , for the case $T(A_n) \in A_{n+1}$.)

PROOF. If $T(A) \in A$, then we are done. Conversely, if the given conditions are met, then $\{A_n\}_{n=0}^{\infty}$ is Cauchy, so let $A \in Y$ be its limit. Thus $T(A_n) \rightarrow T(A)$. If $y \in A$, then

$$d(y, T(A)) \leq d(y, T(A_n)) + d(T(A_n), T(A)), \quad (3)$$

so

$$d(A, T(A)) \leq d(A, T(A_n)) + d(T(A_n), T(A)). \quad (4)$$

Since $d(T(A_n), T(A)) \rightarrow 0$ and we have $d(A, T(A_n)) \leq \rho(A, A_{n+1}) \rightarrow 0$, it follows that $T(A) \in A$.

■

EXAMPLES

(1) Let $X = \mathbb{R}$, with the usual metric. Define $T : CB(\mathbb{R}) \rightarrow \mathbb{R}$ by

$$T(A) = \alpha \sup(A) + (1 - \alpha) \inf(A), \quad (5)$$

where $\alpha \in [0, 1]$. Then T is continuous. If $A \in CB(\mathbb{R})$, then

$$T(A \cup \{T(A)\}) = T(A) \in A \cup \{T(A)\}. \quad (6)$$

(2) Let $X = \mathbb{R}$ as in 1, and let $r : [0, \infty) \rightarrow [0, \infty)$ be such that $r \sim 1_{\mathbb{R}}$, where $1_{\mathbb{R}}$ is the identity on \mathbb{R} . Define $T : CB(\mathbb{R}) \rightarrow \mathbb{R}$ by

$$T(A) = \alpha r(\sup(A)) + (1 - \alpha)r(\inf(A)), \quad (7)$$

where $\alpha \in (0, 1)$. Assuming r is continuous, so is T . Let $A_0 \in CB(\mathbb{R})$, and for $n \in \mathbb{N}$, let

$$A_{n+1} = A_n \cup \left[\inf_{k \leq n} \{T(A_n)\}, \sup_{k \leq n} \{T(A_k)\} \right]. \quad (8)$$

Theorem 1 yields $A \in CB(\mathbb{R})$ with $T(A) \in A$ if

$$\sum_{n=1}^{\infty} \max \left\{ d \left(\inf_{k \leq n} \{T(A_n)\}, A_n \right), d \left(\sup_{k \leq n} \{T(A_k)\}, A_n \right) \right\} < \infty. \quad (9)$$

DEFINITION. Let (X, d) be a metric space and let Y be a subspace of $(CB(X), \rho)$. Let $T : Y \rightarrow X$. Then T is nice if for each $A \in Y$ and each $x \in A$ with $d(x, T(A)) = d(A, T(A))$, there exists a set $B \in Y$ with $T(B) = x$.

EXAMPLES

(3) Let $X = \mathbb{R}^2$, $T : CB(\mathbb{R}^2) \rightarrow \mathbb{R}^2$ defined by

$$T(A) = (\inf(\text{proj}_1(A)), \sup(\text{proj}_1(A))). \quad (10)$$

Let $a > b$ and $A = [0, a] \times [0, b]$. Then $T(A) = (0, a)$, and $(0, b)$ is the only point of A whose distance from $(0, a)$ equals $d(A, T(A))$. Let $B = [0, b]^2$. Then $T(B) = (0, b)$.

(4) Let $X = \mathbb{R}^2$, and for $A \in CB(\mathbb{R}^2)$, let $T(A)$ be the center of the circle which circumscribes A .

Let $r = d(A, T(A))$, and let $x \in A$ with $d(x, T(A)) = r$. Let $B = A \cap \overline{\mathcal{B}(x, \frac{\text{diam}(A)}{2})}$. Then

$$T(B) = x.$$

THEOREM 2. Let (X, d) be a metric space and let Y be a complete subspace of $(CB(X), \rho)$, each member of which is compact. Let $T : Y \rightarrow X$ be continuous. Assume that $K : [0, \infty) \rightarrow [0, \infty)$ is non-decreasing, $K(0) = 0$, and

$$\rho(A, B) \leq K(d(T(A), T(B))) \quad (11)$$

for $A, B \in Y$. If T is nice, then there is $A \in Y$ such that $T(A) \in A$ iff there exists $A_0 \in Y$ for which

$$\sum_{n=1}^{\infty} K^n(d(A_0, T(A_0))) < \infty \quad (*)$$

In this case, we can choose $\{A_n\}_{n=1}^{\infty}$ such that $T(A_{n+1}) \in A_n$ and $A_n \rightarrow A$.

PROOF. If $T(A) \in A$, then we are done. If $A_0 \in Y$ satisfies (*), let $x_1 \in A_0$ with $d(x_1, T(A_0)) = d(A_0, T(A_0))$. Since T is nice, let $A_1 \in Y$ with $T(A_1) = x_1$.

Next, let $x_2 \in A_1$ with $d(x_2, T(A_1)) = d(A_1, T(A_1))$, and then let $A_2 \in Y$ with $T(A_2) = x_2$. Then

$$\begin{aligned} d(T(A_1), T(A_2)) &= d(T(A_1), x_2) \\ &= d(T(A_1), A_1) = d(x_1, A_1) \\ &\leq \rho(A_0, A_1) \leq K(d(T(A_0), T(A_1))), \end{aligned} \quad (12)$$

so that

$$\begin{aligned} K(d(T(A_1), T(A_2))) &\leq K^2(d(T(A_0), T(A_1))) \\ &= K^2(d(T(A_0), x_1)) \\ &= K^2(d(T(A_0), A_0)). \end{aligned} \quad (13)$$

Now, suppose we have $x_n \in A_{n-1}$ and $A_n \in Y$ with $d(x_n, T(A_{n-1})) = d(A_{n-1}, T(A_{n-1}))$ and $T(A_n) = x_n$. Let $x_{n+1} \in A_n$ with $d(x_{n+1}, T(A_n)) = d(A_n, T(A_n))$ and let $A_{n+1} \in Y$ with $T(A_{n+1}) = x_{n+1}$. Then

$$\begin{aligned} d(T(A_n), T(A_{n+1})) &= d(T(A_n), x_{n+1}) \\ &= d(T(A_n), A_n) = d(x_n, A_n) \\ &\leq \rho(A_{n-1}, A_n) \leq K(d(T(A_{n-1}), T(A_n))), \end{aligned} \quad (14)$$

so that

$$\begin{aligned} K(d(T(A_n), T(A_{n+1}))) &\leq K^2(d(T(A_{n-1}), T(A_n))) \\ &= K(K(d(T(A_{n-1}), T(A_n)))) \\ &\leq K(K^2(d(T(A_{n-2}), T(A_{n-1})))) \\ &= K^3(d(T(A_{n-2}), T(A_{n-1}))) \\ &\leq \dots \leq K^n(d(T(A_0), A_0)). \end{aligned} \quad (15)$$

Thus, since

$$\rho(A_n, A_{n+1}) \leq K(d(T(A_n), T(A_{n+1}))), \quad (16)$$

it follows from (*) that

$$\sum_{n=0}^{\infty} \rho(A_n, A_{n+1}) < \infty, \quad (17)$$

and then by Theorem 1, $A_n \rightarrow A$ and $T(A) \in A$. ■

Note that the conditions of theorem 2 force T to be a bijection. In both of these theorems, we have used completeness of the given subspace Y of $CB(X)$ instead of completeness of X . In fact, in theorem 2, since T is a bijection, we may trade completeness of Y back for completeness of X and use the second theorem of Hicks [1].

THEOREM 3. If (X, d) is a complete metric space and Y is any subspace of $(CB(X), \rho)$, each member of which is compact, then for any homeomorphism $T : Y \rightarrow X$ such that

$$\rho(A, B) \leq K(d(T(A), T(B))), \quad (18)$$

where $K : [0, \infty) \rightarrow [0, \infty)$ is nondecreasing, with $K(0) = 0$, there is $A \in Y$ such that $T(A) \in A$ iff there exists $A_0 \in Y$ for which (*) holds.

PROOF. If $A_0 \in Y$ satisfies (*), let $x_0 = T(A_0)$. Apply theorem 2 of Hicks [1] to $T^{-1} : X \rightarrow Y$ to obtain a $p \in X$ such that $p \in T^{-1}(p)$. Let $A = T^{-1}(p)$. Then $T(A) = p$ is in A , so we are done. ■

REFERENCES

- [1] HICKS, T.L. Fixed-Point Theorems for Multivalued Mappings, Indian J. Pure Appl. Math. 20(11) (November 1989), 1077-1079