
01 Jan 2018

Mini Review: A Note on Nonoscillatory Solutions for Higher Dimensional Time Scale Systems

Elvan Akin


Missouri University of Science and Technology, akine@mst.edu

Ozkan Ozturk

Ismail Ugur Tiryaki

Gulsah Yeni

Follow this and additional works at: https://scholarsmine.mst.edu/math_stat_facwork

 Part of the [Mathematics Commons](#), and the [Statistics and Probability Commons](#)

Recommended Citation

E. Akin et al., "Mini Review: A Note on Nonoscillatory Solutions for Higher Dimensional Time Scale Systems," *International Journal of Applied and Experimental Mathematics*, vol. 3, no. 1, Graphy Publications, Jan 2018.



This work is licensed under a [Creative Commons Attribution 3.0 License](#).

This Article - Journal is brought to you for free and open access by Scholars' Mine. It has been accepted for inclusion in Mathematics and Statistics Faculty Research & Creative Works by an authorized administrator of Scholars' Mine. This work is protected by U. S. Copyright Law. Unauthorized use including reproduction for redistribution requires the permission of the copyright holder. For more information, please contact scholarsmine@mst.edu.

A Note on Nonoscillatory Solutions for Higher Dimensional Time Scale Systems

Elvan Akin*, Ozkan Ozturk, Ismail U. Tiryaki and Gulsah Yeni

Department of Mathematics and Statistics, Missouri University of Science and Technology, 400 W 12th Street, Rolla, Missouri 65409 USA

Abstract

In this paper, we focus on nonoscillatory solutions of two (2D) and three (3D) dimensional time scale systems and discuss nonexistence of such solutions.

Publication History:

Received: November 17, 2017

Accepted: January 18, 2018

Published: January 20, 2018

Keywords:

Paraconsistent logic, Linear-time temporal logic, Clinical reasoning verification, Paraconsistent model

Introduction

The motivation of studying dynamic equations on time scales is to unify continuous and discrete analysis and harmonize them in one comprehensive theory and eliminate obscurity from both. A *time scale* T is an arbitrary nonempty closed subset of the real numbers \mathbb{R} . The most well known examples for time scales are \mathbb{R} (leading to differential equations, [10]), \mathbb{Z} (leading to difference equations, [11]), and $q^{\mathbb{N}_0} := \{1, q, q^2, \dots\}$, $q > 1$ (leading to q -difference equations, [9]). In 1988, the theory of time scales was initiated by Stefan Hilger in his Ph.D. thesis, [8]. Since the calculus of time scales has been recently developed, we give a brief introduction to time scales calculus. Nevertheless, an excellent introduction can be found in [5, 6] by Bohner and Peterson.

There are two jump operators: For $t \in T$, the *forward jump operator* $\sigma: T \rightarrow T$ is given by $\sigma(t) := \inf \{s \in T : s > t\}$ for all $t \in T$ while the *backward jump operator* $\rho: T \rightarrow T$ is dened by $\rho(t) := \sup \{s \in T : s < t\}$ for all $t \in T$. Finally, the *graininess function*: $\mu: T \rightarrow [0, \infty)$ is given by $\mu(t) := \sigma(t) - t$ for all $t \in T$.

We dene $\inf 0 = \sup T$. There are four types of points in T . If $\sigma(t) > t$, hen t is called *right - scattered*, while if $\rho(t) < t$, t is called *left - scattered*. If t is right and left - scattered at the same time, then we say that t is *isolated*. If $t < \sup T$ and $\sigma(t) = t$, then t is called *right - dense*, while if $t > \inf T$ and $\rho(t) = t$, we say t is left - dense. Also, if t is right and left - dense at the same time, then we say that t is *dense*.

If $\sup T < \infty$, then $T^k = T \setminus (\rho(\sup T), \sup T]$, and $T^k = T$ if $\sup T = \infty$. Suppose that $f: T \rightarrow \mathbb{R}$ is a function. Then $f^\Delta: T \rightarrow \mathbb{R}$ is dened by $f^\Delta(t) = \frac{f(\sigma(t)) - f(t)}{\mu(t)}$ for all $t \in T$. For any $\varepsilon > 0$, if there exists a $\delta > 0$ such that

$$|f^\sigma(t) - f(s) - f^\Delta(t)(\sigma(t) - s)| \leq \varepsilon |\sigma(t) - s| \text{ for all } s \in (t - \delta, t + \delta) \cap T,$$

then f is called *delta (or Hilger) differentiable* on T^k and f^Δ is called *delta derivative* of f . Let $f: T \rightarrow \mathbb{R}$ be a function with $t \in T^k$. If f is differentiable at t , f is continuous at t . If f is continuous at t and t is right-scattered, then f is differentiable at t and

$$f^\Delta(t) = \frac{f^\sigma(t) - f(t)}{\mu(t)}$$

If t is right dense, then f is differentiable at t if

$$f^\Delta(t) = \lim_{s \rightarrow t} \frac{f^\sigma(t) - f(s)}{t - s}$$

exists as a finite number. If f is differentiable at t , then $f^\Delta(t) = f(t) + \mu(t)f^\Delta(t)$ holds for all type of points in T . Let $f, g: T \rightarrow \mathbb{R}$ be differentiable at $t \in T^k$. Then we have the product and quotient rules for derivatives as follows:

1. $(fg)^\Delta(t) = f^\Delta(t)g(t) + f^\sigma(t)g^\Delta(t) + f^\Delta(t)g^\sigma(t)$
2. If $g(t)g(\sigma(t)) \neq 0$, then $\frac{f}{g}$ is differentiable at t with

$$\left(\frac{f}{g}\right)^\Delta(t) = \frac{f^\Delta(t)g(t) - f(t)g^\Delta(t)}{g(t)g^\sigma(t)}$$

$f: T \rightarrow \mathbb{R}$ is called *right dense continuous* (rd-continuous), denoted by $C_{rd}, C_{rd}(T)$; or $C_{rd}(T, \mathbb{R})$, if it is continuous at right dense points in T and its left sided limits exist as a finite number at left dense points in T . Throughout we denote continuous functions by C . Also, the Cauchy integral is dened by

$$\int_a^b f(t) \Delta t = F(b) - F(a) \text{ for all } a, b \in T$$

Every rd-continuous function has an antiderivative. Moreover, F given by

$$F(t) = \int_{t_0}^t f(s) \Delta s \text{ for } t \in T$$

is an antiderivative of f .

Higher Dimensional Time Scale Systems

The study of higher dimensional time scale systems in nature and society has been motivated by their applications such as population dynamics, genomic and neuron dynamics and epidemiology in biological sciences, [1,16] and in astrophysics, gas dynamics and fluid mechanics, relativistic mechanics, nuclear physics, and chemically reacting systems, [4,7,12,17].

Corresponding Author: Dr. Elvan Akin, Department of Mathematics and Statistics, Missouri University of Science and Technology, 400 W 12th Street, Rolla, Missouri 65409 USA, E-mail: akine@mst.edu

Citation: Akn E, Ozturk O, Tiryaki IU, Yeni G (2018) A Note on Nonoscillatory Solutions for Higher Dimensional Time Scale Systems. Int J Appl Exp Math 3: 126. doi: <https://doi.org/10.15344/2456-8155/2018/126>

Copyright: © 2018 Akn et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Recently, nonoscillatory solutions of 2D and 3D time scale systems have been considered in [13-15] and [2,3], respectively. One of our main goals in these articles is to find integral conditions which eliminate nonoscillatory solutions of given systems. Depending on which dimension we deal with one naturally attempts to find single, double or triple integral conditions. A question we have here is whether we can eliminate all types of nonoscillatory solutions or not. In this contribution, we only satisfy single integral conditions for both time scale systems and other integral conditions can be found in references above. Our approach is based on the sign of components of nonoscillatory solutions of systems and we assume without loss of generality that the first component of such solutions is always positive.

We first start with the following 2D time scale system

$$\begin{cases} x^\Delta(t) = a(t)f(y(t)) \\ y^\Delta(t) = -b(t)g(x(t)), \end{cases} \quad (2.1)$$

where $a, b \in C_{rd}([t_0, \infty)_T, \mathbb{R}^+)$ and $f, g \in C(\mathbb{R}, \mathbb{R})$ satisfying $uf(u) > 0, ug(u) > 0$ for $u \neq 0$ and

$$a, b \in C_{rd}([t_0, \infty)_T, \mathbb{R}^+) \quad (2.2)$$

where F and G are positive constants and Φ_α and Φ_β are odd power functions, i.e., $\Phi_p(u) = |u|^p \operatorname{sgn} u, p > 0$ and $p \in \{\alpha, \beta\}$.

Throughout this article, we assume that T is unbounded above. We call (x, y) a *proper solution* if it is defined on $[t_0, \infty)_T$ and $\sup\{|x(s)|, |y(s)| : s \in [t, \infty)_T\} > 0$ for $t \geq t_0$. By $t \geq t_1$, we mean $t \in [t_1, \infty)_T := [t_1, \infty) \cap T$. A solution (x, y) of (2.1) is said to be *nonoscillatory* if the component functions x and y are both nonoscillatory, i.e., either eventually positive or eventually negative. Otherwise, it is said to be *oscillatory*. Definitions above can be modified for higher dimensional time scale systems.

Assume that (x, y) is a nonoscillatory solution of system (2.1) such that x oscillates but y is eventually positive. Then the first equation of system (2.1) yields $x^\Delta(t) = a(t)f(y(t)) > 0$ eventually one sign for all large $t \geq t_0$, a contradiction. The case where y is eventually negative is similar. Therefore, we have that the component functions x and y are themselves nonoscillatory. In other words, any nonoscillatory solution (x, y) of system (2.1) is one of the following types:

- Type (a): $\operatorname{sgn} x(t) = \operatorname{sgn} y(t)$;
- Type (b): $\operatorname{sgn} x(t) \neq \operatorname{sgn} y(t)$.

For convenience, let us set

$$A(t_0) = \int_{t_0}^\infty a(t)\Delta t \text{ and } B(t_0) = \int_{t_0}^\infty b(t)\Delta t, \quad t_0 \in T \quad (2.3)$$

Theorem

Any nonoscillatory solution of system (2.1) cannot be of

1. Type (a) if $A(t_0) < 1$ and $B(t_0) = 1$.
2. Type (b) if $A(t_0) = 1$ and $B(t_0) < 1$.

Secondly, let's consider 3D time scale systems of the form

$$\begin{cases} x^\Delta(t) = a(t)f(y(t)) \\ y^\Delta(t) = b(t)g(z(t)) \\ z^\Delta(t) = -c(t)h(x(t)) \end{cases} \quad (2.4)$$

where $a, b, c \in C_{rd}([t_0, \infty)_T, \mathbb{R}^+)$ and $f, g, h \in C(\mathbb{R}, \mathbb{R})$ satisfying $uf(u) > 0, ug(u) > 0$, and $uh(u) > 0$ for $u \neq 0$ and

$$\frac{f(u)}{\Phi_\alpha(u)} \geq F, \quad \frac{g(u)}{\Phi_\beta(u)} \geq G, \quad \frac{h(u)}{\Phi_\gamma(u)} \geq H \text{ for all } u \neq 0, \quad (2.5)$$

where F, G , and H are positive constants and $\Phi_p(u) = |u|^p \operatorname{sgn} u, p > 0$ and $p \in \{\alpha, \beta, \gamma\}$. By a similar argument, we can conclude that any nonoscillatory solution (x, y, z) of (2.4) is one of following types:

- Type (a): $\operatorname{sgn} x(t) = \operatorname{sgn} y(t) = \operatorname{sgn} z(t)$,
- Type (b): $\operatorname{sgn} x(t) \neq \operatorname{sgn} y(t) = \operatorname{sgn} z(t)$,
- Type (c): $\operatorname{sgn} x(t) = \operatorname{sgn} y(t) \neq \operatorname{sgn} z(t)$,
- Type (d): $\operatorname{sgn} x(t) = \operatorname{sgn} z(t) \neq \operatorname{sgn} y(t)$.

For convenience let's set

$$C(t_0) = \int_{t_0}^\infty c(t)\Delta t, \quad t_0 \in T$$

Lemma

Any nonoscillatory solution of system (2.4) can not be of

9. Type (a) if $C(t_0) = 1$.
10. Type (c) if $A(t_0) = 1$;
11. Type (d) if $B(t_0) = 1$.

Not only from Lemma 2.2 but also from other results in above references one can observe that Type (b) solution of system (2.1) is not eliminated. In fact, it is worth to emphasize that components of Type (b) solutions have finite limits. Note that when we deal with 2D time scale systems such type does not occur. Here the question comes to our mind: Is it related with dimensions of given systems? When we focus on four dimensional (4D) time scale systems, we realize that nonoscillatory solutions whose components have finite limits do not occur either. Moreover, we have already known that all types of nonoscillatory solutions of 4D time scale systems can be eliminated by some integral conditions. Therefore, we would like to emphasize that this is a matter of considering odd or even dimensional time scale systems.

Competing Interests

The authors declare that there is no competing interest regarding the publication of this article.

References

1. Agarwal RP, O'Regan D, Saker SH (2014) Oscillation and Stability of Delay Models in Biology. Springer.
2. Ekin-Bohner E, Dosla Z, Lawrence B (2012) Almost oscillatory Three Dimensional Dynamic Systems. Adv. Difference Equ 2012: 46.
3. Akin E, Ozturk O (2016) Nonoscillation Criteria for Three Dimensional Time-Scale Systems.
4. Arthur AM, Robinson PD (1969) Complementary variational principle for $\nabla^2 = f(\Phi)$ with applications to the Thomas - Fermi and Liouville equations. Proc Cambridge Philos Soc 65: 535-542.
5. Bohner M, Peterson A (2001) Dynamic Equations on Time Scales: An Introduction with Applications. Birkhauser, Boston.
6. Bohner M, Peterson A (2003) Advances in Dynamic Equations on Time Scales. Birkhauser, Boston.
7. Davis HT (1960) Introduction to Nonlinear Differential Integral Equations U.S. Atomic Energy Commission, Washington DC.
8. Hilger S (1988) Ein Maßkettenkalkül mit Anwendung auf Zentrumsmannigfaltigkeiten [thesis]. Universität Würzburg.

-
9. Kac V, Cheung P (2002) *Quantum calculus*. Universitext, Springer-Verlag, New York.
 10. Kelley WG, Peterson AC (2010) *The Theory of Differential Equations: Classical and Qualitative*, Springer.
 11. Kelley WG, Peterson AC (2001) *Difference Equations, Second Edition: An Introduction with Applications*. Academic Press.
 12. Nehari Z (1963) On a nonlinear differential equation arising in nuclear physics. *Proc. Roy. Irish Academy Sect A*.
 13. Ozturk O, Akin E (2016) Nonoscillation Criteria for Two Dimensional Time-Scale Systems. *Nonauton Dyn Syst* 3: 1-13.
 14. Ozturk O, Akin E (2016) On Nonoscillatory Solutions of Two Dimensional Nonlinear Delay Dynamical Systems. *Opuscula Math* 36: 651-669.
 15. Ozturk O, Akin E, Tiryaki IU (2015) On Nonoscillatory Solutions of Emden-Fowler Dynamic Systems on Time Scales. *Filomat*.
 16. Ruan S, Wolkowicz GSK, Wu J (1999) *Differential Equations with Applications to Biology*. AMS.
 17. Shevchuk VN (1965) Problems methods and fundamental results in the theory of oscillation of solutions of nonlinear nonautonomous ordinary differential equations. *Proc. 2nd All-Union Conf. on Theoretical and Appl Mech, Moscow*.