

01 May 2011

Self-Energy Correction to the Hyperfine Splitting for Excited States

Benedikt J. Wundt

Ulrich D. Jentschura

Missouri University of Science and Technology, ulj@mst.edu

Follow this and additional works at: https://scholarsmine.mst.edu/phys_facwork

 Part of the [Physics Commons](#)

Recommended Citation

B. J. Wundt and U. D. Jentschura, "Self-Energy Correction to the Hyperfine Splitting for Excited States," *Physical Review A - Atomic, Molecular, and Optical Physics*, vol. 83, no. 5, pp. 052501-1-052501-6, American Physical Society (APS), May 2011.

The definitive version is available at <https://doi.org/10.1103/PhysRevA.83.052501>

This Article - Journal is brought to you for free and open access by Scholars' Mine. It has been accepted for inclusion in Physics Faculty Research & Creative Works by an authorized administrator of Scholars' Mine. This work is protected by U. S. Copyright Law. Unauthorized use including reproduction for redistribution requires the permission of the copyright holder. For more information, please contact scholarsmine@mst.edu.

Self-energy correction to the hyperfine splitting for excited states

B. J. Wundt and U. D. Jentschura

Department of Physics, Missouri University of Science and Technology, Rolla, Missouri 65409-0640, USA

(Received 2 February 2011; published 4 May 2011)

The self-energy corrections to the hyperfine splitting is evaluated for higher excited states in hydrogenlike ions using an expansion in the binding parameter $Z\alpha$, where Z is the nuclear-charge number and α is the fine-structure constant. We present analytic results for D , F , and G states, and for a number of highly excited Rydberg states, with principal quantum numbers in the range $13 \leq n \leq 16$, and orbital angular momenta $\ell = n - 2$ and $\ell = n - 1$. A closed-form analytic expression is derived for the contribution of high-energy photons, valid for any state with $\ell \geq 2$ and arbitrary n , ℓ , and total angular momentum j . The low-energy contributions are written in the form of generalized Bethe logarithms and evaluated for selected states.

DOI: [10.1103/PhysRevA.83.052501](https://doi.org/10.1103/PhysRevA.83.052501)

PACS number(s): 31.15.-p, 12.20.Ds, 06.20.Jr

I. INTRODUCTION

The self-energy correction to the hyperfine splitting is the dominant quantum electrodynamic (QED) correction to the magnetic interaction of the bound electron with the field of the nucleus. The hyperfine interaction energy of electron and nucleus is proportional to $g_N \alpha (Z\alpha)^3 m_e^2 / m_N$, where g_N is the nuclear g factor, and m_e and m_N are the electron and nuclear masses, respectively. Relativistic corrections enter at relative order $(Z\alpha)^2$. The dominant QED correction is due to the anomalous magnetic moment of the electron and enters at relative order α . Here, we consider the QED correction of order $\alpha (Z\alpha)^2$, which is the sum of a high- and a low-energy part. Relativistic corrections to the anomalous magnetic interaction give one of the dominant contributions to the high-energy part, which can otherwise be calculated on the basis of a form-factor approach, using a generalized Dirac equation in which the radiative effects and the hyperfine interaction are inserted “by hand.” The low-energy part constitutes a correction to the Bethe logarithm due to the hyperfine interaction. It can be formulated as a hyperfine correction to the self-energy, the effect being equivalent to the self-energy correction to the hyperfine splitting mediated by low-energy virtual photons [up to order $\alpha (Z\alpha)^2$].

In our treatment, we follow the formalism of nonrelativistic QED (NRQED) detailed in Ref. [1], and refer to Refs. [2–7] for a number of previous investigations regarding the treatment of the self-energy correction to the hyperfine splitting in systems with low nuclear-charge number.

Our paper is organized as follows. The general formalism of the hyperfine interaction is described in Sec. II. For the self-energy correction, the low-energy part is treated in Sec. III, and the high-energy part is calculated in Sec. IV. Results and theoretical predictions are discussed in Sec. V. Conclusions are reserved for Sec. VI. Natural units with $\hbar = c = \epsilon_0 = 1$ are used throughout the paper.

II. FORMALISM

Following the derivation in Ref. [8], the magnetic dipole field of the nucleus is described by the vector potential

$$\vec{A}_{\text{hfs}}(\vec{x}) = -\frac{1}{4\pi} \frac{\vec{\mu} \times \vec{x}}{r^3}, \quad (1)$$

where \vec{x} is the coordinate vector and $r = |\vec{x}|$. The curl of this vector potential yields the magnetic field

$$\vec{B}_{\text{hfs}} = \vec{\nabla} \times \vec{A}_{\text{hfs}}(\vec{x}) = -\frac{2}{3} \vec{\mu} \delta(\vec{x}) - \frac{3(\vec{\mu} \cdot \hat{x})\hat{x} - \vec{\mu}}{4\pi r^3}, \quad (2)$$

and the fully relativistic hyperfine interaction Hamiltonian thus reads

$$H_{\text{hfs}} = -e\vec{\alpha} \cdot \vec{A}_{\text{hfs}}(\vec{x}) = \frac{e}{4\pi} \vec{\alpha} \cdot \frac{\vec{\mu} \times \vec{x}}{r^3} = \frac{e}{4\pi} \vec{\mu} \cdot \frac{\vec{x} \times \vec{\alpha}}{r^3}. \quad (3)$$

The hyperfine interaction couples Dirac eigenstates to the magnetic field of the nucleus. The electronic states can be written as $|njm\rangle \equiv |nj\ell m\rangle$, where n is the principal quantum number, and the orbital and total angular momenta of the electron (ℓ and j , respectively) can be mapped to the Dirac angular quantum number $\kappa = (-1)^{j-\ell+\frac{1}{2}}(j+\frac{1}{2})$. Finally, m is the projection of the total electron angular momentum onto the quantization axis. In this paper, we sometimes suppress the orbital angular momentum ℓ in the notation because we consider the coupling of the total electron angular momentum j to the nuclear spin. Nuclear states are denoted as $|IM\rangle$, where I is the nuclear spin and M is its projection onto the quantization axis. They are coupled to the electron eigenstates $|njm\rangle$ by the hyperfine interaction to form states with quantum number $|nfm_f Ij\rangle$, which are eigenstates of the total Dirac plus hyperfine Hamiltonian (f is the total electron plus nuclear angular momentum and m_f is its projection). Using Clebsch–Gordan coefficients $C_{IMjm}^{fm_f}$, the $|nfm_f Ij\rangle$ states can be written as

$$|nfm_f Ij\rangle = \sum_{M,m} C_{IMjm}^{fm_f} |IM\rangle |njm\rangle. \quad (4)$$

The hyperfine energy ΔE_{hfs} thus reads

$$E_{\text{hfs}} = \langle nfm_f Ij | H_{\text{hfs}} | nfm_f Ij \rangle. \quad (5)$$

Using the Wigner–Eckhart theorem, the hyperfine energy can be rewritten as

$$E_{\text{hfs}} = \alpha \frac{g_N}{2} \frac{m_e}{m_N} [f(f+1) - I(I+1) - j(j+1)] \times \left\langle nj \frac{1}{2} \left| \frac{[\vec{x} \times \vec{\alpha}]_0}{m_e r^3} \right| nj \frac{1}{2} \right\rangle, \quad (6)$$

where $|nj\frac{1}{2}\rangle$ is the Dirac eigenstate with a definite angular-momentum projection of $+\frac{1}{2}$, and $[\vec{x} \times \vec{\alpha}]_0$ is the z component (zero component in the spherical basis) of the indicated vector product.

We have thus separated the nuclear from the electronic variables. A detailed analysis of the separation of the nuclear variables can also be found in Ref. [2]. This procedure allows one to reduce the evaluations of the hyperfine structure, and corrections to it, to the evaluation of matrix elements of operators acting solely on electronic states. Specifically, we consider corrections to the state-dependent electronic matrix element Θ_e , where

$$\Theta_e = \left\langle nj\frac{1}{2} \left| \frac{[\vec{x} \times \vec{\alpha}]_0}{m_e r^3} \right| nj\frac{1}{2} \right\rangle. \quad (7)$$

The hyperfine interaction energy thus is

$$E_{\text{hfs}} = \alpha \frac{g_N}{2} \frac{m_e}{m_N} [f(f+1) - I(I+1) - j(j+1)] \Theta_e. \quad (8)$$

Relativistic atomic theory leads to the following result for Θ_e (see Refs. [4,9]):

$$\Theta_e = (Z\alpha)^3 m_e \frac{\kappa [2\kappa(\gamma + n - |\kappa|) - N]}{N^4 (\kappa^2 - \frac{1}{4}) \gamma (4\gamma^2 - 1)}, \quad (9)$$

where $\gamma = \sqrt{\kappa^2 - (Z\alpha)^2}$. The effective principal quantum number is $N = \sqrt{(n - |\kappa|)^2 + 2(n - |\kappa|)\gamma + \kappa^2}$.

III. LOW-ENERGY PART

A. General Formalism

In order to treat low-energy virtual photons, we apply a Foldy-Wouthuysen transformation to the total Hamiltonian H_t , which is the sum of the Dirac-Coulomb Hamiltonian and the relativistic hyperfine interaction Hamiltonian,

$$H_t = H_D + H_{\text{hfs}} = \vec{\alpha} \cdot \vec{p} + \beta m_e - \frac{Z\alpha}{r} - e \vec{\alpha} \cdot \vec{A}_{\text{hfs}}(\vec{x}). \quad (10)$$

The Foldy-Wouthuysen transformation of this Hamiltonian is carried out as described in Refs. [7,8,10–12]. The only difference to the case of the ordinary Dirac Hamiltonian is that the odd operator \mathcal{O} used in the construction of the transformation [11] now reads

$$\mathcal{O} = \vec{\alpha} \cdot \vec{p} - e \vec{\alpha} \cdot \vec{A}_{\text{hfs}}(\vec{x}), \quad (11)$$

instead of $\vec{\alpha} \cdot \vec{p}$. The result of the transformation,

$$H'_t + U H_t U^{-1} = H_{\text{FW}} + H_{\text{HFS}}, \quad (12)$$

is the sum of the Foldy-Wouthuysen Hamiltonian H_{FW} from Ref. [10], and H_{HFS} which is the nonrelativistic hyperfine splitting Hamiltonian [2,8]

$$H_{\text{HFS}} = \frac{e m_e}{4\pi} \vec{\mu} \cdot \vec{h} = \frac{e m_e}{4\pi} \vec{\mu} \cdot (\vec{h}_S + \vec{h}_D + \vec{h}_L). \quad (13)$$

It consists of the three parts

$$\vec{h}_S = \frac{4\pi}{3m_e^2} \vec{\sigma} \delta(\vec{x}), \quad (14a)$$

$$\vec{h}_D = \frac{3(\vec{\sigma} \cdot \hat{x})\hat{x} - \vec{\sigma}}{2m_e^2 r^3}, \quad (14b)$$

$$\vec{h}_L = \frac{\vec{\ell}}{m_e^2 r^3}, \quad (14c)$$

whose designation is inspired by the apparent angular-momentum association of the terms. Following the notation in Ref. [8], lowercase letters in the subscript are used to label relativistic operators, whereas nonrelativistic operators are denoted by uppercase letters in the subscript. However, we use the lowercase notation for the scaled vector quantity \vec{h} in order to denote the electronic operators in the nonrelativistic hyperfine Hamiltonian. Furthermore, $\hat{x} = \vec{x}/|\vec{x}|$ is the position unit vector. The zero component (z component) $h_0 = h_{S,0} + h_{D,0} + h_{L,0}$ of the Hamiltonian “vector” \vec{h} therefore reads as

$$h_0 = \frac{4\pi}{3m_e^2} \sigma_0 \delta(\vec{x}) + \frac{3(\vec{\sigma} \cdot \hat{x})\hat{x}_0 - \sigma_0}{2m_e^2 r^3} + \frac{\ell_0}{m_e^2 r^3}. \quad (15)$$

With the help of h_0 , the nonrelativistic limit of Eq. (9) is obtained as

$$\Theta_e^{\text{NR}} = \left\langle nj\ell\frac{1}{2} \left| h_0 \right| nj\ell\frac{1}{2} \right\rangle = \frac{\kappa}{|\kappa|} \frac{(Z\alpha)^3 m_e}{n^3 (2\kappa + 1) (\kappa^2 - \frac{1}{4})}, \quad (16)$$

which we use to define the nonrelativistic quantity

$$E_F = E_{\text{HFS}}^{\text{NR}} = E_{\text{HFS}} = \alpha \frac{g_N}{2} \frac{m_e}{m_N} \times [f(f+1) - I(I+1) - j(j+1)] \Theta_e^{\text{NR}}, \quad (17)$$

which in turn is commonly referred to as the Fermi energy. The relativistic and QED corrections can be expressed as multiplicative corrections of Θ_e , via the replacement

$$\Theta_e^{\text{NR}} \rightarrow \Theta_e^{\text{NR}} [1 + \delta\Theta_e^{\text{rel}} + \delta\Theta_e^{\text{QED}}]. \quad (18)$$

By expanding Θ_e to second order in $Z\alpha$, we obtain

$$\delta\Theta_e^{\text{rel}} = (Z\alpha)^2 \times \left[\frac{12\kappa^2 - 1}{2\kappa^2(2\kappa - 1)(2\kappa + 1)} + \frac{3}{2n} \frac{1}{|\kappa|} + \frac{3 - 8\kappa}{2n^2(2\kappa - 1)} \right] \quad (19)$$

and the corresponding energy shift

$$\delta E_{\text{hfs}}^{\text{rel}} = E_{\text{HFS}} \delta\Theta_e^{\text{rel}}. \quad (20)$$

The QED term

$$\delta E_{\text{HFS}}^{\text{QED}} = E_{\text{HFS}} \delta\Theta_e^{\text{QED}} \quad (21)$$

is the subject of this paper. For the QED corrections, terms up to relative order $\alpha(Z\alpha)^2$ with respect to the nonrelativistic hyperfine splitting will be considered. In order to do the calculation, we need the three terms from Eq. (13) and a further correction to the electron’s transition current, due to the hyperfine interaction. Namely, in the presence of the hyperfine interaction, the kinetic momentum of the electron finds a modification

$$\frac{\vec{p}}{m_e} \rightarrow \frac{\vec{p}}{m_e} - \frac{e}{m_e} \vec{A}_{\text{hfs}} = \frac{\vec{p}}{m_e} + \frac{|e|m_e}{4\pi} |\vec{\mu}| \delta \vec{J}_{\text{HFS}}. \quad (22)$$

The current

$$\delta \vec{j}_{\text{HFS}} = \frac{\hat{\mu} \times \vec{x}}{m_e^2 r^3} \quad (23)$$

has the zero component

$$\delta j_{0,\text{HFS}} = \frac{1}{m_e^2 r^3} (-y \hat{e}_x + x \hat{e}_y), \quad (24)$$

which is used in the calculations below.

B. Specific Terms

Following Ref. [8], there are four corrections that arise from the correction of the interaction current, the Hamiltonian, the reference-state energy, and the reference-state wave function.

We first treat the hyperfine correction to the interaction current and, to this end, define a useful normalization factor

$$\mathcal{N} = \frac{1}{\langle nj\ell\frac{1}{2} | h_0 | nj\ell\frac{1}{2} \rangle} = \frac{1}{\Theta_e^{\text{NR}}}. \quad (25)$$

The hyperfine correction to the interaction current is then given as

$$\begin{aligned} \delta \Theta_L^{\delta j} &= \frac{4\alpha\mathcal{N}}{3\pi} \int_0^\epsilon d\omega_{\vec{k}} \omega_{\vec{k}} \sum_{n'j'\ell'm'} \left\langle nj\ell\frac{1}{2} \left| \frac{p^i}{m_e} \right| n'j'\ell'm' \right\rangle \\ &\quad \times \frac{1}{E_n - E_{n'} - \omega_{\vec{k}}} \left\langle n'j'\ell'm' \left| \delta j_{0,\text{HFS}}^i \right| nj\ell\frac{1}{2} \right\rangle \\ &= \frac{\alpha}{\pi} (Z\alpha)^2 \frac{4\mathcal{N}}{3(Z\alpha)^2} \sum_{n'j'\ell'm'} (E_n - E_{n'}) \ln \left(\frac{|E_{n'} - E_n|}{m_e(Z\alpha)^2} \right) \\ &\quad \times \left\langle nj\ell\frac{1}{2} \left| \frac{p^i}{m_e} \right| n'j'\ell'm' \right\rangle \left\langle n'j'\ell'm' \left| \delta j_{0,\text{HFS}}^i \right| nj\ell\frac{1}{2} \right\rangle. \end{aligned} \quad (26)$$

The term containing the logarithm of ϵ , which is a scale-separation parameter that cancels when high- and low-energy parts are added [13], vanishes after angular integration in the matrix element. The structure of the logarithmic term here is very similar to the Bethe logarithm encountered in Ref. [14]. Terms of this form will arise for the other corrections in the low-energy part as well. In the following, these terms are denoted as β_{HFS} and are evaluated numerically with the methods described in Ref. [15]. Thus, the low-energy correction due to the nuclear-spin-dependent current is

$$\delta \Theta_L^{\delta j} = \frac{\alpha}{\pi} (Z\alpha)^2 \beta_{\text{HFS}}^{\delta j}. \quad (27)$$

Next, we treat the corrections to the Hamiltonian, the energy, and the wave function. The perturbation due to the hyperfine splitting Hamiltonian yields the term [we define the resolvent $G(\omega_{\vec{k}}) = 1/(E_n - H_S - \omega_{\vec{k}})$]

$$\begin{aligned} \delta \Theta_L^{\delta H} &= \frac{2\alpha\mathcal{N}}{3\pi} \int_0^\epsilon d\omega_{\vec{k}} \omega_{\vec{k}} \\ &\quad \times \left\langle nj\ell\frac{1}{2} \left| \frac{p^i}{m_e} G(\omega_{\vec{k}}) h_0 G(\omega_{\vec{k}}) \frac{p^i}{m_e} \right| nj\ell\frac{1}{2} \right\rangle \end{aligned}$$

$$\begin{aligned} &= \frac{2\alpha\mathcal{N}}{3\pi m_e^2} \ln \left[\frac{\epsilon}{m_e(Z\alpha)^2} \right] \\ &\quad \times \left\langle nj\ell\frac{1}{2} \left| \left(\frac{1}{2} \mathbf{p}^i \cdot [h_0, p^i] + p^2 h_0 \right) \right| nj\ell\frac{1}{2} \right\rangle \\ &\quad + \frac{\alpha}{\pi} (Z\alpha)^2 \beta_{\text{HFS}}^{\delta H}. \end{aligned} \quad (28)$$

The correction to the energy denominator in the Schrödinger propagator can be written as

$$\begin{aligned} \delta \Theta_L^{\delta E} &= -\frac{2\alpha\mathcal{N}}{3\pi} \int_0^\epsilon d\omega_{\vec{k}} \omega_{\vec{k}} \left\langle nj\ell\frac{1}{2} \left| h_0 \right| nj\ell\frac{1}{2} \right\rangle \\ &\quad \times \left\langle nj\ell\frac{1}{2} \left| \frac{p^i}{m_e} [G(\omega_{\vec{k}})]^2 \frac{p^i}{m_e} \right| nj\ell\frac{1}{2} \right\rangle \\ &= -\frac{2\alpha\mathcal{N}}{3\pi m_e^2} \ln \left[\frac{\epsilon}{m_e(Z\alpha)^2} \right] \left\langle nj\ell\frac{1}{2} \left| p^2 \right| nj\ell\frac{1}{2} \right\rangle \\ &\quad \times \left\langle nj\ell\frac{1}{2} \left| h_0 \right| nj\ell\frac{1}{2} \right\rangle + \frac{\alpha}{\pi} (Z\alpha)^2 \beta_{\text{HFS}}^{\delta E}. \end{aligned} \quad (29)$$

Finally, the correction to the wave function due to the hyperfine splitting Hamiltonian is

$$\begin{aligned} \delta \Theta_L^{\delta \Phi} &= \frac{4\alpha\mathcal{N}}{3\pi} \int_0^\epsilon d\omega_{\vec{k}} \omega_{\vec{k}} \\ &\quad \times \left\langle nj\ell\frac{1}{2} \left| \frac{p^i}{m_e} G(\omega_{\vec{k}}) \frac{p^i}{m_e} \left(\frac{1}{E_n - H_S} \right)' h_0 \right| nj\ell\frac{1}{2} \right\rangle \\ &= \frac{4\alpha\mathcal{N}}{3\pi m_e^2} \ln \left[\frac{\epsilon}{m_e(Z\alpha)^2} \right] \\ &\quad \times \left\langle nj\ell\frac{1}{2} \left| p^i (H_S - E_n) p^i \left(\frac{1}{E_n - H_S} \right)' h_0 \right| nj\ell\frac{1}{2} \right\rangle \\ &\quad + \frac{\alpha}{\pi} (Z\alpha)^2 \beta_{\text{HFS}}^{\delta \Phi}, \end{aligned} \quad (30)$$

where the prime indicates the reduced Green function. Using commutator relations, one can finally sum up all four corrections in the low-energy part as

$$\begin{aligned} \delta \Theta_L &= \delta \Theta_L^{\delta j} + \delta \Theta_L^{\delta H} + \delta \Theta_L^{\delta E} + \delta \Theta_L^{\delta \Phi} \\ &= \frac{\alpha\mathcal{N}}{3\pi m_e^2} \ln \left[\frac{\epsilon}{m_e(Z\alpha)^2} \right] \left\langle nj\ell\frac{1}{2} \left| \mathbf{p}^i \cdot [h_0, p^i] \right| nj\ell\frac{1}{2} \right\rangle \\ &\quad + \frac{\alpha}{\pi} (Z\alpha)^2 \beta_{\text{HFS}}, \end{aligned} \quad (31)$$

where β_{HFS} is the sum

$$\beta_{\text{HFS}} = \beta_{\text{HFS}}^{\delta j} + \beta_{\text{HFS}}^{\delta H} + \beta_{\text{HFS}}^{\delta E} + \beta_{\text{HFS}}^{\delta \Phi}. \quad (32)$$

The double commutator

$$\left\langle nj\ell\frac{1}{2} \left| \mathbf{p}^i \cdot [h_0, p^i] \right| nj\ell\frac{1}{2} \right\rangle = \left\langle nj\ell\frac{1}{2} \left| \vec{\nabla}^2 h_0 \right| nj\ell\frac{1}{2} \right\rangle \quad (33)$$

vanishes for states with $\ell \geq 2$ up to and including order $(Z\alpha)^5$, and hence $\delta \Theta_L$ takes the very simple form

$$\delta \Theta_L = \frac{\alpha}{\pi} (Z\alpha)^2 \beta_{\text{HFS}}. \quad (34)$$

IV. HIGH-ENERGY PART

Up to relative order $\alpha(Z\alpha)^2 E_F$, it is sufficient [8] to consider the problem on the level of the modified Dirac Hamiltonian,

$$H_D^{(m)} = \vec{\alpha}[\vec{p} - eF_1(\vec{\nabla}^2)\vec{A}] + \beta m_e + F_1(\vec{\nabla}^2)V + F_2(\vec{\nabla}^2)\frac{e}{2m_e}(i\vec{\gamma} \cdot \vec{E} - \beta \vec{\Sigma} \cdot \vec{B}), \quad (35)$$

where F_1 and F_2 are the one-loop Dirac and Pauli form factors of the electron, respectively. Their expressions are known (see Chapter 7 of Ref. [16]).

The F_1 form-factor slope gives rise to the following effective interaction:

$$-eF_1'(0)\vec{\nabla}^2\vec{\alpha} \cdot \vec{A}_{\text{hfs}} = \frac{\alpha}{3\pi} \left[\ln\left(\frac{m_e}{2\epsilon}\right) + \frac{11}{24} \right] \vec{\nabla}^2 H_{\text{hfs}}. \quad (36)$$

Up to the order $\alpha(Z\alpha)^2 E_F$, we may write the correction in terms of the nonrelativistic hyperfine Hamiltonian h_0 ,

$$\delta\Theta_{H,1} = \frac{\alpha\mathcal{N}}{3\pi m_e^2} \left[\ln\left(\frac{m_e}{2\epsilon}\right) + \frac{11}{24} \right] \left\langle nj\ell\frac{1}{2} \left| \vec{\nabla}^2 h_0 \right| nj\ell\frac{1}{2} \right\rangle. \quad (37)$$

However, as already pointed out, the matrix element of $\vec{\nabla}^2 h_0$ vanishes on states with $\ell \geq 2$, which are relevant to our investigations, and so

$$\delta\Theta_{H,1} = 0. \quad (38)$$

The second correction is a second-order perturbation involving the F_1 correction to the Coulomb potential,

$$\delta\Theta_{H,2} = \frac{2\alpha\mathcal{N}}{3\pi m_e^2} \left[\ln\left(\frac{m_e}{2\epsilon}\right) + \frac{11}{24} \right] \times \left\langle nj\ell\frac{1}{2} \left| \vec{\nabla}^2 V \left(\frac{1}{E_n - H_S} \right)' h_0 \right| nj\ell\frac{1}{2} \right\rangle. \quad (39)$$

Again, $\vec{\nabla}^2 V$ is proportional to the Dirac δ and therefore vanishes for states with $\ell \geq 1$. Accordingly, for states with $\ell \geq 2$, we have

$$\delta\Theta_{H,2} = 0. \quad (40)$$

The Pauli F_2 form factor gives rise to a second-order perturbation involving a magnetic-moment correction to the Coulomb potential,

$$\delta\Theta_{H,3} = 2\mathcal{N} F_2(0) \times \left\langle \psi^\dagger \left| \frac{-i}{2m_e} \vec{\gamma} \cdot \vec{\nabla} V \left(\frac{1}{E_\psi - H_D} \right)' H_{\text{hfs}} \right| \psi \right\rangle, \quad (41)$$

where F_2 is the magnetic form factor. For $F_2(0)$, the Schwinger value $F_2(0) = \frac{\alpha}{2\pi}$ may be used. After a Foldy–Wouthuysen transformation, we can write $\delta\Theta_{H,3}$ as the sum of two terms. The first of these, $\delta\Theta_{H,3n}$, involves no mixing of upper and lower components in the Dirac wave function and reads

$$\delta\Theta_{H,3n} = \frac{\alpha\mathcal{N}}{2\pi m_e^2} \times \left\langle nj\ell\frac{1}{2} \left| \frac{Z\alpha}{r^3} \vec{\sigma} \cdot \vec{\ell} \left(\frac{1}{E_n - H_S} \right)' H_{\text{HFS}} \right| nj\ell\frac{1}{2} \right\rangle. \quad (42)$$

We find the following general result for states with $\ell \geq 2$:

$$\delta\Theta_{H,3n} = \frac{\alpha}{\pi} (Z\alpha)^2 \left(\frac{1}{2\ell\kappa} \frac{60\ell^4 + 120\ell^3 + 55\ell^2 - 5\ell - 3}{(2\ell + 1)^2 (4\ell^3 + 8\ell^2 + \ell - 3)} + \frac{3}{2n} \frac{1}{\kappa(2\ell + 1)} - \frac{3}{8n^2} \frac{\ell(\ell + 1)}{\kappa(\ell + \frac{3}{2})(\ell - \frac{1}{2})} \right). \quad (43)$$

Lower components of the Dirac wave function give rise to the mixing term

$$\delta\Theta_{H,3m} = -i \frac{\alpha\mathcal{N}}{2\pi m_e} \left\langle nj\ell\frac{1}{2} \left| \frac{Z\alpha}{r^3} (\vec{\gamma} \cdot \vec{x}) \frac{1}{2m_e} H_{\text{hfs}} \right| nj\ell\frac{1}{2} \right\rangle = \frac{\alpha\mathcal{N}}{\pi} \frac{-\kappa}{8j(j+1)} \left\langle nj\ell\frac{1}{2} \left| \frac{Z\alpha}{m_e^3 r^4} \right| nj\ell\frac{1}{2} \right\rangle. \quad (44)$$

We find the general result

$$\delta\Theta_{H,3m} = \frac{\alpha}{\pi} (Z\alpha)^2 \frac{|\kappa|}{4j(j+1)} \frac{(2\kappa + 1)(\kappa^2 - \frac{1}{4})}{(2\ell - 1)(2\ell + 3)(\ell + \frac{1}{2})} \times \left[\frac{1}{n^2} - \frac{3}{\ell(\ell + 1)} \right]. \quad (45)$$

The F_2 correction to the magnetic photon exchange of electron and nucleus gives rise to the effective interaction

$$-F_2(\vec{\nabla}^2) \frac{e}{2m_e} \beta \vec{\Sigma} \cdot \vec{B}_{\text{hfs}} = F_2(\vec{\nabla}^2) \left[\frac{em_e}{4\pi} \beta \vec{\mu} \cdot (\vec{h}_s + \vec{h}_d) \right]. \quad (46)$$

Here, \vec{h}_s and \vec{h}_d are the generalizations of \vec{h}_S and \vec{h}_D to 4×4 matrices,

$$\vec{h}_s = \frac{4\pi}{3m_e^2} \vec{\Sigma} \delta(\vec{x}), \quad (47a)$$

$$\vec{h}_d = \frac{3(\vec{\Sigma} \cdot \hat{x}) \hat{x} - \vec{\Sigma}}{2m_e^2 r^3}. \quad (47b)$$

Taking $F_2(\vec{\nabla}^2) \approx F_2(0)$ in Eq. (46), we obtain the correction

$$\delta\Theta_{H,4} = \mathcal{N} F_2(0) \langle \psi | \beta (h_{s,0} + h_{d,0}) | \psi \rangle = \frac{\alpha\mathcal{N}}{2\pi} \langle \psi | \beta (h_{s,0} + h_{d,0}) | \psi \rangle. \quad (48)$$

Generalizing results from Ref. [17] for the term of relative order α , we find the result

$$\delta\Theta_{H,4} = \frac{\alpha}{\pi} \left\{ \frac{1}{4\kappa} + (Z\alpha)^2 \left[\frac{24\kappa^3 + 18\kappa^2 - \kappa - 1}{8\kappa^3(4\kappa^3 + 4\kappa^2 - \kappa - 1)} + \frac{3}{8n} \frac{1}{|\kappa|\kappa} + \frac{1}{n^2} \frac{1}{2\kappa} \frac{1 - 3\kappa}{2\kappa - 1} \right] \right\}. \quad (49)$$

Taking the slope of F_2 in Eq. (46), we obtain

$$F_2'(0) \frac{e}{2m_e} \beta \vec{\nabla}^2 \vec{\Sigma} \cdot \vec{B}_{\text{hfs}} = \frac{\alpha}{12\pi} \left\{ \frac{em_e}{4\pi} \beta \vec{\mu} \cdot [\vec{\nabla}^2 (\vec{h}_s + \vec{h}_d)] \right\}, \quad (50)$$

with $F_2'(0) = \alpha/12\pi$. As $F_2'(0)\vec{\nabla}^2$ already is of relative order $\alpha(Z\alpha)^2$, this operator only has to be applied to the

nonrelativistic wave function where it vanishes for states with $\ell \geq 2$, and thus

$$\delta\Theta_{H,5} = \frac{\alpha\mathcal{N}}{12\pi} \left\langle nj\ell \frac{1}{2} \left| \vec{\nabla}^2 (h_{S,0} + h_{D,0}) \right| nj\ell \frac{1}{2} \right\rangle = 0. \quad (51)$$

Finally, since

$$\delta\Theta_{H,1} = \delta\Theta_{H,2} = \delta\Theta_{H,5} = 0, \quad (52)$$

we have, for the high-energy part, the result

$$\delta\Theta_H = \delta\Theta_{H,3n} + \delta\Theta_{H,3m} + \delta\Theta_{H,4}. \quad (53)$$

It is quite surprising that the result obtained by adding the above expressions,

$$\begin{aligned} \delta\Theta_H = & \frac{\alpha}{\pi} \left\{ \frac{1}{4\kappa} + (Z\alpha)^2 \left[\frac{1}{8\kappa^3} \frac{24\kappa^3 + 18\kappa^2 - \kappa - 1}{4\kappa^3 + 4\kappa^2 - \kappa - 1} \right. \right. \\ & + \frac{1}{2\ell\kappa} \frac{60\ell^4 + 120\ell^3 + 55\ell^2 - 5\ell - 3}{(2\ell + 1)^2 (4\ell^3 + 8\ell^2 + \ell - 3)} \\ & - \frac{3}{\ell(\ell + 1)} \frac{j + \frac{1}{2}}{4j(j + 1)} \frac{(2\kappa + 1)(\kappa^2 - \frac{1}{4})}{(2\ell - 1)(2\ell + 3)(\ell + \frac{1}{2})} \\ & + \frac{1}{n} \frac{3}{8\kappa} \frac{4(j + \frac{1}{2}) + (2\ell + 1)}{(2\ell + 1)(j + \frac{1}{2})} + \frac{1}{8n^2} \left(\frac{4}{1 - 2\kappa} - \frac{4}{\kappa} \right. \\ & \left. \left. + \frac{(2j + 1)(2\kappa - 1)(2\kappa + 1)^2}{2j(j + 1)(2\ell + 3)(4\ell^2 - 1)} - \frac{3\ell(\ell + 1)}{\kappa(\ell + \frac{3}{2})(\ell - \frac{1}{2})} \right) \right\}, \end{aligned} \quad (54)$$

can actually be simplified quite considerably as

$$\begin{aligned} \delta\Theta_H = & \frac{\alpha}{\pi} \left\{ \frac{1}{4\kappa} + (Z\alpha)^2 \left[\frac{1}{n} \frac{3}{8} \frac{\kappa}{|\kappa|} \frac{6\kappa + 1}{\kappa^2(2\kappa + 1)} + \frac{1}{n^2} \right. \right. \\ & \left. \left. \times \frac{4\kappa - 1}{2\kappa(1 - 2\kappa)} + \frac{(4\kappa + 1)(6\kappa + 1)(6\kappa^2 + 3\kappa - 1)}{8\kappa^3(2\kappa + 1)^2(2\kappa - 1)(\kappa + 1)} \right] \right\}. \end{aligned} \quad (55)$$

V. RESULTS AND PREDICTIONS

The total self-energy correction to the hyperfine splitting is obtained as the sum of the high- and low-energy parts given in Eqs. (55) and (34), and reads

$$\begin{aligned} \delta\Theta = & \frac{\alpha}{\pi} \left\{ \frac{1}{4\kappa} + (Z\alpha)^2 \left[\frac{1}{8\kappa^3} \frac{(4\kappa + 1)(6\kappa + 1)(6\kappa^2 + 3\kappa - 1)}{(2\kappa + 1)^2(2\kappa - 1)(\kappa + 1)} \right. \right. \\ & \left. \left. + \frac{1}{n} \frac{3}{8} \frac{\kappa}{|\kappa|} \frac{6\kappa + 1}{\kappa^2(2\kappa + 1)} + \frac{1}{n^2} \frac{4\kappa - 1}{2\kappa(1 - 2\kappa)} + \beta_{\text{HFS}} \right] \right\}. \end{aligned} \quad (56)$$

TABLE I. Low-energy contribution β_{HFS} of the self-energy correction for the hyperfine splitting for D states ($\ell = 2$). The numbers in parentheses are standard uncertainties in the last figure.

n	$\beta_{\text{HFS}}(nD_{3/2})$	$\beta_{\text{HFS}}(nD_{5/2})$
3	$-2.068\,39(5) \times 10^{-2}$	$-3.455\,22(5) \times 10^{-2}$
4	$-1.302\,99(5) \times 10^{-2}$	$-3.793\,94(5) \times 10^{-2}$
5	$-1.025\,92(5) \times 10^{-2}$	$-3.965\,95(5) \times 10^{-2}$
6	$-0.943\,28(5) \times 10^{-2}$	$-4.084\,82(5) \times 10^{-2}$

TABLE II. Low-energy contribution β_{HFS} of the self-energy correction for the hyperfine splitting for F states ($\ell = 3$). The numbers in parentheses are standard uncertainties in the last figure.

n	$\beta_{\text{HFS}}(nF_{5/2})$	$\beta_{\text{HFS}}(nF_{7/2})$
4	$-1.021\,46(5) \times 10^{-2}$	$-0.953\,04(5) \times 10^{-2}$
5	$-0.683\,94(5) \times 10^{-2}$	$-1.077\,62(5) \times 10^{-2}$
6	$-0.504\,64(5) \times 10^{-2}$	$-1.141\,28(5) \times 10^{-2}$
7	$-0.407\,32(5) \times 10^{-2}$	$-1.182\,43(5) \times 10^{-2}$

The Bethe-logarithm-type correction β_{HFS} is given in Eq. (32).

Restoring the reduced-mass dependence [we define $r(\mathcal{N}) \equiv m_e/m_{\mathcal{N}}$] and adding the relativistic correction of relative order $(Z\alpha)^2$, we find that

$$\begin{aligned} v_{\text{hfs}} = & R_{\infty} c \frac{Z^3 \alpha^2}{n^3} \frac{r(\mathcal{N})}{[1 + r(\mathcal{N})]^3} \frac{\kappa}{|\kappa|} \frac{g_{\mathcal{N}}}{(2\kappa + 1)(\kappa^2 - \frac{1}{4})} \\ & \times [f(f + 1) - I(I + 1) - j(j + 1)] \left\{ 1 + (Z\alpha)^2 \right. \\ & \times \left[\frac{12\kappa^2 - 1}{2\kappa^2(2\kappa - 1)(2\kappa + 1)} + \frac{3}{2n} \frac{1}{|\kappa|} + \frac{3 - 8\kappa}{2n^2(2\kappa - 1)} \right] \\ & + \frac{\alpha}{\pi} \frac{1}{4\kappa} + \frac{\alpha}{\pi} (Z\alpha)^2 \left[\frac{3}{8n} \frac{\kappa}{|\kappa|} \frac{6\kappa + 1}{\kappa^2(2\kappa + 1)} + \frac{1}{n^2} \frac{4\kappa - 1}{2\kappa(1 - 2\kappa)} \right. \\ & \left. \left. + \frac{1}{8\kappa^3} \frac{(4\kappa + 1)(6\kappa + 1)(6\kappa^2 + 3\kappa - 1)}{(2\kappa + 1)^2(2\kappa - 1)(\kappa + 1)} + \beta_{\text{HFS}} \right] \right\}. \end{aligned} \quad (57)$$

Numerical data for β_{HFS} for D , F , and G states, and selected Rydberg states, can be found in Tables I, II, III, and IV, respectively. Those in Table IV are relevant for a possible determination of fundamental constants from Rydberg-state spectroscopy in hydrogenlike ions of medium nuclear-charge number (see Ref. [18]).

The following is a numerical example: For the transition $|1\rangle \leftrightarrow |2\rangle$ in atomic hydrogen ($Z = 1$), where

$$|1\rangle = |n = 15, \ell = 14, j = 29/2, f = 15\rangle, \quad (58)$$

$$|2\rangle = |n = 16, \ell = 15, j = 31/2, f = 16\rangle, \quad (59)$$

we find the following frequency shift from the Dirac value indicated in Eq. (2) of Ref. [17],

$$\Delta v_{\text{hfs},1 \rightarrow 2} = 96.759\,863\,0(8) \text{ Hz} + 78.764\,693(13) \text{ Hz}, \quad (60)$$

where the first term is due to the hyperfine effects calculated here, and the second term is due to relativistic recoil and QED

TABLE III. Low-energy contribution β_{HFS} of the self-energy correction for the hyperfine splitting for G states ($\ell = 4$). The numbers in parentheses are standard uncertainties in the last figure.

n	$\beta_{\text{HFS}}(nG_{7/2})$	$\beta_{\text{HFS}}(nG_{9/2})$
5	$-0.368\,23(5) \times 10^{-2}$	$-0.341\,76(5) \times 10^{-2}$
6	$-0.147\,39(5) \times 10^{-2}$	$-0.388\,58(5) \times 10^{-2}$
7	$-0.012\,93(5) \times 10^{-2}$	$-0.414\,24(5) \times 10^{-2}$
8	$0.072\,16(5) \times 10^{-2}$	$-0.430\,98(5) \times 10^{-2}$

TABLE IV. Low-energy contribution β_{HFS} of the self-energy correction for the hyperfine splitting for highly excited states. The numbers in parentheses are standard uncertainties in the last figure. The exponent of the numerical data is chosen to be the same as in Tables I–III, illustrating the monotonic decrease of the coefficients with the angular-momentum quantum numbers.

n	ℓ	$2j$	κ	β_{HFS}	$2j$	κ	β_{HFS}
16	15	29	15	$0.006\,310(5) \times 10^{-2}$	31	−16	$-0.002\,130(5) \times 10^{-2}$
16	14	27	14	$0.016\,397(5) \times 10^{-2}$	29	−15	$-0.003\,041(5) \times 10^{-2}$
15	14	27	14	$0.006\,888(5) \times 10^{-2}$	29	−15	$-0.002\,795(5) \times 10^{-2}$
15	13	25	13	$0.019\,372(5) \times 10^{-2}$	27	−14	$-0.004\,086(5) \times 10^{-2}$
14	13	25	13	$0.007\,420(5) \times 10^{-2}$	27	−14	$-0.003\,741(5) \times 10^{-2}$
14	12	23	12	$0.023\,029(5) \times 10^{-2}$	25	−13	$-0.005\,617(5) \times 10^{-2}$
13	12	23	12	$0.007\,794(5) \times 10^{-2}$	25	−13	$-0.005\,122(5) \times 10^{-2}$

effects calculated in Ref. [18]. The final theoretical prediction for the shift from the Dirac value

$$\Delta\nu_{\text{hfs},1\rightarrow 2} = 175.524\,556(13) \text{ Hz}, \quad (61)$$

where the fundamental constants of the Committee on Data for Science and Technology (CODATA) 2006 [19] have been used in the numerical evaluation. The next higher-order term neglected here is the recoil correction of relative order $(Z\alpha)^2 r(\mathcal{N})$, for which a general expression has been derived in Ref. [20] (the corresponding expression also is given in Eq. (42) of Ref. [4]). The recoil correction is numerically suppressed for $Z = 1$.

VI. CONCLUSIONS

Rydberg states of hydrogenlike ions with medium nuclear-charge number have been proposed as a device for the determination of fundamental constants [18]. Here, we demonstrate that it is possible to obtain accurate theoretical predictions for transition frequencies, even in cases where the nucleus carries spin. To this end, we calculate the self-energy correction to the hyperfine splitting of the high-lying states. Vacuum polarization effects can be neglected for states with $\ell \geq 2$ to the order relevant for the current investigation.

We split the calculation into a low-energy part, which contains Bethe-logarithm-type corrections (Sec. III), and a

high-energy part, which can be treated on the basis of electron form factors (Sec. IV). For the low-energy part, we find that the net result can be expressed as the sum of corrections due to the hyperfine Hamiltonian, the energy correction, the wave-function correction, and the hyperfine modification of the electron's transition current. For the high-energy part, we find a sum of two terms, one of which is due to a second-order effect involving the Pauli form-factor correction to the Coulomb field, and the second of which is an anomalous-magnetic-moment correction to the hyperfine splitting, evaluated on relativistic wave functions. The first correction can be split into two terms, which involve and do not involve mixing of the upper and lower components of the Dirac wave function, respectively. Quite surprisingly, the high-energy contribution can be expressed in closed analytic form, valid for an arbitrary excited state [see Eq. (55)]. For the Bethe-logarithm-type corrections relevant to the low-energy part, a numerical approach is indispensable.

Finally, as indicated in Tables I–III, we also find results for D , F , and G states, which are of general interest to high-precision spectroscopy.

ACKNOWLEDGMENTS

This research has been supported by the National Science Foundation and by a Precision Measurement Grant from the National Institute of Standards and Technology.

-
- [1] W. E. Caswell and G. P. Lepage, *Phys. Lett. B* **167**, 437 (1986).
[2] S. J. Brodsky and R. G. Parsons, *Phys. Rev.* **163**, 134 (1967).
[3] V. A. Yerokhin, A. N. Artemyev, V. M. Shabaev, and G. Plunien, *Phys. Rev. A* **72**, 052510 (2005).
[4] U. D. Jentschura and V. A. Yerokhin, *Phys. Rev. A* **73**, 062503 (2006).
[5] V. A. Yerokhin and U. D. Jentschura, *Phys. Rev. Lett.* **100**, 163001 (2008).
[6] V. A. Yerokhin and U. D. Jentschura, *Phys. Rev. A* **81**, 012502 (2010).
[7] D. Zwanziger, *Phys. Rev.* **121**, 1128 (1961).
[8] U. D. Jentschura and V. A. Yerokhin, *Phys. Rev. A* **81**, 012503 (2010).
[9] P. Pyykkö, E. Pajanne, and M. Inokuti, *Int. J. Quantum Chem.* **7**, 785 (1973).
[10] L. L. Foldy and S. A. Wouthuysen, *Phys. Rev.* **78**, 29 (1950).
[11] J. D. Bjorken and S. D. Drell, *Relativistische Quantenmechanik* (Bibliographisches Institut, Mannheim/Wien/Zürich, 1966).
[12] K. Pachucki, *Phys. Rev. A* **69**, 052502 (2004).
[13] K. Pachucki, *Ann. Phys. NY* **226**, 1 (1993).
[14] H. A. Bethe, *Phys. Rev.* **72**, 339 (1947).
[15] S. Salomonson and P. Öster, *Phys. Rev. A* **40**, 5559 (1989).
[16] C. Itzykson and J. B. Zuber, *Quantum Field Theory* (McGraw-Hill, New York, 1980).
[17] B. J. Wundt and U. D. Jentschura, *J. Phys. B* **43**, 115002 (2010).
[18] U. D. Jentschura, P. J. Mohr, J. N. Tan, and B. J. Wundt, *Phys. Rev. Lett.* **100**, 160404 (2008).
[19] P. J. Mohr, B. N. Taylor, and D. B. Newell, *Rev. Mod. Phys.* **80**, 633 (2008).
[20] M. M. Sternheim, *Phys. Rev.* **130**, 211 (1963).