

---

01 Aug 2012

## Three-Body Dynamics in Single Ionization of Atomic Hydrogen by 75 keV Proton Impact

Aaron LaForge

Kisra Egodapitiya

Sachin Sharma

Michael Schulz

Missouri University of Science and Technology, [schulz@mst.edu](mailto:schulz@mst.edu)

Follow this and additional works at: [https://scholarsmine.mst.edu/phys\\_facwork](https://scholarsmine.mst.edu/phys_facwork)

 Part of the [Physics Commons](#)

---

### Recommended Citation

A. LaForge et al., "Three-Body Dynamics in Single Ionization of Atomic Hydrogen by 75 keV Proton Impact," *Proceedings of the 27th International Conference on Photonic, Electronic and Atomic Collisions (2011, Belfast)*, vol. 388, no. PART 8, Institute of Physics - IOP Publishing, Aug 2012.

The definitive version is available at <https://doi.org/10.1088/1742-6596/388/8/082044>

This Article - Conference proceedings is brought to you for free and open access by Scholars' Mine. It has been accepted for inclusion in Physics Faculty Research & Creative Works by an authorized administrator of Scholars' Mine. This work is protected by U. S. Copyright Law. Unauthorized use including reproduction for redistribution requires the permission of the copyright holder. For more information, please contact [scholarsmine@mst.edu](mailto:scholarsmine@mst.edu).

## Three-Body Dynamics in Single Ionization of Atomic Hydrogen by 75 keV Proton Impact

Aaron LaForge\*<sup>1</sup>, Kisra Egodapitiya\*, Sachin Sharma\*, and Michael Schulz\*

\* School of Physics, Missouri University of Science and Technology, Rolla, Mo USA

**Synopsis** We have measured and calculated doubly differential cross sections for ionization of atomic hydrogen using 75-keV proton impact for fixed projectile energy losses as a function of scattering angle.

We have measured and calculated doubly differential cross sections for ionization of atomic hydrogen using 75-keV proton impact for fixed projectile energy losses as a function of scattering angle. This collision system represents a pure three-body system and thus offers an accurate test of the theoretical description of the few-body dynamics without any complications presented by electron correlation in many-electron targets. Comparison between experiment and several theoretical models reveals that the projectile-target nucleus interaction is best described by the operator of a second-order term of the transition amplitude. Higher-order contributions in the projectile-electron interaction, on the other hand, are more appropriately accounted for in the final-state wave function.

---

<sup>1</sup> E-mail: [laforge@mpi-hd.mpg.de](mailto:laforge@mpi-hd.mpg.de)