


01 Jan 1971

Existence And Uniqueness For Nonlinear Neutral-differential Equations

L. J. Grimm

Missouri University of Science and Technology

Follow this and additional works at: https://scholarsmine.mst.edu/math_stat_facwork

 Part of the [Mathematics Commons](#), and the [Statistics and Probability Commons](#)

Recommended Citation

L. J. Grimm, "Existence And Uniqueness For Nonlinear Neutral-differential Equations," *Bulletin of the American Mathematical Society*, vol. 77, no. 3, pp. 374 - 376, American Mathematical Society, Jan 1971. The definitive version is available at <https://doi.org/10.1090/S0002-9904-1971-12701-5>

This Article - Journal is brought to you for free and open access by Scholars' Mine. It has been accepted for inclusion in Mathematics and Statistics Faculty Research & Creative Works by an authorized administrator of Scholars' Mine. This work is protected by U. S. Copyright Law. Unauthorized use including reproduction for redistribution requires the permission of the copyright holder. For more information, please contact scholarsmine@mst.edu.

EXISTENCE AND UNIQUENESS FOR NONLINEAR NEUTRAL-DIFFERENTIAL EQUATIONS¹

BY L. J. GRIMM

Communicated by Wolfgang Wasow, August 14, 1970

ABSTRACT. Fixed point theorems are used to prove existence and uniqueness of the C^1 solution of the initial-value problem for a functional-differential equation of neutral type.

1. Introduction. In this paper we consider the initial-value problem (IVP) for the functional-differential equation of neutral type

$$(1) \quad x'(t) = f(t, x(t), x(g(t, x(t))), x'(h(t, x(t)))),$$

with the initial condition

$$(2a) \quad x(0) = x_0.$$

Here $f(t, x, y, z)$, $g(t, x)$ and $h(t, x)$ are continuous functions with $g(0, x_0) = h(0, x_0) = 0$. We assume further that the algebraic equation $z = f(0, x_0, x_0, z)$ has a real root z_0 , and we require that

$$(2b) \quad x'(0) = z_0.$$

Existence theorems for IVP's for equation (1) have been proved by R. D. Driver [1] for the case where $h(t, x) < t$, and recently by V. P. Skripnik [2] under the hypotheses that f is sufficiently small, $h(t, x)$ is independent of x , and f is linear in the argument $x'(h(t))$. Our existence theorem requires none of these hypotheses. Under some additional conditions we obtain a local uniqueness theorem, and obtain as a corollary a result on existence of continuous solutions of certain nonlinear functional equations.

2. Existence. Let $\alpha > 0$ and let $J = [-\alpha, \alpha]$. We shall make the following assumptions concerning the IVP (1)–(2a)–(2b):

(i) $f(t, x, y, z)$ is continuous in some region in R^4 containing

$$P = \{(t, x, y, z) : |t| \leq \alpha, |x - x_0| \leq \beta, |y - x_0| \leq \beta, |z| \leq M\}$$

where α , β and $M > |z_0|$ are positive constants. We assume that $\alpha \leq \beta/M$ and that $\sup_{(t, x, y, z) \in P} |f(t, x, y, z)| \leq M$.

AMS 1970 subject classifications. Primary 34K05; Secondary 34K05.

Key words and phrases. Neutral-differential equations, functional differential equations.

¹ Research supported by NSF Grant GP 20194.

(ii) $g(t, x)$ and $h(t, x)$ are continuous in the projection \tilde{R} of P into the (t, x) space; g and h both map \tilde{R} into J , with $g(0, x_0) = h(0, x_0) = 0$, and $h(t, x)$ satisfies the Lipschitz condition

$$|h(t_1, x_1) - h(t_2, x_2)| \leq k_1 |t_1 - t_2| + k_2 |x_1 - x_2|$$

for all $(t_1, x_1), (t_2, x_2) \in R$, where k_1 and k_2 are nonnegative constants with $k_1 + k_2 M \leq 1$.

(iii) The function $f(t, x, y, z)$ satisfies the Lipschitz condition

$$|f(t, x, y, z_1) - f(t, x, y, z_2)| \leq L_z |z_1 - z_2|$$

for all $(t, x, y, z_1), (t, x, y, z_2) \in P$, where $L_z < 1$.

The Schauder fixed-point theorem yields

THEOREM 1. *Under the hypotheses (i)–(iii), the IVP (1)–(2a)–(2b) has at least one solution which is continuously differentiable on J .*

3. Uniqueness. In case $h(t, x)$ is independent of x , we obtain the following theorem:

THEOREM 2. *In addition to the hypotheses of Theorem 1, suppose that:*

(iv) $h(t, x)$ is independent of x ;

(v) f and g satisfy the Lipschitz conditions:

$$|f(t, x_1, y_1, z_1) - f(t, x_2, y_2, z_2)| \leq L \{ |x_1 - x_2| + |y_1 - y_2| \} + L_z |z_1 - z_2|$$

where L and L_z are nonnegative constants, with $L_z < 1$;

$$|g(t, x_1) - g(t, x_2)| \leq L_g |x_1 - x_2|$$

with L_g a nonnegative constant, uniformly in their respective domains.

Then there exists γ_0 , $0 < \gamma_0 \leq \alpha$, such that there is a unique continuously differentiable solution of the IVP (1)–(2a)–(2b) on the interval $[-\gamma_0, \gamma_0]$.

The proof follows from the contraction mapping principle.

4. Nonlinear functional equations. As a corollary to our existence and uniqueness results, we note that if $f(t, x, y, z)$ is independent of x and y , and $h(t, x)$ is independent of x , the problem (1)–(2b) has the form of the functional equation

$$(3) \quad z(t) = f(t, z(h(t))),$$

$$(4) \quad z(0) = z_0,$$

where z_0 is a root of $z = f(0, z)$. Theorems 1 and 2 then yield at once:

THEOREM 3. *Let $f(t, z)$ be continuous in some region in R^2 containing $P_1 = \{t: |t| \leq \alpha, |z| \leq M\}$, where α and M are positive constants such that $\sup_{(t,z) \in P_1} |f(t, z)| < M$, and $M > |z_0|$ where z_0 is a real root of $z = f(0, z)$. Let f satisfy the Lipschitz condition $|f(t, z_1) - f(t, z_2)| \leq L_z |z_1 - z_2|$ for all $(t, z_1), (t, z_2) \in P_1$, with $L_z < 1$. Let $h(t)$ be continuous for $|t| \leq \alpha$, $h(0) = 0$, and $|h(t_1) - h(t_2)| \leq |t_1 - t_2|$ for $t_1, t_2 \in [-\alpha, \alpha]$.*

Then the problem (3)–(4) has at least one continuous solution on $[-\alpha, \alpha]$, and this is the unique continuous solution on this interval if α is sufficiently small.

REFERENCES

1. R. D. Driver, *A functional-differential system of neutral type arising in a two-body problem of classical electrodynamics*, Internat. Sympos. Nonlinear Differential Equations and Nonlinear Mechanics, New York, Academic Press, 1963, pp. 474–484. MR 26 #4008.
2. V. P. Skripnik, *On some systems with deviating argument of neutral type*, Izv. Vysš. Učebn. Zaved. Matematika 1968, no. 8 (75), 80–87. (Russian)

UNIVERSITY OF MISSOURI-ROLLA, ROLLA, MISSOURI 65401