

01 Jan 1981

## Integral Identities Of Norms And Characterization Of Inner Product Spaces

Časlav V. Stanojević  
*Missouri University of Science and Technology*

Ana M. Suchanek

Follow this and additional works at: [https://scholarsmine.mst.edu/math\\_stat\\_facwork](https://scholarsmine.mst.edu/math_stat_facwork)



Part of the [Mathematics Commons](#), and the [Statistics and Probability Commons](#)

---

### Recommended Citation

Č. V. Stanojević and A. M. Suchanek, "Integral Identities Of Norms And Characterization Of Inner Product Spaces," *Proceedings of the American Mathematical Society*, vol. 81, no. 1, pp. 101 - 103, American Mathematical Society, Jan 1981.

The definitive version is available at <https://doi.org/10.1090/S0002-9939-1981-0589146-3>

This Article - Journal is brought to you for free and open access by Scholars' Mine. It has been accepted for inclusion in Mathematics and Statistics Faculty Research & Creative Works by an authorized administrator of Scholars' Mine. This work is protected by U. S. Copyright Law. Unauthorized use including reproduction for redistribution requires the permission of the copyright holder. For more information, please contact [scholarsmine@mst.edu](mailto:scholarsmine@mst.edu).

## INTEGRAL IDENTITIES OF NORMS AND CHARACTERIZATION OF INNER PRODUCT SPACES

ČASLAV V. STANOJEVIĆ AND ANA M. SUCHANEK

**ABSTRACT.** Integral identities of norms over compact groups are used to characterize inner product spaces. These integral identities also generalize the Penico-Stanojević [1] integral form of the parallelogram law.

1. Recently Penico and Stanojević [1] proved that a normed complex linear space  $N$  is an inner product space if and only if

$$\frac{1}{2\pi} \int_0^{2\pi} \|(\cos t)x + (\sin t)y\|^2 dt < 1,$$

for all unit vectors  $x$  and  $y$  in  $N$ . This characterization is an integral analogue to the Schoenberg [2] version of the classical M. M. Day [3] result.

In this paper we shall consider a more general type of integral of norms than the one in the prior inequality, and show that certain integral identities characterize inner product spaces among normed linear spaces.

2. Throughout the paper  $N$  will denote a normed complex linear space,  $C$  the field of complex numbers and  $\Gamma = \{z \mid |z| = 1\}$ , the unit circle of  $C$ .

In the next theorem we shall give necessary and sufficient conditions for  $N$  to be an inner product space in terms of norm integrals over compact groups.

**THEOREM.** *Let  $G$  be a compact topological group with normalized Haar measure  $\mu$ , and let  $\gamma \neq 1$  be a continuous homomorphism of  $G$  into  $\Gamma$ . Then  $N$  is an inner product space if and only if*

$$\int_G \|x + \gamma(g)y\|^2 d\mu = \|x\|^2 + \|y\|^2 \tag{*}$$

for all  $x, y \in N$ .

**PROOF.** The "only if" part. Notice that from

$$\int_G \gamma(g) d\mu = \int_G \gamma(ag) d\mu = \gamma(a) \int_G \gamma(g) d\mu$$

it follows that

$$\int_G \gamma(g) d\mu = 0 \tag{†}$$

for  $\gamma \neq 1$ .

Received by the editors January 8, 1980.

AMS (MOS) subject classifications (1970). Primary 46C10.

Key words and phrases. Inner product spaces.

Assume that  $N$  is an inner product space. Then

$$\int_G \|x + \gamma(g)y\|^2 d\mu = \|x\|^2 + \|y\|^2 + \int_G \operatorname{Re}[\overline{\gamma(g)}(x, y)] d\mu,$$

where  $(\cdot, \cdot)$  is the inner product in  $N$ . The integral on the right-hand side is zero because of  $(\dagger)$ . Hence  $(*)$  follows.

The "if" part. Any complex two-dimensional subspace  $N_2$  of  $N$  can be considered as a real four-dimensional space, and the unit sphere  $S_2$  of  $N_2$ , as a central convex body in that real four-dimensional space. Let  $E$  be an ellipsoid inscribed in  $S_2$  and having maximal four-dimensional volume. Then, from a result of Dvoretzky and Rogers [4], it follows that there are two complex linearly independent vectors  $x$  and  $y$  in  $E \cap S_2$ .

The ellipsoid  $E$  induces a euclidean norm  $\|\cdot\|_E$  on  $N_2$ , and  $E$  is the unit sphere of  $N_2$  with respect to  $\|\cdot\|_E$ . Thus

$$\|x\| = \|y\| = \|x\|_E = \|y\|_E = 1.$$

Since  $E \subset S_2$ , we have that  $\|u\| < \|u\|_E$  for all  $u \in N_2$ .

Let  $x$  and  $y$  be the previous complex linearly independent vectors, and let  $\alpha$  and  $\beta$  be any two complex numbers. Since  $N_2$  is an inner product space with respect to  $\|\cdot\|_E$ , it follows that  $(*)$  holds for any  $\alpha x, \beta y \in N_2$ , i.e.

$$\int_G \|\alpha x + \gamma(g)\beta y\|_E^2 d\mu = \|\alpha x\|_E^2 + \|\beta y\|_E^2 = |\alpha|^2 + |\beta|^2.$$

Assume  $(*)$  and recall that  $\|\cdot\| < \|\cdot\|_E$ . Then

$$\begin{aligned} |\alpha|^2 + |\beta|^2 &= \|\alpha x\|^2 + \|\beta y\|^2 = \int_G \|\alpha x + \gamma(g)\beta y\|^2 d\mu \\ &< \int_G \|\alpha x + \gamma(g)\beta y\|_E^2 d\mu = |\alpha|^2 + |\beta|^2. \end{aligned}$$

Hence

$$\int_G (\|\alpha x + \gamma(g)\beta y\|_E^2 - \|\alpha x + \gamma(g)\beta y\|^2) d\mu = 0.$$

Since for every open subset  $U$  of  $G$ ,  $\mu(U) > 0$ , and since the integrand of the last integral is a nonnegative continuous function in  $g$ , we have that

$$\|\alpha x + \gamma(g)\beta y\|_E = \|\alpha x + \gamma(g)\beta y\|$$

for all  $g \in G$ . In particular for  $g = e$ ,  $\|\alpha x + \beta y\|_E = \|\alpha x + \beta y\|$  for all  $\alpha, \beta \in C$ .

Every vector  $u \in N_2$  is a linear combination of  $x$  and  $y$ , i.e.  $u = \alpha x + \beta y$ . Hence we proved that  $\|u\|_E = \|u\|$  for all  $u \in N_2$ .

From the standard result it follows that  $N$  is an inner product space. This completes the proof of the theorem.

**3.** The scope of our theorem is best viewed through the following examples.

*Example 1.* Let  $G = \mathbf{R}/2\pi\mathbf{Z}$ , with the quotient topology  $\gamma(t) = e^{it}$  and let  $\mu$  be the Lebesgue measure on  $G$ . Then  $N$  is an inner product space if and only if

$$\int_G \|x + e^{it}y\|^2 dt = \frac{1}{2\pi} \int_0^{2\pi} \|x + e^{it}y\|^2 dt = \|x\|^2 + \|y\|^2,$$

for all  $x, y \in N$ .

*Example 2.* Let  $\mu$  be the normalized arc length on  $\Gamma$ . On the  $n$ -dimensional torus  $G = \prod_{j=1}^n \Gamma_j$ , for each  $n$ -tuple  $n = (m_1, \dots, m_n)$  of integers, the function

$$\gamma_m(z_1, \dots, z_n) = \prod_{j=1}^n z_j^{m_j}$$

is a continuous homomorphism from  $G$  into  $C$ . Then  $N$  is an inner product space if and only if

$$\int_{\Gamma} \int_{\Gamma} \dots \int_{\Gamma} \left\| x + \left( \prod_{j=1}^n z_j^{m_j} \right) y \right\|^2 |dz_1| |dz_2| \dots |dz_n| = \|x\|^2 + \|y\|^2$$

for all  $x, y \in N$ .

In particular we have that  $N$  is an inner product space if and only if

$$\int_{\Gamma} \|x + z^n y\|^2 |dz| = \|x\|^2 + \|y\|^2$$

for all  $x, y \in N$ .

*Example 3.* A normed complex linear space  $N$  is an inner product space if and only if

$$\frac{1}{(2\pi)^n} \int_0^{2\pi} \int_0^{2\pi} \dots \int_0^{2\pi} \left\| \sum_{k=1}^n e^{it_k} x_k \right\|^2 dt_1 dt_2 \dots dt_n = \sum_{k=1}^n \|x_k\|^2$$

for all  $x_k \in N$ .

REFERENCES

1. A. J. Penico and Č. V. Stanojević, *An integral analogue to parallelogram law*, Proc. Amer. Math. Soc. **79** (1980), 427-430.
2. I. J. Schoenberg, *A remark on M. M. Day's characterization of inner-product spaces and a conjecture of L. M. Blumenthal*, Proc. Amer. Math. Soc. **3** (1952), 361-364.
3. M. M. Day, *Some characterizations of inner-product spaces*, Trans. Amer. Math. Soc. **82** (1947), 320-337.
4. A. Dvoretzky and C. A. Rogers, *Absolute and conditional convergence in normed linear spaces*, Proc. Nat. Acad. Sci. U.S.A. **36** (1950), 192-197.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MISSOURI-ROLLA, ROLLA, MISSOURI 65401