



01 Jan 1988

## A Fixed Point Theorem Revisited

Alberta Bollenbacher

Troy L. Hicks

*Missouri University of Science and Technology*

Follow this and additional works at: [https://scholarsmine.mst.edu/math\\_stat\\_facwork](https://scholarsmine.mst.edu/math_stat_facwork)



Part of the [Mathematics Commons](#), and the [Statistics and Probability Commons](#)

---

### Recommended Citation

A. Bollenbacher and T. L. Hicks, "A Fixed Point Theorem Revisited," *Proceedings of the American Mathematical Society*, vol. 102, no. 4, pp. 898 - 900, American Mathematical Society, Jan 1988.

The definitive version is available at <https://doi.org/10.1090/S0002-9939-1988-0934863-2>

This Article - Journal is brought to you for free and open access by Scholars' Mine. It has been accepted for inclusion in Mathematics and Statistics Faculty Research & Creative Works by an authorized administrator of Scholars' Mine. This work is protected by U. S. Copyright Law. Unauthorized use including reproduction for redistribution requires the permission of the copyright holder. For more information, please contact [scholarsmine@mst.edu](mailto:scholarsmine@mst.edu).

## A FIXED POINT THEOREM REVISITED

ALBERTA BOLLENBACHER AND T. L. HICKS

(Communicated by John B. Conway)

**ABSTRACT.** A version of a theorem commonly referred to as Caristi's Theorem is given. It has an elementary constructive proof and it includes many generalizations of Banach's fixed point theorem. Several examples illustrate the diversity that can occur.

**THEOREM 1 (CARISTI [1]).** *Suppose  $T: X \rightarrow X$  and  $\phi: X \rightarrow [0, \infty)$ , where  $X$  is a complete metric space and  $\phi$  is lower semicontinuous. If for each  $x$  in  $X$ ,*

$$(C) \quad d(x, Tx) \leq \phi(x) - \phi(Tx),$$

*then  $T$  has a fixed point.*

**THEOREM 2 [2].** *Suppose  $T: X \rightarrow X$ , where  $X$  is a metric space. Then there exists  $\phi: X \rightarrow [0, \infty)$  such that (C) holds if and only if the series*

$$(1) \quad \sum_{n=0}^{\infty} d(T^n x, T^{n+1} x)$$

*converges for all  $x \in X$ .*

For  $x \in X$ ,  $O(x, \infty) = \{x, Tx, T^2x, \dots\}$  is called the orbit of  $x$ .  $G: X \rightarrow [0, \infty)$  is  $T$ -orbitally lower semicontinuous at  $x$  if  $\{x_n\}$  is a sequence in  $O(x, \infty)$  and  $\lim x_n = \bar{x}$  implies  $G(\bar{x}) \leq \liminf G(x_n)$ . If (C) holds for every  $y \in O(x, \infty)$ , (1) holds for this  $x$ , since the sequence of partial sums is nondecreasing and bounded above by  $\phi(x)$ . This fact is used in the next theorem.

**THEOREM 3.** *Let  $(X, d)$  be a metric space. Suppose  $T: X \rightarrow X$  and  $\phi: X \rightarrow [0, \infty)$ . Suppose there exists an  $x$  such that*

$$(C) \quad d(y, Ty) \leq \phi(y) - \phi(Ty)$$

*for every  $y \in O(x, \infty)$ , and any Cauchy sequence in  $O(x, \infty)$  converges to a point in  $X$ . Then:*

- (1)  $\lim T^n x = \bar{x}$  exists.
- (2)  $d(T^n x, \bar{x}) \leq \phi(T^n x)$ .
- (3)  $T\bar{x} = \bar{x}$  iff  $G(x) = d(x, Tx)$  is  $T$ -orbitally lower semicontinuous at  $x$ .
- (4)  $d(T^n x, x) \leq \phi(x)$  and  $d(\bar{x}, x) \leq \phi(x)$ .

---

Received by the editors January 28, 1987.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 47H10; Secondary 54H25.

*Key words and phrases.* Fixed point theorem, Caristi's theorem, Banach's theorem.

©1988 American Mathematical Society  
 0002-9939/88 \$1.00 + \$.25 per page

PROOF. Observe that  $\{T^n x\}$  is Cauchy. For if  $m > n$ ,

$$\begin{aligned} d(T^n x, T^m x) &\leq d(T^n x, T^{n+1} x) + \cdots + d(T^{m-1} x, T^m x) \\ &= \sum_{k=n}^{m-1} d(T^k x, T^{k+1} x). \end{aligned}$$

We noted above that the series  $\sum_{k=0}^{\infty} d(T^k x, T^{k+1} x)$  converges. Thus, (1) follows.

$$\begin{aligned} 0 \leq d(T^n x, T^m x) &\leq \sum_{k=n}^{m-1} d(T^k x, T^{k+1} x) \\ &\leq \sum_{k=n}^{m-1} [\phi(T^k x) - \phi(T^{k+1} x)] = \phi(T^n x) - \phi(T^m x) \leq \phi(T^n x). \end{aligned}$$

Letting  $m \rightarrow \infty$  gives (2).

(3)  $x_n = T^n x \rightarrow \bar{x}$  and  $G$  is  $T$ -orbitally lower semicontinuous at  $x$  implies

$$0 \leq d(\bar{x}, T\bar{x}) = G(\bar{x}) \leq \liminf G(x_n) = \liminf d(T^n x, T^{n+1} x) = 0.$$

Thus  $T\bar{x} = \bar{x}$ .

Assume that  $T\bar{x} = \bar{x}$  and  $\{x_n\}$  is a sequence in  $O(x, \infty)$  with  $\lim x_n = \bar{x}$ . Then  $G(\bar{x}) = d(\bar{x}, T\bar{x}) = 0 \leq \liminf d(x_n, Tx_n) = \liminf G(x_n)$ .

$$\begin{aligned} d(x, T^n x) &\leq d(x, Tx) + d(Tx, T^2 x) + \cdots + d(T^{n-1} x, T^n x) \\ (4) \quad &\leq [\phi(x) - \phi(Tx)] + [\phi(Tx) - \phi(T^2 x)] + \cdots + [\phi(T^{n-1} x) - \phi(T^n x)] \\ &= \phi(x) - \phi(T^n x) \leq \phi(x). \end{aligned}$$

Letting  $n \rightarrow \infty$  gives  $d(x, \bar{x}) \leq \phi(x)$ .

COROLLARY [3]. Let  $(X, d)$  be a complete metric space and  $0 < k < 1$ . Suppose  $T: X \rightarrow X$  and there exists an  $x$  such that

$$(A) \quad d(Ty, T^2 y) \leq kd(y, Ty)$$

for all  $y \in O(x, \infty)$ . Then:

- (1)  $\lim T^n x = \bar{x}$  exists.
- (2)  $d(T^n x, \bar{x}) \leq k^n(1-k)^{-1}d(x, Tx)$ .
- (3)  $T\bar{x} = \bar{x}$  if and only if  $G(x) = d(x, Tx)$  is  $T$ -orbitally lower semicontinuous at  $x$ .
- (4)  $d(T^n x, x) \leq (1-k)^{-1}d(x, Tx)$  and  $d(\bar{x}, x) \leq (1-k)^{-1}d(x, Tx)$ .

PROOF. Set  $\phi(y) = (1-k)^{-1}d(y, Ty)$  for  $y \in O(x, \infty)$ . Let  $y = T^n x$  in (A). Then

$$d(T^{n+1} x, T^{n+2} x) \leq kd(T^n x, T^{n+1} x)$$

and

$$d(T^n x, T^{n+1} x) - kd(T^n x, T^{n+1} x) \leq d(T^n x, T^{n+1} x) - d(T^{n+1} x, T^{n+2} x).$$

Thus,

$$d(T^n x, T^{n+1} x) \leq (1-k)^{-1}[d(T^n x, T^{n+1} x) - d(T^{n+1} x, T^{n+2} x)]$$

or  $d(y, Ty) \leq \phi(y) - \phi(Ty)$ . (1), (3), and (4) are immediate.

For (2), (A) implies  $d(T^n x, T^{n+1} x) \leq k^n d(x, Tx)$ . Thus,

$$d(T^n x, \bar{x}) \leq \phi(T^n x) = (1 - k)^{-1} d(T^n x, T^{n+1} x) \leq k^n (1 - k)^{-1} d(x, Tx).$$

REMARKS. (1)  $\phi$  is not required to be lower semicontinuous, and (C) need only hold on  $O(x, \infty)$  for some  $x$ . Also, it may be as easy to check the lower semicontinuity of  $G$  as it is of  $\phi$ . Even when  $\phi$  is lower semicontinuous and (C) holds for all  $x$ , Caristi's theorem does not give  $T\bar{x} = \bar{x}$ , but  $Tx_0 = x_0$  for some  $x_0$  in  $X$ .

(2) Furthermore (2) and (4) give useful bounds that reduce to the usual Banach bounds in the Corollary and in Banach's theorem. It is shown in [3 and 4] that the Corollary includes many generalizations of Banach's fixed point theorem.

EXAMPLE 1. In this example,  $T\bar{x} \neq \bar{x}$ . However,  $\phi$  is continuous,  $T$  is lower semicontinuous,  $T$  is continuous except at one point, and  $d(x, Tx) = \phi(x) - \phi(Tx)$ .

Let  $X = [0, 1]$  and  $\phi(x) = x$  for  $x \in X$ . Put  $Tx = 0$  if  $x \in [0, \frac{1}{2}]$ , and  $Tx = \frac{1}{2}x + \frac{1}{4}$  if  $x \in (\frac{1}{2}, 1]$ . If  $x \in [0, \frac{1}{2}]$ ,  $d(x, Tx) = d(x, 0) = x$ , and  $\phi(x) - \phi(Tx) = \phi(x) - 0 = x - 0 = x$ . Similarly, if  $x \in (\frac{1}{2}, 1]$ ,  $d(x, Tx) = \frac{1}{2}x - \frac{1}{4} = \phi(x) - \phi(Tx)$ . Thus  $d(x, Tx) = \phi(x) - \phi(Tx)$  for all  $x$ . Clearly, 0 is the only fixed point of  $T$ . If  $x > \frac{1}{2}$ ,  $\lim T^n x = \frac{1}{2} \neq T(\frac{1}{2})$ .

EXAMPLE 2. This example shows that even if  $\phi$  and  $T$  are both continuous, you may have many fixed points. Let  $X$  be the subset of the plane, with the usual metric  $d$ , defined by  $X = \{(x, y) : 0 \leq x, y \leq 1\}$ . Let  $T(x, y) = (x, 0)$ .  $T(Tp) = Tp$  for every  $p \in X$ .  $0 = d(Tp, T^2p) \leq \frac{1}{2}d(p, Tp)$ . As in the Corollary, let  $\phi(p) = 2d(p, Tp)$  and  $d(p, Tp) \leq \phi(p) - \phi(Tp)$ .

EXAMPLE 3. In this example,  $T$  and  $\phi$  are discontinuous,  $\phi(x) = cd(x, Tx)$ ,  $\phi$  is lower semicontinuous, and  $d(x, Tx) \leq \phi(x) - \phi(Tx)$ .  $T: [-1, 1] \rightarrow [-1, 1]$  is defined by  $Tx = -1$ , if  $x < 0$  and  $Tx = x/4$  if  $x \geq 0$ . Note that  $d(Tx, T^2x) \leq \frac{1}{4}d(x, Tx)$ . As in the Corollary, put  $\phi(x) = \frac{4}{3}d(x, Tx)$ . If  $x < 0$ ,  $\lim T^n x = -1 = T(-1)$ . If  $x > 0$ ,  $\lim T^n x = 0 = T(0)$ . Also, 0 and  $-1$  are the only fixed points of  $T$ .

## BIBLIOGRAPHY

1. J. Caristi, *Fixed point theorems for mappings satisfying inwardness conditions*, Trans. Amer. Math. Soc. **215** (1976), 241-251.
2. J. Eisenfeld and V. Lakshmikantham, *Fixed point theorems on closed sets through abstract cones*, Appl. Math. Comput. **3** (1977), 155-167.
3. T. L. Hicks and B. E. Rhoades, *A Banach type fixed point theorem*, Math. Japon. **24** (1979), 327-330.
4. B. E. Rhoades, *Contractive definitions revisited*, Topological Methods in Nonlinear Functional Analysis, Contemp. Math., vol. 21, Amer. Math. Soc., Providence, R. I., 1983, pp. 190-205.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MISSOURI-ROLLA, ROLLA, MISSOURI 65401 (Current address of T. L. Hicks)

Current address (Alberta Bollenbacher): Department of Mathematics, Southeast Missouri State University, Cape Girardeau, Missouri 63701