
01 Jan 2023

OSCILLATION of SECOND-ORDER HALF-LINEAR NEUTRAL NONCANONICAL DYNAMIC EQUATIONS

Martin Bohner

Missouri University of Science and Technology, bohner@mst.edu

Hassan El-Morshedy

Said Grace

Irena Jadlovská

Follow this and additional works at: https://scholarsmine.mst.edu/math_stat_facwork



Part of the [Mathematics Commons](#), and the [Statistics and Probability Commons](#)

Recommended Citation

M. Bohner et al., "OSCILLATION of SECOND-ORDER HALF-LINEAR NEUTRAL NONCANONICAL DYNAMIC EQUATIONS," *Journal of Applied Analysis and Computation*, vol. 13, no. 5, pp. 2646 - 2658, Journal of Applied Analysis and Computation, Jan 2023.

The definitive version is available at <https://doi.org/10.11948/20220484>

This Article - Journal is brought to you for free and open access by Scholars' Mine. It has been accepted for inclusion in Mathematics and Statistics Faculty Research & Creative Works by an authorized administrator of Scholars' Mine. This work is protected by U. S. Copyright Law. Unauthorized use including reproduction for redistribution requires the permission of the copyright holder. For more information, please contact scholarsmine@mst.edu.

OSCILLATION OF SECOND-ORDER HALF-LINEAR NEUTRAL NONCANONICAL DYNAMIC EQUATIONS

Martin Bohner^{1,†}, Hassan El-Morshedy²,
Said Grace³ and Irena Jadlovská⁴

Abstract In this paper, we shall establish some new criteria for the oscillation of certain second-order noncanonical dynamic equations with a sublinear neutral term. This task is accomplished by reducing the involved nonlinear dynamic equation to a second-order linear dynamic inequality. We also establish some new oscillation theorems involving certain integral conditions. Three examples, illustrating our results, are presented. Our results generalize results for corresponding differential and difference equations.

Keywords Half-linear dynamic equation, delay, second-order, noncanonical, oscillation.

MSC(2010) 34C10, 34K11.

1. Introduction

The usual notation and terminology for time scales as can be found in Bohner and Peterson [10] will be used throughout. Here, we are concerned with obtaining some new criteria for the oscillation of second-order half-linear dynamic equations with a sublinear neutral term of the form

$$\left(a(y^\Delta)^\alpha\right)^\Delta(t) + q(t)f(x(g(t))) = 0, \quad t \geq t_0, \quad (\text{E})$$

where $y(t) = x(t) + p(t)x^\beta(\delta(t))$. For an arbitrary time scale \mathbb{T} (i.e., a nonempty closed subset of the real numbers) with $\sup \mathbb{T} = \infty$, we set $[t_0, \infty)_{\mathbb{T}} = [t_0, \infty) \cap \mathbb{T}$, and we assume throughout that

(H₁) $a, p, q \in C_{\text{rd}}([t_0, \infty)_{\mathbb{T}}, (0, \infty))$;

[†]The corresponding author.

¹Department of Mathematics and Statistics, Missouri S&T, Rolla, MO 65409-0020, USA

²Department of Mathematics, Faculty of Science, Damietta University, New Damietta 34517, Egypt

³Department of Engineering Mathematics, Faculty of Engineering, Cairo University, Orman, Giza 12000, Egypt

⁴Mathematical Institute, Slovak Academy of Sciences, Grešákova 6, 040 01 Košice, Slovakia

Email: bohner@mst.edu (M. Bohner), elmorshedy@yahoo.com (H. El-Morshedy), saidgrace@yahoo.com (S. Grace), jadlovaska@saske.sk (I. Jadlovská)

(H₂) $g, \delta \in C_{rd}([t_0, \infty)_{\mathbb{T}}, \mathbb{T})$ are nondecreasing with $\delta(t) \leq t, g(t) \leq t$, and

$$\lim_{t \rightarrow \infty} \delta(t) = \lim_{t \rightarrow \infty} g(t) = \infty;$$

(H₃) $\alpha \geq 1$ and $0 < \beta \leq 1$ are ratios of positive odd integers;

(H₄) $f \in C(\mathbb{R}, \mathbb{R})$ and $f(x)/x^\alpha \geq M > 0$ for $x \neq 0$.

Our presented results will also employ the hypotheses

$$A(t) := \int_t^\infty \frac{\Delta s}{a^{1/\alpha}(s)} < \infty \quad \text{for all } t \geq t_0; \tag{H_5}$$

$$\int_{t_0}^\infty q(s) \Delta s = \infty; \tag{H_6}$$

$$\lim_{t \rightarrow \infty} \frac{p(t)A^\beta(\delta(t))}{A(t)} = 0; \tag{H_7}$$

$$\lim_{t \rightarrow \infty} p(t) = 0. \tag{H_8}$$

Recall that a solution of (E) is a nontrivial real-valued function x satisfying (E) for sufficiently large t . Solutions vanishing identically in some neighborhood of infinity will be excluded from our consideration. A solution x of (E) is said to be oscillatory if it is neither eventually positive nor eventually negative; otherwise, it is called nonoscillatory. An equation itself is said to be oscillatory if all its solutions are oscillatory.

The problem of investigating oscillation criteria for various types of dynamic equations has been a very active research area over the past two decades, see [2, 5, 11, 30]. A large number of papers and monographs has been devoted to this problem; for some recent contributions, we refer to [1, 3, 6, 13, 14, 16–18, 25–29, 31] and the references contained therein. In particular, oscillatory behavior of solutions to half-linear equations has been the subject of numerous studies; see, e.g., the papers [7–9, 20–23] for more details. We point out that analysis of qualitative behavior of half-linear equations is important not only for the further development of oscillation theory, but for practical reasons too since half-linear equations have numerous applications in the study of p -Laplace equations, and so forth; see, e.g., the papers [7, 8, 19, 24] (we also refer to [19, 24] for models from mathematical biology, where oscillation and/or delay actions may be formulated by means of cross-diffusion terms).

The purpose of this paper is to provide some new oscillation criteria for (E) in noncanonical form, i.e., satisfying (H₅), via a comparison with second-order linear dynamic inequalities. We also establish new theorems involving integral conditions that ensure the oscillation of (E). For related results in the case of corresponding differential equations, i.e., $\mathbb{T} = \mathbb{R}$, we refer to [12, 20, 21]. For related results in the case of corresponding difference equations, i.e., $\mathbb{T} = \mathbb{Z}$, we refer to [4]. For related results in the time scales case, we also refer to [7–9, 15, 23].

After this introduction, Section 2 gives three auxiliary results that are needed in the proofs of our five main theorems in Section 3. We conclude the paper in Section 4 with three examples, illustrating our theoretical findings.

2. Auxiliary Results

In this section, we give some auxiliary results that are used in the remainder of this paper.

Lemma 2.1. *If (H₁)–(H₆) and (H₈) hold, then any eventually positive solution of (E) is eventually decreasing.*

Proof. Let x be an eventually positive solution of (E), say

$$x(t) > 0, \quad x(\delta(t)) > 0, \quad x(g(t)) > 0 \quad \text{for } t \in [t_1, \infty)_{\mathbb{T}} \quad (2.1)$$

for some $t_1 \geq t_0$. Then (E) implies $(a(y^\Delta)^\alpha)^\Delta < 0$ on $[t_1, \infty)_{\mathbb{T}}$, so that

$$a(y^\Delta)^\alpha \quad \text{is decreasing on } [t_1, \infty)_{\mathbb{T}}. \quad (2.2)$$

By (2.2), we either have

$$a(y^\Delta)^\alpha > 0, \quad \text{i.e., } y^\Delta > 0 \quad \text{on } [t_1, \infty)_{\mathbb{T}}, \quad (2.3)$$

or otherwise there exists $t_2 > t_1$ such that

$$a(y^\Delta)^\alpha < 0, \quad \text{i.e., } y^\Delta < 0 \quad \text{on } [t_2, \infty)_{\mathbb{T}}. \quad (2.4)$$

Assume now that (2.3) holds. By (2.2), we have

$$a(t)(y^\Delta(t))^\alpha \leq a(t_1)(y^\Delta(t_1))^\alpha \quad \text{for all } t \geq t_1.$$

Rearranging, we have

$$y^\Delta(t) \leq \left(\frac{a(t_1)(y^\Delta(t_1))^\alpha}{a(t)} \right)^{1/\alpha} \quad \text{for all } t \geq t_1.$$

Integrating this inequality from t_1 to $t \geq t_1$, we get

$$\begin{aligned} y(t) &\leq y(t_1) + a^{1/\alpha}(t_1)y^\Delta(t_1) \int_{t_1}^t \frac{\Delta s}{a^{1/\alpha}(s)} \\ &\stackrel{(2.3)}{\leq} y(t_1) + a^{1/\alpha}(t_1)y^\Delta(t_1)A(t_1) \quad \text{for all } t \geq t_1. \end{aligned}$$

Thus, due to (H₅), y is bounded above, and together with (2.3), y has a finite positive limit, say $\ell \in (0, \infty)$, at infinity. This implies that x is bounded and

$$\lim_{t \rightarrow \infty} x(t) = \lim_{t \rightarrow \infty} y(t) = \ell.$$

Hence, there exist $\varepsilon > 0$ and $t^* \geq t_1$ such that $x(g(t)) > \ell - \varepsilon > 0$ for all $t \geq t^*$. Therefore, (E) and (H₄) imply that

$$\left(a(y^\Delta)^\alpha \right)^\Delta(t) \leq -(\ell - \varepsilon)^\alpha M q(t) \quad \text{for all } t \geq t^*.$$

Integrating from t^* to $t \geq t^*$, we find

$$a(t)(y^\Delta(t))^\alpha \leq a(t^*)(y^\Delta(t^*))^\alpha - (\ell - \varepsilon)^\alpha M \int_{t^*}^t q(s) \Delta s$$

for all $t \geq t_*$. Letting $t \rightarrow \infty$ implies, in view of (H₆), that

$$\lim_{t \rightarrow \infty} a(t) (y^\Delta(t))^\alpha = -\infty,$$

contradicting (2.3). Hence, (2.4) holds, completing the proof. □

Lemma 2.2. *Assume (H₁)–(H₆) and (H₈). Let x be an eventually positive solution of (E). Then $w = a^{1/\alpha}y^\Delta$ satisfies eventually $w \leq 0$, $w^\Delta \leq 0$, and*

$$\alpha w^\Delta (w^\sigma)^{\alpha-1} \leq (w^\alpha)^\Delta \leq \alpha w^\Delta w^{\alpha-1}. \tag{2.5}$$

Proof. Suppose x satisfies (2.1). Lemma 2.1 yields $w \leq 0$ on $[t_2, \infty)_\mathbb{T}$. By Pötzsche’s chain rule [10, Theorem 1.90], we have

$$0 \stackrel{(E)}{\geq} (w^\alpha)^\Delta = \alpha w^\Delta \int_0^1 ((1-h)w + hw^\sigma)^{\alpha-1} dh. \tag{2.6}$$

Note that the integral on the right-hand side of (2.6) is nonnegative as $\alpha - 1$ is an even nonnegative integer divided by an odd positive integer. Thus, from (2.6), we obtain $w^\Delta \leq 0$. This implies $w^\sigma \leq w$ eventually. Hence,

$$w^\sigma \leq (1-h)w + hw^\sigma \leq w \quad \text{for all } h \in [0, 1].$$

Therefore, we obtain

$$w^{\alpha-1} \leq ((1-h)w + hw^\sigma)^{\alpha-1} \leq (w^\sigma)^{\alpha-1} \quad \text{for all } h \in [0, 1],$$

and thus,

$$w^{\alpha-1} \leq \int_0^1 ((1-h)w + hw^\sigma)^{\alpha-1} dh \leq (w^\sigma)^{\alpha-1}.$$

Hence

$$\alpha w^\Delta (w^\sigma)^{\alpha-1} \leq \alpha w^\Delta \int_0^1 ((1-h)w + hw^\sigma)^{\alpha-1} dh \leq \alpha w^\Delta w^{\alpha-1},$$

proving (2.5). □

Remark 2.1. Assuming (H₁)–(H₅), (H₇) implies (H₈). To see that, in view of (H₁)–(H₅), we have $\lim_{t \rightarrow \infty} A(t) = 0$ and

$$\frac{A^\beta(\delta(t))}{A(t)} \geq \frac{A^\beta(t)}{A(t)} = \frac{1}{A^{1-\beta}(t)}.$$

Then, (H₇) implies (H₈). Thus, in this situation, we may apply Lemmas 2.1 and 2.2.

Lemma 2.3. *Assume (H₁)–(H₇). Let x be an eventually positive solution of (E). Let $w = a^{1/\alpha}y^\Delta$. Let $P \in (0, 1)$. Then, eventually,*

$$y + wA \geq 0, \tag{2.7}$$

$$x \geq Py, \tag{2.8}$$

$$y/A \text{ is nondecreasing,} \tag{2.9}$$

and there exists $\gamma > 0$ such that

$$y/A \geq \gamma. \tag{2.10}$$

Proof. Suppose x satisfies (2.1). Lemma 2.2 yields $w \leq 0$ and $w^\Delta \leq 0$ on $[t_2, \infty)_{\mathbb{T}}$. Hence, $w(s) \leq w(t)$ for all $s \geq t \geq t_2$, so

$$y^\Delta(s) \leq \frac{w(t)}{a^{1/\alpha}(s)} \quad \text{for all } s \geq t \geq t_2.$$

Integrating this inequality from t to $u \geq t \geq t_2$, we find

$$-y(t) \leq y(u) - y(t) \leq w(t) \int_t^u \frac{\Delta s}{a^{1/\alpha}(s)} \quad \text{for all } u \geq t \geq t_2,$$

and letting $u \rightarrow \infty$ yields

$$-y(t) \leq w(t)A(t) \quad \text{for all } t \geq t_2,$$

which proves (2.7). Next, on $[t_2, \infty)_{\mathbb{T}}$, we have

$$\left(\frac{y}{A}\right)^\Delta = \frac{y^\Delta A - yA^\Delta}{AA^\sigma} = \frac{y^\Delta A + a^{-1/\alpha}y}{AA^\sigma} = \frac{wA + y}{a^{1/\alpha}AA^\sigma} \stackrel{(2.7)}{\geq} 0,$$

and so (2.9) holds, while

$$\frac{y(t)}{A(t)} \stackrel{(2.9)}{\geq} \frac{y(t_2)}{A(t_2)} =: \gamma > 0 \quad \text{for all } t \geq t_2$$

proves (2.10). Now, since $y(t) = x(t) + p(t)x^\beta(\delta(t)) \geq x(t)$, we get

$$\begin{aligned} x(t) &= y(t) - p(t)x^\beta(\delta(t)) \geq y(t) - p(t)y^\beta(\delta(t)) \\ &\stackrel{(2.9)}{\geq} y(t) - p(t) \left(\frac{A(\delta(t))y(t)}{A(t)} \right)^\beta \\ &= y(t) \left[1 - p(t) \frac{A^\beta(\delta(t))}{A(t)} \left(\frac{y}{A} \right)^{\beta-1}(t) \right] \\ &\stackrel{(2.10)}{\geq} y(t) \left[1 - \gamma^{\beta-1} p(t) \frac{A^\beta(\delta(t))}{A(t)} \right] \end{aligned}$$

for all $t \geq t_2$. From (H₇), there exists $t_3 \geq t_2$ such that

$$p(t) \frac{A^\beta(\delta(t))}{A(t)} \leq (1 - P)\gamma^{1-\beta} \quad \text{for all } t \geq t_3,$$

and thus (2.8) holds on $[t_3, \infty)_{\mathbb{T}}$. □

3. Main Results

Now, we present our five main oscillation results for (E). Throughout this section, we use the notation

$$Q(t) := Mq(t), \quad t \geq t_0.$$

The first two results give new criteria by comparing with second-order inequalities.

Theorem 3.1. *Assume (H₁)–(H₇). Let $P \in (0, 1)$. If*

$$\left(a^{1/\alpha}y^\Delta\right)^\Delta(t) + \frac{P^\alpha}{\alpha}A^{\alpha-1}(\sigma(t))Q(t)y(g(t)) \leq 0 \tag{3.1}$$

has no eventually positive decreasing solution, then (E) is oscillatory.

Proof. Assume x is a nonoscillatory solution of (E), satisfying (2.1). Let $w = a^{1/\alpha}y^\Delta$. Observe that Lemmas 2.1, 2.2, and 2.3 hold. On $[t_3, \infty)_{\mathbb{T}}$, we thus have

$$\begin{aligned} 0 &\stackrel{(E)}{=} \left[(w^\alpha)^\Delta + q(f \circ x \circ g)\right] \frac{(w^\sigma)^{1-\alpha}}{\alpha} \\ &\stackrel{(2.5)}{\geq} \left[\alpha w^\Delta (w^\sigma)^{\alpha-1} + q(f \circ x \circ g)\right] \frac{(w^\sigma)^{1-\alpha}}{\alpha} \\ &= w^\Delta + q \frac{(w^\sigma)^{1-\alpha}}{\alpha} (f \circ x \circ g) \\ &\stackrel{(H_4)}{\geq} w^\Delta + q \frac{(w^\sigma)^{1-\alpha}}{\alpha} M(x \circ g)^\alpha \\ &= w^\Delta + Q \frac{(w^\sigma)^{1-\alpha}}{\alpha} (x \circ g)^\alpha \\ &\stackrel{(2.8)}{\geq} w^\Delta + Q \frac{(w^\sigma)^{1-\alpha}}{\alpha} P^\alpha (y \circ g)^\alpha \\ &= w^\Delta + Q \left(\frac{w^\sigma}{y \circ g}\right)^{1-\alpha} \frac{P^\alpha}{\alpha} (y \circ g) \\ &\stackrel{(H_2)}{\geq} w^\Delta + Q \left(\left(\frac{w}{y}\right)^\sigma\right)^{1-\alpha} \frac{P^\alpha}{\alpha} (y \circ g) \\ &\stackrel{(2.7)}{\geq} w^\Delta + Q (A^\sigma)^{\alpha-1} \frac{P^\alpha}{\alpha} (y \circ g), \end{aligned}$$

so that (3.1) indeed has an eventually positive decreasing solution. This contradiction shows that no nonoscillatory solution of (E) can exist, and thus (E) is oscillatory. \square

Theorem 3.2. *Assume (H₁)–(H₇). Let $P \in (0, 1)$. If, for all $\gamma > 0$,*

$$\left(a(y^\Delta)^\alpha\right)^\Delta(t) + \gamma^{\alpha-1}P^\alpha A^{\alpha-1}(g(t))Q(t)y(g(t)) \leq 0 \tag{3.2}$$

has no eventually positive decreasing solution, then (E) is oscillatory.

Proof. Assume x is a nonoscillatory solution of (E), satisfying (2.1). Let $w = a^{1/\alpha}y^\Delta$. Observe that Lemmas 2.1, 2.2, and 2.3 hold. On $[t_3, \infty)_{\mathbb{T}}$, we thus have

$$\begin{aligned} (w^\alpha)^\Delta &\stackrel{(E)}{=} -q(f \circ x \circ g) \\ &\stackrel{(H_4)}{\leq} -qM(x \circ g)^\alpha \\ &\stackrel{(2.8)}{\leq} -qMP^\alpha(y \circ g)^\alpha \\ &= -P^\alpha Q \left(\frac{y}{A} \circ g\right)^{\alpha-1} (A \circ g)^{\alpha-1} (y \circ g) \end{aligned}$$

$$\stackrel{(2.10)}{\leq} -\gamma^{\alpha-1} P^\alpha Q(A \circ g)^{\alpha-1}(y \circ g),$$

so that (3.2) indeed has an eventually positive decreasing solution. This contradiction shows that no nonoscillatory solution of (E) can exist, and thus (E) is oscillatory. \square

The next three results deal with sufficient integral conditions to ensure oscillation of (E).

Theorem 3.3. *Assume (H₁)–(H₅) and (H₇). If*

$$\limsup_{t \rightarrow \infty} \left[A(t) \int_{t_1}^t A^{\alpha-1}(\sigma(s))Q(s)\Delta s + \frac{1}{A(g(t))} \int_t^\infty A^\alpha(\sigma(s))A(g(s))Q(s)\Delta s \right] > \alpha, \quad (3.3)$$

then (E) is oscillatory.

Proof. First note that it follows from (3.3) that there exists $P \in (0, 1)$ such that

$$\limsup_{t \rightarrow \infty} \left[A(t) \int_{t_1}^t A^{\alpha-1}(\sigma(s))Q(s)\Delta s + \frac{1}{A(g(t))} \int_t^\infty A^\alpha(\sigma(s))A(g(s))Q(s)\Delta s \right] > \frac{\alpha}{P^\alpha}. \quad (3.4)$$

Assume x is a nonoscillatory solution of (E), satisfying (2.1). Let $w = a^{1/\alpha}y^\Delta$. Observe that Lemmas 2.1, 2.2, and 2.3 hold. Notice that (3.4) implies that (H₆) holds. If not, i.e., if $\int_{t_0}^t q(s)\Delta s$ is bounded, then

$$A(t) \int_{t_1}^t A^{\alpha-1}(\sigma(s))Q(s)\Delta s \leq A(t) \int_{t_1}^t Q(s)\Delta s \rightarrow 0 \quad \text{as } t \rightarrow \infty$$

and

$$\begin{aligned} \frac{1}{A(g(t))} \int_t^\infty A^\alpha(\sigma(s))A(g(s))Q(s)\Delta s &\leq \int_t^\infty A^\alpha(\sigma(s))Q(s)\Delta s \\ &\leq A^\alpha(\sigma(t)) \int_t^\infty Q(s)\Delta s \rightarrow 0 \quad \text{as } t \rightarrow \infty, \end{aligned}$$

where we used (H₅) and the decreasing nature of A . However, this contradicts (3.4) and hence we proved our claim. Note also that we now reach (3.1) as in the proof of Theorem 3.1. On $[t_3, \infty)_{\mathbb{T}}$, we thus have

$$(wA + y)^\Delta = w^\Delta A^\sigma + wA^\Delta + y^\Delta = w^\Delta A^\sigma \stackrel{(3.1)}{\leq} -\frac{P^\alpha}{\alpha} (A^\sigma)^\alpha Q(y \circ g),$$

and integrating this inequality from t to $u \geq t \geq t_3$ yields

$$\begin{aligned} -(wA + y)(t) &\stackrel{(2.7)}{\leq} (wA + y)(u) - (wA + y)(t) = \int_t^u (wA + y)^\Delta(s)\Delta s \\ &\leq -\frac{P^\alpha}{\alpha} \int_t^u A^\alpha(\sigma(s))Q(s)y(g(s))\Delta s \\ &= -\frac{P^\alpha}{\alpha} \int_t^u A^\alpha(\sigma(s))Q(s) \frac{y(g(s))}{A(g(s))} A(g(s))\Delta s \\ &\stackrel{(2.9)}{\leq} -\frac{P^\alpha}{\alpha} \int_t^u A^\alpha(\sigma(s))Q(s) \frac{y(g(t))}{A(g(t))} A(g(s))\Delta s \end{aligned}$$

$$\stackrel{(H_2)}{\leq} -\frac{P^\alpha}{\alpha} \int_t^u A^\alpha(\sigma(s))Q(s) \frac{y(t)}{A(g(t))} A(g(s))\Delta s,$$

which upon letting $u \rightarrow \infty$ becomes

$$(wA + y)(t) \geq \frac{P^\alpha y(t)}{\alpha A(g(t))} \int_t^\infty A^\alpha(\sigma(s))Q(s)A(g(s))\Delta s \tag{3.5}$$

for all $t \geq t_3$. On the other hand, for $t \geq t_3$, we get

$$\begin{aligned} w(t) &\leq w(t) - w(t_3) = \int_{t_3}^t w^\Delta(s)\Delta s \\ &\stackrel{(3.1)}{\leq} -\frac{P^\alpha}{\alpha} \int_{t_3}^t A^{\alpha-1}(\sigma(s))Q(s)y(g(s))\Delta s \\ &\stackrel{(H_2)}{\leq} -\frac{P^\alpha}{\alpha} \int_{t_3}^t A^{\alpha-1}(\sigma(s))Q(s)y(t)\Delta s, \end{aligned}$$

i.e.,

$$-(wA)(t) \geq \frac{P^\alpha}{\alpha} A(t)y(t) \int_{t_3}^t A^{\alpha-1}(\sigma(s))Q(s)\Delta s \tag{3.6}$$

for all $t \geq t_3$. Combining now (3.5) and (3.6) yields

$$\begin{aligned} y(t) &= (wA + y)(t) - (wA)(t) \\ &\geq \frac{P^\alpha y(t)}{\alpha A(g(t))} \int_t^\infty A^\alpha(\sigma(s))Q(s)A(g(s))\Delta s \\ &\quad + \frac{P^\alpha}{\alpha} A(t)y(t) \int_{t_3}^t A^{\alpha-1}(\sigma(s))Q(s)\Delta s, \end{aligned}$$

which upon division by $y(t)$ results in

$$\begin{aligned} \frac{\alpha}{P^\alpha} &\geq \frac{1}{A(g(t))} \int_t^\infty A^\alpha(\sigma(s))Q(s)A(g(s))\Delta s \\ &\quad + A(t) \int_{t_3}^t A^{\alpha-1}(\sigma(s))Q(s)\Delta s \end{aligned}$$

for all $t \geq t_3$. Taking \limsup as $t \rightarrow \infty$ contradicts (3.4) and completes the proof. \square

Theorem 3.4. Assume (H_1) – (H_5) and (H_7) . Let $P \in (0, 1)$. If for any $\ell_1 \in [t_1, \infty)_{\mathbb{T}}$, there exists $\ell \in [\ell_1, \infty)_{\mathbb{T}}$ such that

$$\limsup_{t \rightarrow \infty} \int_\ell^t \left[\frac{P^\alpha}{\alpha} A^\alpha(\sigma(s))Q(s) - \frac{a^{-1/\alpha}(s)}{4A(\sigma(s))} \right] \Delta s > 1, \tag{3.7}$$

then (E) is oscillatory.

Proof. Assume x is a nonoscillatory solution of (E), satisfying (2.1). Let $w = a^{1/\alpha}y^\Delta$. Observe that Lemmas 2.1, 2.2, and 2.3 hold. Notice that (3.7) implies that (H_6) holds. For if $\int_\ell^t q(s)\Delta s$ is bounded, then $\int_\ell^t A^\alpha(\sigma(s))Q(s)\Delta s$ is also bounded, which, given that ℓ_1 is arbitrarily large, contradicts (3.7). Note also that we now

reach (3.1) as in the proof of Theorem 3.1. For this proof, we also introduce $v = w/y$. On $[t_3, \infty)_{\mathbb{T}}$, we have

$$\begin{aligned} v^\Delta &= \frac{w^\Delta y - wy^\Delta}{yy^\sigma} = \frac{w^\Delta}{y^\sigma} - \frac{wa^{1/\alpha}y^\Delta a^{-1/\alpha}}{yy^\sigma} \\ &= \frac{w^\Delta}{y^\sigma} - \frac{w^2 a^{-1/\alpha}}{yy^\sigma} = \frac{w^\Delta}{y^\sigma} - v^2 a^{-1/\alpha} \frac{y}{y^\sigma} \\ &\stackrel{(3.1)}{\leq} -\frac{P^\alpha}{\alpha} (A^\sigma)^{\alpha-1} Q \frac{y \circ g}{y^\sigma} - v^2 a^{-1/\alpha} \frac{y}{y^\sigma} \\ &\stackrel{(H_2)}{\leq} -\frac{P^\alpha}{\alpha} (A^\sigma)^{\alpha-1} Q - v^2 a^{-1/\alpha}, \end{aligned}$$

so that

$$\begin{aligned} (Av)^\Delta &= A^\Delta v + A^\sigma v^\Delta = -va^{-1/\alpha} + A^\sigma v^\Delta \\ &\leq -va^{-1/\alpha} - \frac{P^\alpha}{\alpha} (A^\sigma)^\alpha Q - A^\sigma v^2 a^{-1/\alpha} \\ &= -\frac{P^\alpha}{\alpha} (A^\sigma)^\alpha Q - \frac{a^{-1/\alpha}}{4A^\sigma} (1 + 2A^\sigma v)^2 + \frac{a^{-1/\alpha}}{4A^\sigma} \\ &\leq -\frac{P^\alpha}{\alpha} (A^\sigma)^\alpha Q + \frac{a^{-1/\alpha}}{4A^\sigma}. \end{aligned}$$

Integrating this last inequality from ℓ to $t \geq \ell \geq t_3$, we get

$$\begin{aligned} \int_\ell^t \left[\frac{P^\alpha}{\alpha} A^\alpha(\sigma(s))Q(s) - \frac{a^{-1/\alpha}(s)}{4A(\sigma(s))} \right] \Delta s &\leq A(\ell)v(\ell) - A(t)v(t) \\ &\leq -A(t)v(t) \stackrel{(2.7)}{\leq} -1, \end{aligned}$$

which contradicts (3.7) and completes the proof. \square

Theorem 3.5. Assume (H_1) – (H_5) and (H_7) . If

$$\int_{t_1}^\infty \left[\frac{1}{a(s)} \int_{t_1}^s A^\alpha(g(u))q(u)\Delta u \right]^{1/\alpha} \Delta s = \infty, \quad (3.8)$$

then (E) is oscillatory.

Proof. Assume x is a nonoscillatory solution of (E), satisfying (2.1). Let $w = a^{1/\alpha}y^\Delta$. Observe that Lemmas 2.1, 2.2, and 2.3 hold. Notice that (3.8) implies that (H_6) holds. Note therefore that we reach (3.2) as in the proof of Theorem 3.2. On $[t_3, \infty)_{\mathbb{T}}$, we have

$$\begin{aligned} (w^\alpha)^\Delta &\stackrel{(3.2)}{\leq} -\gamma^{\alpha-1} P^\alpha (A \circ g)^{\alpha-1} Q(y \circ g) \\ &= -\gamma^{\alpha-1} P^\alpha (A \circ g)^\alpha Q\left(\frac{y}{A} \circ g\right) \\ &\stackrel{(2.10)}{\leq} -\gamma^\alpha P^\alpha (A \circ g)^\alpha Q, \end{aligned}$$

which upon integrating from t_3 to $t \geq t_3$ yields

$$w^\alpha(t) \leq w^\alpha(t) - w^\alpha(t_3) = \int_{t_3}^t (w^\alpha)^\Delta(s) \Delta s$$

$$\begin{aligned} &\leq -\gamma^\alpha P^\alpha \int_{t_3}^t A^\alpha(g(s))Q(s)\Delta s \\ &= -\gamma^\alpha P^\alpha M \int_{t_3}^t A^\alpha(g(s))q(s)\Delta s. \end{aligned}$$

Rearranging yields

$$y^\Delta(t) \leq -\gamma P M^{1/\alpha} \left[\frac{1}{a(t)} \int_{t_3}^t A^\alpha(g(s))q(s)\Delta s \right]^{1/\alpha}$$

for all $t \geq t_3$. Integrating this inequality again from t_3 to $t \geq t_3$ gives

$$\begin{aligned} -y(t_3) \leq y(t) - y(t_3) &= \int_{t_3}^t y^\Delta(s)\Delta s \\ &\leq -\gamma P M^{1/\alpha} \int_{t_3}^t \left[\frac{1}{a(s)} \int_{t_3}^s A^\alpha(g(u))q(u)\Delta u \right]^{1/\alpha} \Delta s, \end{aligned}$$

and so

$$\int_{t_3}^t \left[\frac{1}{a(s)} \int_{t_3}^s A^\alpha(g(u))q(u)\Delta u \right]^{1/\alpha} \Delta s \leq \frac{y(t_3)}{\gamma P M^{1/\alpha}}$$

for all $t \geq t_3$, contradicting (3.8) and completing the proof. □

4. Examples

We conclude this paper by giving three illustrating examples, one for $\mathbb{T} = \mathbb{R}$, then for $\mathbb{T} = \mathbb{Z}$, then for a general time scale.

Example 4.1. Consider the second-order neutral differential equation

$$\left(t^6 \left(\left[x(t) + \frac{1}{t} x^{1/3} \left(\frac{t}{2} \right) \right]' \right)^3 \right)' + t^\lambda x^3 \left(\frac{t}{3} \right) = 0 \tag{4.1}$$

for $t \in [3, \infty)_{\mathbb{R}}$. Here, $\alpha = 3$, $\beta = 1/3$, $p(t) = 1/t$, $\delta = t/2$, $g(t) = t/3$, $M = 1$, $Q(t) = q(t) = t^\lambda$ for $\lambda \in \mathbb{R}$. Moreover, $A(t) = 1/t$ and (H_4) is easily verified. Using Theorem 3.5, we see that (4.1) is oscillatory for all $\lambda \geq 5$.

Example 4.2. Consider the second-order neutral difference equation

$$\begin{aligned} &\Delta \left((t^2 + t)^5 \Delta \left(\left[x(t) + \frac{1}{t} x^\beta(t-1) \right] \right)^5 \right) \\ &+ (t+1)^6 x^5(t-2)(2 + \sin x(t-2)) = 0, \quad t \in \mathbb{N}, \end{aligned} \tag{4.2}$$

where $\beta \in (0, 1]$ is a ratio of positive odd integers. Here, $\alpha = 5$, $a(t) = (t^2 + t)^5$, $p(t) = 1/t$, $\delta(t) = t - 1$, and $g(t) = t - 2$. We also have $M = 1$, $Q(t) = (t + 1)^6$, and $A(t) = \frac{1}{t}$. Therefore,

$$\lim_{t \rightarrow \infty} \frac{p(t)A^\beta(\delta(t))}{A(t)} = \lim_{t \rightarrow \infty} \frac{1}{(t-1)^\beta} = 0.$$

Thus (H_7) is satisfied. Note that

$$\frac{P^\alpha}{\alpha} A^\alpha(\sigma(s))Q(s) - \frac{a^{-1/\alpha}}{4A(\sigma(s))} = \frac{P^5}{5}(s+1) - \frac{1}{4s}.$$

Then (3.7) is clearly satisfied, and hence (4.2) is oscillatory according to Theorem 3.4.

Example 4.3. Consider the second-order neutral dynamic equation

$$\left(\sigma^3(t)t^3 \left(\left[x(t) + \frac{1}{t}x^{1/3} \left(\frac{t}{2} \right) \right]^\Delta \right)^3 \right)^\Delta + q_0\sigma^2(t)x^3(\lambda t) = 0 \quad (4.3)$$

for $t \in [1, \infty)_{\mathbb{T}}$, where $\lambda \in (0, 1]$ and $q_0 > 0$. Here, $\alpha = 3$, $\beta = 1/3$, $p(t) = 1/t$, $\delta(t) = t/2$, $g(t) = \lambda t$, $M = 1$, $Q(t) = q(t) = q_0\sigma^2(t)$, and

$$A(t) = \int_t^\infty \frac{1}{\sigma(s)s} \Delta s = \frac{1}{t}.$$

Condition (H_7) is clearly satisfied, since

$$\lim_{t \rightarrow \infty} \frac{p(t)A^\beta(\delta(t))}{A(t)} = \lim_{t \rightarrow \infty} \left(\frac{2}{t} \right)^{1/3} = 0.$$

Now, since

$$A(t) \int_{t_1}^t A^{\alpha-1}(\sigma(s))Q(s) \Delta s = \frac{1}{t} \int_{t_1}^t \frac{q_0}{\sigma^{3-1}(s)} \sigma^2(s) \Delta s = q_0 \frac{t-t_1}{t}$$

and

$$\frac{1}{A(g(t))} \int_t^\infty A^\alpha(\sigma(s))A(g(s))Q(s) \Delta s = \lambda t \int_t^\infty \frac{q_0\sigma^2(s)}{\lambda s\sigma^3(s)} \Delta s = q_0,$$

(3.3) takes the form

$$q_0 > \frac{3}{2}. \quad (4.4)$$

Hence, by Theorem 3.3, (4.3) is oscillatory if (4.4) holds. Similarly, condition (3.7) takes the form

$$\limsup_{t \rightarrow \infty} \int_{t_1}^t \left[\frac{P^3 q_0}{3} \frac{s}{\sigma(s)} - \frac{1}{4} \right] \frac{1}{s} \Delta s > 1,$$

and so Theorem 3.4 requires, for the oscillation of (4.3), that there exist $P \in (0, 1)$ and $\varepsilon > 0$ such that

$$\frac{P^3 q_0}{3} \frac{t}{\sigma(t)} > \frac{1}{4} + \varepsilon.$$

References

- [1] S. Abbas, S. S. Negi, S. R. Grace, et al., *Survey on qualitative theory of dynamic equations on time scale*, Mem. Differ. Equ. Math. Phys., 2021, 84, 1–67.
- [2] R. P. Agarwal, M. Bohner, T. Li and C. Zhang, *Comparison theorems for oscillation of second-order neutral dynamic equations*, Mediterr. J. Math., 2014, 11(4), 1115–1127.

- [3] H. A. H. Agwa, H. M. Arafa, M. Bohner and M. A. A. Naby, *Oscillation of second-order integro-dynamic equations with damping and distributed deviating arguments*, J. Nonlinear Convex Anal., 2022, 23(6), 1275–1288.
- [4] M. Bohner, H. A. El-Morshedy, S. R. Grace and I. Saĝer, *Oscillation of second-order nonlinear difference equations with sublinear neutral term*, Math. Morav., 2019, 23(1), 1–10.
- [5] M. Bohner, L. Erbe and A. Peterson, *Oscillation for nonlinear second order dynamic equations on a time scale*, J. Math. Anal. Appl., 2005, 301(2), 491–507.
- [6] M. Bohner, S. R. Grace and I. Jadlovská, *Asymptotic behavior of solutions of forced third-order dynamic equations*, Analysis (Berlin), 2019, 39(1), 1–6.
- [7] M. Bohner, T. S. Hassan and T. Li, *Fite–Hille–Wintner-type oscillation criteria for second-order half-linear dynamic equations with deviating arguments*, Indag. Math. (N.S.), 2018, 29(2), 548–560.
- [8] M. Bohner and T. Li, *Oscillation of second-order p -Laplace dynamic equations with a nonpositive neutral coefficient*, Appl. Math. Lett., 2014, 37, 72–76.
- [9] M. Bohner and T. Li, *Kamenev-type criteria for nonlinear damped dynamic equations*, Sci. China Math., 2015, 58(7), 1445–1452.
- [10] M. Bohner and A. Peterson, *Dynamic equations on time scales*, Birkhäuser Boston, Inc., Boston, MA, 2001. An introduction with applications.
- [11] M. Bohner and S. H. Saker, *Oscillation of second order nonlinear dynamic equations on time scales*, Rocky Mountain J. Math., 2004, 34(4), 1239–1254.
- [12] J. Džurina, S. R. Grace, I. Jadlovská and T. Li, *Oscillation criteria for second-order Emden–Fowler delay differential equations with a sublinear neutral term*, Math. Nachr., 2020, 293(5), 910–922.
- [13] S. G. Georgiev, *Functional dynamic equations on time scales*, Springer, Cham., 2019.
- [14] S. R. Grace, S. Abbas and J. R. Graef, *Oscillation of even order nonlinear dynamic equations on time-scales*, Math. Morav., 2022, 26(1), 47–55.
- [15] S. R. Grace, R. P. Agarwal, M. Bohner and D. O’Regan, *Oscillation of second-order strongly superlinear and strongly sublinear dynamic equations*, Commun. Nonlinear Sci. Numer. Simul., 2009, 14(8), 3463–3471.
- [16] B. Karpuz, *Sharp oscillation and nonoscillation tests for delay dynamic equations*, Math. Methods Appl. Sci., 2019, 42(9), 2993–3001.
- [17] Z. Kayar and B. Kaymakçalan, *Applications of the novel diamond alpha Hardy–Copson type dynamic inequalities to half linear difference equations*, J. Difference Equ. Appl., 2022, 28(4), 457–484.
- [18] N. Kılıç and O. Öcalan, *Oscillation criteria for first-order dynamic equations with nonmonotone delays*, Hacet. J. Math. Stat., 2021, 50(2), 318–325.
- [19] T. Li, N. Pintus and G. Viglialoro, *Properties of solutions to porous medium problems with different sources and boundary conditions*, Z. Angew. Math. Phys., 2019, 70(3), 1–18.
- [20] T. Li and Y. V. Rogovchenko, *Oscillation of second-order neutral differential equations*, Math. Nachr., 2015, 288(10), 1150–1162.

- [21] T. Li and Y. V. Rogovchenko, *Oscillation criteria for second-order superlinear Emden-Fowler neutral differential equations*, *Monatsh. Math.*, 2017, 184(3), 489–500.
- [22] T. Li and Y. V. Rogovchenko, *On the asymptotic behavior of solutions to a class of third-order nonlinear neutral differential equations*, *Appl. Math. Lett.*, 2020, 105, 1–7.
- [23] T. Li and S. H. Saker, *A note on oscillation criteria for second-order neutral dynamic equations on isolated time scales*, *Commun. Nonlinear Sci. Numer. Simul.*, 2014, 19(12), 4185–4188.
- [24] T. Li and G. Viglialoro, *Boundedness for a nonlocal reaction chemotaxis model even in the attraction-dominated regime*, *Differential Integral Equations*, 2021, 34(5–6), 315–336.
- [25] S. S. Negi, S. Abbas, M. Malik and S. R. Grace, *New oscillation criteria for p -Laplacian dynamic equations on time scales*, *Rocky Mountain J. Math.*, 2020, 50(2), 659–670.
- [26] Y. Qiu, K. S. Chiu, I. Jadlovská and T. Li, *Existence of nonoscillatory solutions to nonlinear higher-order neutral dynamic equations*, *Adv. Difference Equ.*, 2020, 475, 1–16.
- [27] P. Řehák, *A note on transformations of independent variable in second order dynamic equations*, in *Difference equations and discrete dynamical systems with applications*, 312 of Springer Proc. Math. Stat., Springer, Cham., 2020, 335–353.
- [28] S. H. Saker, M. M. Osman and D. R. Anderson, *Two weighted norm dynamic inequalities with applications on second order half-linear dynamic equations*, *Qual. Theory Dyn. Syst.*, 2022, 21(1), 1–26.
- [29] Y. Sui and S. Sun, *Oscillation of third order nonlinear damped dynamic equation with mixed arguments on time scales*, *Adv. Difference Equ.*, 2018, 233, 1–17.
- [30] C. Zhang, R. P. Agarwal, M. Bohner and T. Li, *Oscillation of second-order nonlinear neutral dynamic equations with noncanonical operators*, *Bull. Malays. Math. Sci. Soc.*, 2015, 38(2), 761–778.
- [31] Y. Zhou, A. Alsaedi and B. Ahmad, *Oscillation and nonoscillation theorems of neutral dynamic equations on time scales*, *Adv. Difference Equ.*, 2019, 404, 1–11.