
01 Sep 2015

Theoretical and experimental study of (e,2e) ionization of the CO₂ (1π_g) molecule at 250 eV

Mevlut Dogan

Zehra Nur Ozer

Murat Yavuz

Osman Alwan

et. al. For a complete list of authors, see https://scholarsmine.mst.edu/phys_facwork/1518

Follow this and additional works at: https://scholarsmine.mst.edu/phys_facwork

 Part of the [Physics Commons](#)

Recommended Citation

M. Dogan et al., "Theoretical and experimental study of (e,2e) ionization of the CO₂ (1π_g) molecule at 250 eV," *Journal of Physics: Conference Series*, vol. 635, no. 7, Institute of Physics - IOP Publishing, Sep 2015. The definitive version is available at <https://doi.org/10.1088/1742-6596/635/7/072070>

This Article - Conference proceedings is brought to you for free and open access by Scholars' Mine. It has been accepted for inclusion in Physics Faculty Research & Creative Works by an authorized administrator of Scholars' Mine. This work is protected by U. S. Copyright Law. Unauthorized use including reproduction for redistribution requires the permission of the copyright holder. For more information, please contact scholarsmine@mst.edu.

Theoretical and experimental study of (e,2e) ionization of the CO₂ (1 π_g) molecule at 250 eV

Mevlut Dogan*¹, Zehra N. Ozer*, Murat Yavuz*, O. Alwan^{†‡}, A. Naja[†], B. B. Joulakian[‡],
Esam Ali⁺, C. Ning⁺⁺, Don Madison⁺

*Department of Physics, e-COL Laboratory, Afyon Kocatepe University, 03200, Afyon, Turkey

[†]Lebanese University, LPM, EDST, Tripoli, Lebanon

[‡]Université de Lorraine, SRSMC (UMR CNRS 7565), 1 bld Arago, bat. ICPM 57078 Metz Cedex 3, France

⁺Missouri University of Science & Technology, Rolla, MO 65409, USA

⁺⁺Tsinghua University, Beijing 100084, People's Republic of China

Synopsis Triple differential cross sections (TDCSs) of the electron-impact ionization of carbon dioxide are measured in the coplanar asymmetric geometry, with incident electron energy value of 250eV, and ejected electron of 37eV. We will report the experimental results in comparison with the theoretical calculations of the M3DW and TCC (type 5) calculations.

Not only its importance in applied fields from astrophysics to plasma chemistry, but also being the main component in the atmospheres of Venus and Mars, Carbon dioxide is an important molecule and it is very desirable to have available sets of cross sections for this gas. On the other side, information on cross sections for ionization of such atmospheric molecules by electron impact is important both for understanding the basic processes of atomic collisions and controlling events in the ionosphere.

In recent years the two-center Coulomb continuum (TCC) model was extended to three-center targets like CO₂ for low equal energy sharing regime and also relatively higher incident energy asymmetric regime [1-4]. The molecular three body distorted wave (M3DW) model contains the post collision interaction (PCI) between scattered and ejected electrons to all orders of perturbation theory and the distorted wave Born approximations (DWBA) contains PCI only to first order. The M3DW and DWBA have yielded good with experiment for several different molecular targets. The aim of this work is to compare experimental and theoretical results for (e,2e) ionization of CO₂. For coplanar asymmetric kinematics, two outgoing electrons from the same ionization event are detected with specified kinetic energies and angles. The scattered electron was detected at three fixed angles (-10⁰, -20⁰, -30⁰), while the ejected electron energy is fixed to 37 eV. The results are measured for ionization of the CO₂ (1 π_g) orbital at an incident electron energy of 250 eV.

Figure 1 compares experimental results with the predictions of the DWBA, M3DW and TCC

(type5) model. We will discuss the characteristics of the molecular results in combination with experiment and theory during the conference.




Figure 1. TDCS for CO₂ plotted as function of the ejection angle for the 37 eV ejected electron. The experimental results are compared to TCC, M3DW, and DWBA calculations.

Acknowledgments

This work was supported by TUBITAK through grant 109T738 and AKU, BAPK through grant 14.HIZ.DES.80. The theoretical work of EA and DM was supported by the US National Science Foundation under Grant. No. PHY-1068237.

References

- [1] O. Chuluunbaatar and B. Joulakian 2010 *J. Phys. B: At. Mol. Opt. Phys.* **43** 155201.
- [2] M.J. Hussey and A.J. Murray 2005 *J. Phys. B: At. Mol. Phys.* **38** 2965.
- [3] A. Lahmam-Bennani et al 2009 *J. Phys. B: At. Mol. Opt. Phys.* **42** 235205.
- [4] O Alwan et al 2014 *J. Phys. B: At. Mol. Opt. Phys.* **47** 225201

¹ E-mail: mdogan@aku.edu.tr

