
01 Sep 2015

First Evidence of Interference Effects in the Ionization of N₂ Molecule

Zehra Nur Ozer

Hari Chaluvadi

Mevlut Dogan

James Colgan

et. al. For a complete list of authors, see https://scholarsmine.mst.edu/phys_facwork/1541

Follow this and additional works at: https://scholarsmine.mst.edu/phys_facwork



Part of the [Physics Commons](#)

Recommended Citation

Z. N. Ozer et al., "First Evidence of Interference Effects in the Ionization of N₂ Molecule," *Journal of Physics: Conference Series*, vol. 635, no. 7, Institute of Physics - IOP Publishing, Sep 2015.

The definitive version is available at <https://doi.org/10.1088/1742-6596/635/7/072071>

This Article - Conference proceedings is brought to you for free and open access by Scholars' Mine. It has been accepted for inclusion in Physics Faculty Research & Creative Works by an authorized administrator of Scholars' Mine. This work is protected by U. S. Copyright Law. Unauthorized use including reproduction for redistribution requires the permission of the copyright holder. For more information, please contact scholarsmine@mst.edu.

First Evidence of Interference effects in the ionization of N₂ molecule

Zehra N. Ozer^{*1}, Hari Chaluvadi[†], Mevlut Dogan^{*}, James Colgan[‡], Don Madison[†]

^{*} Department of Physics, e-COL Laboratory, Afyon Kocatepe University, 03200, Afyon, Turkey

[†] Department of Physics, Missouri University of Science and Technology, Rolla, Missouri, USA

[‡] Theoretical Division, Los Alamos National Laboratory, Los Alamos, New Mexico, USA

Synopsis We will present an experimental and theoretical investigation of triple differential cross sections for electron-impact ionization of nitrogen molecules at intermediate energies. A discussion of interference effects contained in the theoretical and experimental interference factors will be presented.

Double slit experiments with light depict the fundamentally probabilistic nature of quantum mechanical phenomena. The interference effects for quantum particles can be seen in triple differential cross sections (TDCSs) for electron impact ionization of diatomic molecules. Observation of such effects helps provide a detailed understanding of collision-induced reactions. The existence of Young-type interference patterns was recently observed in the ionization of diatomic molecules [1-2] and theoretically supported by calculations obtained using a molecular distorted-wave model. Evidence for double-slit interference effects was recently found for electron-H₂ scattering [3-5]. Although there have been several experimental attempts to find double-slit interference effects for electron-N₂ scattering, no experimental evidence had been found in prior work [6-8].

The I-factor, which is the ratio of molecular and atomic cross sections, represents a better test for interference than looking directly at the TDCS since it is not clear how interference effects will be manifested directly in the cross sections. In this work, we report a study looking for evidence of Young's double-slit interference in (e,2e) ionization of N₂ (3σ_g) by looking at the I-factor which, in this case, is the ratio of the molecular N₂ cross sections divided by the atomic nitrogen N cross sections [9]. Similar to the H₂ study, we do not have experimental data for ionization of atomic N. In this case, we use theoretical N cross sections calculated in the M3DW (molecular 3-body distorted wave) approximation as the denominator for both experiment and theory (as has been routinely done for heavy particle scattering [1,2]).

We compared experimental and theoretical (e,2e) cross sections and I-factors for

intermediate energy electron-impact ionization of the N₂ molecule in the scattering plane. We found reasonably good agreement between the theoretical M3DW TDCS results and experiment. We will discuss the evidence for Young's double-slit interference in (e,2e) ionization of N₂ (3σ_g) at intermediate energies by looking at the I-factor during the conference.

Acknowledgments

This work was supported by the Scientific and Technological Research Council of Turkey (TUBITAK) through grant 109T738 and AKU, BAPK through grant 14.FENED.06 and 14.HIZ.DES.80. The theoretical work of HC and DM was supported by US National Science Foundation under Grant. No. PHY-1068237.

References

- [1] D. Misra *et al* 2006 *Physical Review A*, **74**(6), [060701](#).
- [2] N. Stolterfoht *et al* 2001 *Physical Review Letters*, **87**(2), [023201](#).
- [3] D. Milne-Brownlie *et al* 2006 *Physical Review Letters*, **96**(23), [233201](#).
- [4] E. S. Casagrande *et al* 2008 *Journal of Physics B: Atomic, Molecular and Optical Physics*, **41**(2), [025204](#).
- [5] Z. N. Ozer *et al* 2013 *Physical Review A*, **87**(4), [042704](#).
- [6] J. Gao *et al* 2005 *Physical Review A*, **72**(3), [032721](#).
- [7] A. J. Murray *et al* 2006 *J. Phys. B: At. Mol. Opt. Phys.*, **39**, [3945-3956](#).
- [8] L. R. Hargreaves *et al* 2009 *Physical Review A*, **80**(6), [062704](#).
- [9] H. Chaluvadi *et al* 2015 submitted to *J. Phys. B: At. Mol. Opt. Phys.*

¹E-mail: zehraengil@aku.edu.tr

²E-mail: mdogan@aku.edu.tr

