

01 Jun 2007

Signatures of Black Holes at the LHC

Marco Cavaglia

Missouri University of Science and Technology, cavagliam@mst.edu

Romulus Godang

Lucien M. Cremaldi

Donald J. Summers

Follow this and additional works at: https://scholarsmine.mst.edu/phys_facwork

 Part of the [Physics Commons](#)

Recommended Citation

M. Cavaglia et al., "Signatures of Black Holes at the LHC," *Journal of High Energy Physics*, vol. 2007, no. 6, Institute of Physics - IOP Publishing, Jun 2007.

The definitive version is available at <https://doi.org/10.1088/1126-6708/2007/06/055>



This work is licensed under a [Creative Commons Attribution 4.0 License](#).

This Article - Journal is brought to you for free and open access by Scholars' Mine. It has been accepted for inclusion in Physics Faculty Research & Creative Works by an authorized administrator of Scholars' Mine. This work is protected by U. S. Copyright Law. Unauthorized use including reproduction for redistribution requires the permission of the copyright holder. For more information, please contact scholarsmine@mst.edu.

Signatures of black holes at the LHC

To cite this article: Marco Cavaglia *et al* JHEP06(2007)055

View the [article online](#) for updates and enhancements.

Related content

- [Seeing through the string landscape—a string hunter's companion in particle physics and cosmology](#)
Dieter Lüst
- [Supersymmetry and the LHC inverse problem](#)
Nima Arkani-Hamed, Gordon L. Kane, Jesse Thaler *et al.*
- [LHC signature of mirage mediation](#)
Won Sang Cho, Yeong Gyun Kim, Kang Young Lee *et al.*

Recent citations

- [Semiclassical geons at particle accelerators](#)
Gonzalo J. Olmo and D. Rubiera-Garcia
- [Critical formation of trapped surfaces in collisions of non-expanding gravitational shock waves in de Sitter space-time](#)
I.Ya. Arefeva *et al*
- [Hawking evaporation of cosmogenic black holes in TeV-gravity models](#)
P Draggiotis *et al*

Signatures of black holes at the LHC*

Marco Cavaglià,^a Romulus Godang,^{ab} Lucien M. Cremaldi^a and Donald J. Summers^a

^a*Department of Physics and Astronomy, University of Mississippi,
University, MS 38677-1848, U.S.A.*

^b*University of South Alabama, Department of Physics,
Mobile, AL 36688, U.S.A.*

E-mail: cavaglia@phy.olemiss.edu, godang@slac.stanford.edu,
cremaldi@phy.olemiss.edu, summers@phy.olemiss.edu

ABSTRACT: Signatures of black hole events at CERN's Large Hadron Collider are discussed. Event simulations are carried out with the Fortran Monte Carlo generator CATFISH. Inelasticity effects, exact field emissivities, color and charge conservation, corrections to semiclassical black hole evaporation, gravitational energy loss at formation and possibility of a black hole remnant are included in the analysis.

KEYWORDS: Large Extra Dimensions, Black Holes.

*Work supported in part by the U.S. Department of Energy contract DE-FG05-91ER40622.

Contents

1.	Introduction	1
2.	A quick look at the physics of mini black holes	2
3.	The CATFISH generator	3
4.	Analysis of black hole events	4
4.1	Visible and missing transverse momentum	4
4.2	Event shape	7
4.3	Jet parameters	8
5.	Conclusions and further developments	9

1. Introduction

If the fundamental scale of gravity is of the order of few TeVs [1], proton-proton collisions at CERN's LHC could lead to the formation of mini Black Holes (BHs) [2] and branes [3] (For reviews and further references, see refs. [4, 5]). The cross section for creation of a BH or brane with radius R is expected to be approximately equal to the geometrical Black Disk (BD) cross section $\sigma_{BD}(s, n) = \pi R^2(s, n)$, where \sqrt{s} is the Center of Mass (CM) energy of the colliding quanta and n is the number of extra dimensions. The semiclassical Hawking effect [6] provides a decay mechanism for BHs which makes them visible to a detector. The spectrum of massive excitations in string theories suggests that branes may also decay thermally [7]. Under the most favorable circumstances, the BH event rate at the LHC should be comparable to the $t\bar{t}$ event rate.

Until now, numerical studies of observational signatures have implemented the semiclassical picture outlined above. However, recent results have significantly modified our understanding of BH formation and evolution. It is thus timely and worthwhile to examine the observational signatures of BH events beyond the simple semiclassical picture. To this purpose, we have analyzed BH events at the LHC with the Fortran Monte Carlo (MC) generator CATFISH, which implements many of the accepted theoretical results in the literature [8, 9] and allows the comparison of different theoretical models of BH production and decay. MC generators with similar characteristics of CATFISH have already been successfully utilized to simulate BH production in ultrahigh-energy cosmic ray air showers [10] and in lepton colliders [11].

2. A quick look at the physics of mini black holes

Thorne’s hoop conjecture [12] states that an event horizon forms when a mass M is compacted into a region with circumference smaller than twice the Schwarzschild radius $R(M)$ in any direction. At the LHC, this process can be achieved by scattering two partons with CM energy larger than M and impact parameter smaller than R . Analytic and numerical results show that the BH event is inelastic due to emission of gravitational radiation [5]. If the collision is elastic, the hoop conjecture implies that the parton cross section for BH production is equal to the geometrical cross section σ_{BD} . Otherwise, the cross section is smaller and depends on the impact parameter. The collisional energy loss depends on the impact parameter and increases as the number of spacetime dimensions increases. Consensus is that the BH mass monotonically decreases with the impact parameter from a maximum of about 60-70% of the CM energy for head-on collisions [13–15]. However, other independent estimates suggest that the gravitational energy loss could be smaller [16, 17]. Note that these treatments are rigorous only for BHs larger than the Compton length of the colliding quanta [18]. Moreover, mass, spin, charge and finite-size effects of the incoming partons are neglected. Size and spin effects are expected to be mostly relevant around the Planck energy. Charge effects could dominate at higher energy. The pointlike approximation fails for directions transversal to the motion [19].

The total cross section for a super-Planckian BH event involving two nucleons is obtained by integrating the parton cross section over the Parton Distribution Functions (PDFs). If the BH mass depends on the impact parameter, the generally accepted formula for the total cross section in a proton-proton collision is

$$\sigma_{pp \rightarrow BH}(s, n) = \sum_{ij} \int_0^1 2zdz \int_{x_m}^1 dx \int_x^1 \frac{dx'}{x'} f_i(x', Q) f_j(x/x', Q) F \sigma_{BD}(xs, n), \quad (2.1)$$

where $f_i(\cdot, Q)$ are the PDFs with four-momentum transfer squared Q [20, 21] and z is the impact parameter normalized to its maximum value. The cutoff at small x is $x_m = M_{\min}^2 / (sy^2(z))$, where $y(z)$ and M_{\min} are the fraction of CM energy trapped into the BH and the minimum-allowed mass of the gravitational object, respectively. F is a form factor. The total cross section for the BD model is obtained by setting $F = 1$ and $y^2(z) = 1$. The momentum transfer is usually set to be M_{BH} or the Schwarzschild radius inverse. The lower cutoff on the fraction of the nucleon momentum carried by the partons is set by the minimum-allowed formation mass of the gravitational object, M_{\min} . This threshold is usually considered to be roughly equal to the minimum mass for which the semiclassical description of the BH is valid. However, this argument is based on Hawking’s semiclassical theory and may not be valid at energies equal to few times the Planck mass. For example, the existence of a minimum spacetime length l_m implies the lower bound on the BH mass [22, 23]:

$$M_{ml} = \frac{n+2}{8\Gamma\left(\frac{n+3}{2}\right)} (\sqrt{\pi} l_m M_{\star}/2)^{n+1} M_{\star}, \quad (2.2)$$

where M_{\star} is the fundamental Planck mass. BHs with mass less than M_{ml} do not exist, since their horizon radius would fall below the minimum-allowed length.

After its formation, the mini BH is believed to radiate excess multipole moments (balding phase), spin-down and then classically evaporate through the Hawking mechanism. At the end of the Hawking evaporation, the BH may undergo a non-thermal decay in a number n_p of hard quanta or leave a remnant. Although some progress has been made, a complete quantitative description of the BH evolution is not fully known. The better understood stage is the Hawking phase, for which (classical) field emissivities have recently been calculated for *all* Standard Model (SM) fields [24]. (For earlier works on spin-0, -1/2 and -1 fields see refs. [25].) For the minimal $SU(3) \times SU(2) \times U(1)$ SM, most of the BH mass is radiated as SM quanta on the brane, although the gravitational emission in the bulk cannot be neglected for high n . It should be stressed, however, that the effect of rotation and quantum corrections on BH emissivities is not clear. Onset of additional evaporation channels at trans-Planckian energies could also lead to a larger emission of undetectable non-SM quanta during the decay phase even in absence of rotation [26, 27]. Quantum gravitational effects and BH recoil [28] could also affect the emission of visible quanta on the brane. Examples of quantum gravitational effects are quantum thermal fluctuations and corrections to Hawking thermodynamics due to the existence of a minimum length [23]. In absence of a BH remnant, the final non-thermal decay is usually described phenomenologically by setting a cutoff on the BH mass of the order of the Planck mass, $Q_{\min} \sim M_{\star}$, and democratically distributing the energy to the quanta. The existence of a minimum length gives a natural means to set Q_{\min} . In this case, the modified thermodynamical quantities determine the endpoint of Hawking evaporation when the BH mass reaches M_{ml} .

3. The CATFISH generator

In this section we review the main characteristics of the CATFISH generator. CATFISH includes three models for BH formation and cross section: BD, Yoshino-Nambu (YN) graviton loss model [13], and Yoshino-Rychkov (YR) graviton loss improved model [14]. Since the differences between the YN and YR models are not significant, only the latter has been used in the analysis below. The distribution of the initial BH masses is sampled from the differential cross section. CATFISH uses the cteq5m1 PDF distribution [20, 29]. (The use of different PDF distributions does not significantly affect the total and differential cross sections. For a detailed discussion on the uncertainties in the cross section due to the PDFs, see ref. [30].) Following earlier studies [31], the momentum transfer is set to $Q = \min \{M_{BH} \text{ or } R(M_{BH}), Q_{\max}\}$, where Q_{\max} is the maximum value allowed by the PDFs. The part of CM energy of the pp collision which is not trapped or lost in gravitational radiation forms the beam remnant, which is hadronized by PYTHIA [32]. Energy losses in the balding and spin-down phases are assumed to be either negligible or included in the energy loss during formation.

Exact classical emissivities of non-rotating spherically-symmetric BHs are implemented in the Hawking phase [24]. The particle content at trans-Planckian energy is assumed to be the minimal $SU(3) \times SU(2) \times U(1)$ SM with three families and a single Higgs boson on a thin brane. For black holes with mass \sim few TeV the Hawking temperature is generally above 100 GeV. Therefore, all SM degrees of freedom are considered massless. Presence of

a minimum length may affect the evaporation phase and is implemented in CATFISH. The MC uses the dimensionless parameter $\alpha = l_m M_\star/2$ to determine the minimum length [22, 23]. If there is no minimum length, the MC evaporates the BH according to the Hawking theory with varying temperature. Alternatively, the BH evolution proceeds according to the modified thermodynamics of ref. [22, 23]. The evaporation ends with a stable BH remnant or an explosive n_p -body decay when the BH reaches the mass Q_{\min} . Color charge is always conserved in the decay process. Conservation of EM charge can be turned off to make the BH remnant electrically charged. Four-momentum is conserved at each step in the evaporation process by taking into account the recoil of the BH on the brane due to the emission of the Hawking quanta. The initial energy of the BH is distributed democratically among all the Hawking quanta with a tolerance of $\pm 10\%$. Beam remnant, fragmentation, and initial- and final-state radiation are dealt with PYTHIA.

4. Analysis of black hole events

We focus on a purely statistical analysis of variables which allows an easy comparison with previous results [33–38] which have been obtained with the TRUENOIR [39] or CHARYBDIS [40] generators. A more refined analysis of other detector response-dependent signatures such as back-to-back di-jet suppression, di-lepton events ($\mu^+\mu^-$, μ^+e^- , μ^+e^+ , ...) will be presented in a future publication.

4.1 Visible and missing transverse momentum

Figure 1 shows missing transverse momentum (\cancel{P}_T) and visible transverse momentum of leptons and hadrons for 10,000 events at the LHC with the following parameters (benchmark):

$$n = 6, \quad M_{\min} = Q_{\min} = M_\star, \quad n_p = 4, \quad \alpha = 0,$$

BD cross section and conservation of EM charge. The momentum transfer is chosen as the Schwarzschild radius inverse. P_T cuts of 5 GeV on leptons (e, μ) and 15 GeV on photons + hadrons (γ, h) have been imposed to remove the beams and initial-state radiation. (These choices of cuts and momentum transfer apply to all simulations.) The plots show the total visible energy distribution, \cancel{P}_T and the visible transverse momentum of leptons (e, μ) and photons + jets (γ, h) with varying fundamental scale $M_\star = 1 \dots 3$ TeV. Figure 2 shows the results for three extra dimensions ($n = 3$). The results in figure 1 and figure 2 are in good agreement with simulations based on different BH generators [34].

A handful of BH events shows a large amount of transverse momentum up to several TeV, depending on the value of the fundamental scale and the number of extra dimensions. In the absence of a BH remnant, this missing transverse momentum is due to the emission of gravitons and other invisible quanta (e.g. neutrinos) in the various evolutionary phases of the BH (formation, Hawking evaporation and final explosive phase). The bulk of BH events is characterized by light, low-entropy BHs. Since the graviton and invisible channels accounts only for a small fraction of the total multiplicity in the decay phase, only rare high-mass events show a large amount of missing transverse momentum. A rough counting




Figure 1: Visible energy, \cancel{P}_T and visible transverse momentum of leptons and photons+jets (GeV) for the black disk model (BD) and fundamental Planck scale $M_* = 1, 2, 3$ TeV. The number of extra dimensions is $n = 6$ and the final BH decay is in four hard quanta.




Figure 2: Visible energy, \cancel{P}_T and visible transverse momentum of leptons and photons+jets (GeV) for the black disk model (BD) and fundamental Planck scale $M_* = 1, 2, 3$ TeV. The number of extra dimensions is $n = 3$ and the final BH decay is in four hard quanta.

of degrees of freedom shows that the hadronic-to-leptonic decay ratio of a BH event should be approximately 5:1. The prevalence of the hadronic channel on the leptonic channel is evident from the right panels of figure 1 and figure 2. Figures 1 and 2 also show the effect of the fundamental scale on visible energy and missing and visible transverse momentum. Increasing M_* leads to more massive BHs, i.e., higher multiplicity and harder quanta in




Figure 3: Visible energy, \cancel{P}_T and visible transverse momentum of leptons and photons+jets (GeV) for the black disk model (BD) and the Yoshino-Rychkov TS model (YR) in a ten-dimensional spacetime ($n = 6$) with fundamental Planck scale $M_\star = 1 \text{ TeV}$. The minimum formation mass of the BH is $M_{\min} = 1 \text{ TeV}$ or $M_{\min} = 2 \text{ TeV}$. The final BH decay is in four hard quanta ($n_p = 4$).

the Hawking phase. Therefore, higher values of M_\star tend to produce larger \cancel{P}_T . Visible transverse momenta show a similar pattern. Observation of events with high \cancel{P}_T would indicate high values of M_\star , independently of the details of BH formation and the number of extra dimensions. If BHs are observed at the LHC, M_\star could be measured to a certain degree of precision.

Missing and visible energy outputs depend on the initial BH mass, and thus from the number of extra dimensions. Graviton emission in the Hawking phase also increases with n [24], leading to a decrease in visible energy for higher-dimensional BHs (compare the upper-left panels of figure 1 and figure 2.) However, the variation in \cancel{P}_T due to spacetime dimensionality is much less significant than the change due to M_\star because of the high degree of sphericity of BH events (lower-left panels). Effects due to the dimensionality of spacetime are more evident for massive BHs, whereas most of the BHs produced at the LHC are very light. Therefore, it is unlikely that statistics alone will allow measurement of the number of extra dimensions.

Figure 3 shows the effects of changes in the minimum mass cutoff. Simulations separate quite easily different values of M_{\min} . However, since M_{\min} is a lower bound on the BH mass, increases in M_{\min} are akin to increases in M_\star (compare the upper-left panels of figure 1 and figure 3). Changes in M_{\min} are also entangled with the initial graviton emission, specially for massive events. In the BD model, larger values of M_{\min} (at fixed M_\star) lead to more massive BHs, and thus to higher visible transverse momenta. If the initial gravitational emission is turned on, this increase may be balanced by a decrease due to lower multiplicity (compare $M_{\min} = 1 \text{ TeV}$ for the BD model with $M_{\min} = 2 \text{ TeV}$ for the YR model). A measure of M_{\min} might prove to be difficult at the LHC.




Figure 4: Visible energy, \cancel{P}_T and visible transverse momentum of leptons and photons+jets (GeV) for the Yoshino-Rychkov TS model (YR) in a ten-dimensional spacetime ($n = 6$) with fundamental Planck scale $M_* = 1$ TeV and different final decay modes: neutral remnant ($n_p = 0$), two hard quanta ($n_p = 2$) and four hard quanta ($n_p = 4$).

Figure 4 displays the effects of the final explosive stage. Simulations show no statistical difference between decay in $n_p = 2$ and $n_p = 4$ quanta. Since the degrees of freedom in the final explosive phase are democratically chosen, a spectral analysis of the energy and the number of emitted quanta is required to distinguish the two models. Detection of a BH remnant stands a better chance because of larger \cancel{P}_T and smaller visible momentum due to the remnant undetectability. (See also refs. [41, 33].) Note that a large fraction of events with remnant produces very little visible output; most of the BHs are initially so light that the Hawking phase does not take place. On the contrary, the energy carried by the decay products is much larger than the invisible energy carried by the remnant for massive events.

Figure 5 compares BH events in a smooth spacetime ($\alpha = 0$) and a spacetime with minimum length equal to the fundamental Planck scale inverse ($\alpha = 0.5$). The simulations show no significant statistical differences between the two cases. The effects of a small distance cutoff becomes only relevant when the minimum scale is very close to the threshold of complete suppression of BH production. In this case, the minimum allowed mass eq. (2.2) is so large that BHs cannot form at the LHC CM energy. Therefore, observation of minimum length effects at the LHC requires a certain degree of fine tuning. It is unlikely that any information on quantum effects at the Planck scale can be extracted from LHC data.

4.2 Event shape

BH events are expected to be highly spherical because of the spherical nature of Hawking evaporation. The event shape can be quantified by means of the sphericity S and aplanarity




Figure 5: Visible energy, \cancel{P}_T and visible transverse momentum of leptons and photons+jets (GeV) for the black disk model (BD) and the Yoshino-Rychkov TS (YR) model in a ten-dimensional spacetime ($n = 6$) with fundamental Planck scale $M_* = 1$ TeV and zero ($\alpha = 0$) or M_*^{-1} ($\alpha = 0.5$) minimum length. The final BH decay is in two hard quanta ($n_p = 2$).

A [42], thrust and oblateness T [43], and Fox-Wolfram moment $R_1 \dots R_4$ variables [44]. Figure 6 shows sphericity, aplanarity, oblateness and thrust for a ten-dimensional model with fundamental Planck scale equal to 1 TeV, $M_{\min} = Q_{\min} = M_*$, no minimum length, different formation and final decay models. (Rare) massive BH events are characterized by very high sphericity and isotropy. A similar conclusion is reached by examining the second Fox-Wolfram moment (see first panel of figure 7). Increasing M_{\min} makes the events even more spherical because of the higher multiplicity in the decay phase.

Comparison between formation models at fixed n_p shows that more spherical events are obtained if the graviton loss is neglected; BHs are more massive and emit more quanta in the Hawking phase. The higher sphericity of BD events is evident from the central-right part of the plots, where Hawking emission dominates the emission in the final explosive phase. This makes the statistical difference between the formation models more clear. Comparison between $n_p = 2$ and $n_p = 4$ at given formation model shows that the former are less spherical than the latter. This effect is better displayed in the region of the plots corresponding to light BHs, where emission in the final phase dominates over Hawking emission. However, it should be stressed that the distinction between $n_p = 2$ and $n_p = 4$ at the LHC might be difficult due to the presence of non-BH background (e.g. $q\bar{q}$ events). Discrimination between alternative models of BH formation should be possible by selecting massive spectacular events with high sphericity.

4.3 Jet parameters

The upper-right and the lower panels of figure 7 show the number of jets and the heavy and light jet mass [32] for the choice of parameters discussed above, respectively. These plots




Figure 6: Sphericity (upper-left), Aplanarity (upper-right), Oblateness (lower-left) and thrust (lower-right) for the black disk model (BD) and the Yoshino-Rychkov TS model (YR) in a ten-dimensional spacetime ($n = 6$). The final black hole decay is in two hard quanta ($n_p = 2$) or four hard quanta ($n_p = 4$).

include initial- and final-state radiation jets in addition to the jets originated in the BH decay phase. As is expected, the BD model produces on average more jets than the model with graviton loss at formation (upper-right panel of figure 7). This is also evident from the right portions of the jet mass distributions, where the BD model is characterized by more massive jets than the YR model at fixed n_p . Therefore, measurement of high jet mass allows determination of the BH formation model independently of the shape variables. The left portions of the jet mass distributions are sensitive to the final BH decay. Final decay in $n_p = 2$ jets produces more heavy jets than final decay in $n_p = 4$ jets. Therefore, the measurement of low jet mass may give important information on the physics of the final BH phase.

5. Conclusions and further developments

The study of BH production at the TeV scale is now a few years old and entering the mature stage. With the LHC scheduled to begin operations soon, accurate simulations of BH events




Figure 7: R_2 (upper-left), Number of jets (upper-right), Heavy Jets mass (lower-left), Light Jets mass (lower-right). Fox-Wolfram moment R_2 , number of jets, heavy and light jet mass for the black disk model (BD) and the Yoshino-Rychkov TS model (YR) in a ten-dimensional spacetime ($n = 6$). The final black hole decay is in two hard quanta ($n_p = 2$) or four hard quanta ($n_p = 4$).

are a pressing need. These simulations should check the stability of the overall picture of BH production against improvements in the theory and give independent confirmation of previous results. In this paper we have investigated the signatures of BH events at the LHC with the MC generator CATFISH. CATFISH implements several features of BH production at the TeV scale which were not included in previous generators [9]. Our analysis has shown that the main signatures of BH production at the LHC (missing transverse momentum, high sphericity, high jet multiplicity) do not depend significantly on the fine details of BH formation and evolution. Measurement of the fundamental Planck scale and detection of a BH remnant could possibly be extracted from LHC data. On the other hand, discerning different models of BH formation and evolution at the LHC might prove difficult on a purely statistical basis

Several other interesting signatures of BH formation in particle colliders have been proposed in the literature (see, e.g., refs. [33, 34, 36–38]). In particular, suppression of high-energy back-to-back-correlated di-jets with energy above the fundamental scale and

di-lepton production with large transverse momentum are expected to be two of the most interesting signatures of BH production at the LHC. Investigation of these signatures with CATFISH is in progress. Detector response and event reconstruction are also fundamental issues to be addressed in a complete analysis of BH events at the LHC. Further work along these lines is currently being pursued.

Acknowledgments

The authors would like to thank Vitor Cardoso, David Cline, Greg Landsberg, Alexander Melnitchouk, Robert Palmer, Hans Wenzel and Graham Wilson for discussions and many useful suggestions. This work was supported in part by the U.S. Department of Energy contract DE-FG05-91ER40622.

References

- [1] N. Arkani-Hamed, S. Dimopoulos and G.R. Dvali, *The hierarchy problem and new dimensions at a millimeter*, *Phys. Lett.* **B 429** (1998) 263 [[hep-ph/9803315](#)];
I. Antoniadis, N. Arkani-Hamed, S. Dimopoulos and G.R. Dvali, *New dimensions at a millimeter to a Fermi and superstrings at a TeV*, *Phys. Lett.* **B 436** (1998) 257 [[hep-ph/9804398](#)];
N. Arkani-Hamed, S. Dimopoulos and G.R. Dvali, *Phenomenology, astrophysics and cosmology of theories with sub-millimeter dimensions and TeV scale quantum gravity*, *Phys. Rev.* **D 59** (1999) 086004 [[hep-ph/9807344](#)].
- [2] T. Banks and W. Fischler, *A model for high energy scattering in quantum gravity*, [hep-th/9906038](#);
S.B. Giddings and S.D. Thomas, *High energy colliders as black hole factories: the end of short distance physics*, *Phys. Rev.* **D 65** (2002) 056010 [[hep-ph/0106219](#)];
S. Dimopoulos and G.L. Landsberg, *Black holes at the LHC*, *Phys. Rev. Lett.* **87** (2001) 161602 [[hep-ph/0106295](#)];
K.-m. Cheung, *Black hole production and large extra dimensions*, *Phys. Rev. Lett.* **88** (2002) 221602 [[hep-ph/0110163](#)];
A. Chamblin and G.C. Nayak, *Black hole production at LHC: string balls and black holes from $p p$ and lead lead collisions*, *Phys. Rev.* **D 66** (2002) 091901 [[hep-ph/0206060](#)].
- [3] E.-J. Ahn, M. Cavaglià and A.V. Olinto, *Brane factories*, *Phys. Lett.* **B 551** (2003) 1 [[hep-th/0201042](#)];
E.-J. Ahn and M. Cavaglià, *A new era in high-energy physics*, *Gen. Rel. Grav.* **34** (2002) 2037 [[hep-ph/0205168](#)];
K. Cheung and C.-H. Chou, *p -brane production in fat brane or universal extra dimension scenario*, *Phys. Rev.* **D 66** (2002) 036008 [[hep-ph/0205284](#)].
- [4] M. Cavaglià, *Black hole and brane production in TeV gravity: a review*, *Int. J. Mod. Phys.* **A 18** (2003) 1843 [[hep-ph/0210296](#)];
G.L. Landsberg, *Black holes at future colliders and beyond*, *J. Phys.* **G32** (2006) R337 [[hep-ph/0607297](#)];
R. Emparan, *Black hole production at a TeV*, [hep-ph/0302226](#);
P. Kanti, *Black holes in theories with large extra dimensions: a review*, *Int. J. Mod. Phys.* **A 19** (2004) 4899 [[hep-ph/0402168](#)];

- S. Hossenfelder, *What black holes can teach us*, hep-ph/0412265.
- [5] V. Cardoso, E. Berti and M. Cavaglià, *What we (don't) know about black hole formation in high- energy collisions*, *Class. and Quant. Grav.* **22** (2005) L61 [hep-ph/0505125].
- [6] S.W. Hawking, *Particle creation by black holes*, *Commun. Math. Phys.* **43** (1975) 199 [Erratum *ibid.* **46** (1976) 206].
- [7] D. Amati and J.G. Russo, *Fundamental strings as black bodies*, *Phys. Lett.* **B 454** (1999) 207 [hep-th/9901092].
- [8] M. Cavaglià, R. Godang, L. Cremaldi and D. Summers, *CATFISH: a Monte Carlo simulator for black holes at the LHC*, *Comput. Phys. Commun.* in press, hep-ph/0609001.
- [9] D.M. Gingrich, *Comparison of black hole generators for the LHC*, hep-ph/0610219.
- [10] E.-J. Ahn and M. Cavaglià, *Simulations of black hole air showers in cosmic ray detectors*, *Phys. Rev. D* **73** (2006) 042002 [hep-ph/0511159].
- [11] R. Godang et al., *Resolution of nearly mass degenerate Higgs bosons and production of black hole systems of known mass at a muon collider*, *Int. J. Mod. Phys. A* **20** (2005) 3409 [hep-ph/0411248].
- [12] K.S. Thorne, in: *Magic without magic: John Archibald Wheeler*, edited by J. Klauder, Freeman, San Francisco (1972), gr-qc/0909003.
- [13] H. Yoshino and Y. Nambu, *Black hole formation in the grazing collision of high- energy particles*, *Phys. Rev. D* **67** (2003) 024009 [gr-qc/0209003].
- [14] H. Yoshino and V.S. Rychkov, *Improved analysis of black hole formation in high-energy particle collisions*, *Phys. Rev. D* **71** (2005) 104028 [hep-th/0503171].
- [15] O.I. Vasilenko, *Trap surface formation in high-energy black holes collision*, hep-th/0305067.
- [16] V. Cardoso, O.J.C. Dias and J.P.S. Lemos, *Gravitational radiation in d-dimensional spacetimes*, *Phys. Rev. D* **67** (2003) 064026 [hep-th/0212168].
- [17] E. Berti, M. Cavaglià and L. Gualtieri, *Gravitational energy loss in high energy particle collisions: ultrarelativistic plunge into a multidimensional black hole*, *Phys. Rev. D* **69** (2004) 124011 [hep-th/0309203].
- [18] M.B. Voloshin, *Semiclassical suppression of black hole production in particle collisions*, *Phys. Lett. B* **518** (2001) 137 [hep-ph/0107119]; *More remarks on suppression of large black hole production in particle collisions*, *Phys. Lett. B* **524** (2002) 376 [Erratum *ibid.* **B 605** (2005) 426] [hep-ph/0111099];
T.G. Rizzo, *Black hole production at the LHC: effects of Voloshin suppression*, *JHEP* **02** (2002) 011 [hep-ph/0201228];
V.S. Rychkov, *Black hole production in particle collisions and higher curvature gravity*, *Phys. Rev. D* **70** (2004) 044003 [hep-ph/0401116];
S.B. Giddings and V.S. Rychkov, *Black holes from colliding wavepackets*, *Phys. Rev. D* **70** (2004) 104026 [hep-th/0409131].
- [19] E. Kohlprath and G. Veneziano, *Black holes from high-energy beam-beam collisions*, *JHEP* **06** (2002) 057 [gr-qc/0203093].
- [20] CTEQ collaboration, R. Brock et al., *Handbook of perturbative QCD: version 1.0*, *Rev. Mod. Phys.* **67** (1995) 157.

- [21] PARTICLE DATA GROUP W.-M. Yao et al., *Review of particle physics*, *J. Phys. G* **33** (2006) 1.
- [22] M. Cavaglià, S. Das and R. Maartens, *Will we observe black holes at LHC?*, *Class. and Quant. Grav.* **20** (2003) L205 [[hep-ph/0305223](#)].
- [23] M. Cavaglià and S. Das, *How classical are TeV-scale black holes?*, *Class. and Quant. Grav.* **21** (2004) 4511 [[hep-th/0404050](#)].
- [24] V. Cardoso, M. Cavaglià and L. Gualtieri, *Black hole particle emission in higher-dimensional spacetimes*, *Phys. Rev. Lett.* **96** (2006) 071301 [*Erratum ibid.* **96** (2006) 219902] [[hep-th/0512002](#)]; *Hawking emission of gravitons in higher dimensions: non-rotating black holes*, *JHEP* **02** (2006) 021 [[hep-th/0512116](#)].
- [25] P. Kanti and J. March-Russell, *Calculable corrections to brane black hole decay. I: the scalar case*, *Phys. Rev. D* **66** (2002) 024023 [[hep-ph/0203223](#)];
 D. Ida, K.-y. Oda and S.C. Park, *Rotating black holes at future colliders: Greybody factors for brane fields*, *Phys. Rev. D* **67** (2003) 064025 [[hep-th/0212108](#)];
 P. Kanti and J. March-Russell, *Calculable corrections to brane black hole decay. II: greybody factors for spin 1/2 and 1*, *Phys. Rev. D* **67** (2003) 104019 [[hep-ph/0212199](#)];
 C.M. Harris and P. Kanti, *Hawking radiation from a $(4+n)$ -dimensional black hole: exact results for the schwarzschild phase*, *JHEP* **10** (2003) 014 [[hep-ph/0309054](#)]; *Hawking radiation from a $(4+n)$ -dimensional rotating black hole*, *Phys. Lett. B* **633** (2006) 106 [[hep-th/0503010](#)];
 D. Ida, K.-Y. Oda and S.C. Park, *Rotating black holes at future colliders. II: anisotropic scalar field emission*, *Phys. Rev. D* **71** (2005) 124039 [[hep-th/0503052](#)];
 E. Jung and D.K. Park, *Bulk versus brane in the absorption and emission: 5D rotating black hole case*, *Nucl. Phys. B* **731** (2005) 171 [[hep-th/0506204](#)].
- [26] M. Cavaglià, *Black hole multiplicity at particle colliders (Do black holes radiate mainly on the brane?)*, *Phys. Lett. B* **569** (2003) 7 [[hep-ph/0305256](#)].
- [27] A. Chamblin, F. Cooper and G.C. Nayak, *SUSY production from TeV scale blackhole at LHC*, *Phys. Rev. D* **70** (2004) 075018 [[hep-ph/0405054](#)].
- [28] V.P. Frolov and D. Stojkovic, *Black hole radiation in the brane world and recoil effect*, *Phys. Rev. D* **66** (2002) 084002 [[hep-th/0206046](#)].
- [29] CTEQ collaboration, H.L. Lai et al., *Global QCD analysis of parton structure of the nucleon: CTEQ5 parton distributions*, *Eur. Phys. J. C* **12** (2000) 375 [[hep-ph/9903282](#)].
- [30] E.-J. Ahn, M. Ave, M. Cavaglià and A.V. Olinto, *TeV black hole fragmentation and detectability in extensive air-showers*, *Phys. Rev. D* **68** (2003) 043004 [[hep-ph/0306008](#)].
- [31] L.A. Anchordoqui, J.L. Feng, H. Goldberg and A.D. Shapere, *Black holes from cosmic rays: probes of extra dimensions and new limits on TeV-scale gravity*, *Phys. Rev. D* **65** (2002) 124027 [[hep-ph/0112247](#)].
- [32] T. Sjostrand, S. Mrenna and P. Skands, *PYTHIA 6.4 physics and manual*, *JHEP* **05** (2006) 026 [[hep-ph/0603175](#)].
- [33] H. Stocker, *Stable TeV - black hole remnants at the LHC: discovery through di-jet suppression, mono-jet emission and a supersonic boom in the quark-gluon plasma*, [hep-ph/0605062](#).

- [34] C.M. Harris et al., *Exploring higher dimensional black holes at the Large Hadron Collider*, *JHEP* **05** (2005) 053 [[hep-ph/0411022](#)];
B. Webber, *Black holes at accelerators*, *ECONF C0507252* (2005) T030 [[hep-ph/0511128](#)].
- [35] G.L. Alberghi et al., *Probing quantum gravity effects in black holes at LHC*, [hep-ph/0601243](#).
- [36] J. Tanaka, T. Yamamura, S. Asai and J. Kanzaki, *Study of black holes with the ATLAS detector at the LHC*, *Eur. Phys. J. C* **41** (2005) 19 [[hep-ph/0411095](#)].
- [37] L. Lonnblad, M. Sjudahl and T. Akesson, *QCD-suppression by black hole production at the LHC*, *JHEP* **09** (2005) 019 [[hep-ph/0505181](#)].
- [38] G.C. Nayak and J. Smith, *Higgs boson production from black holes at the LHC*, *Phys. Rev. D* **74** (2006) 014007 [[hep-ph/0602129](#)].
- [39] S. Dimopoulos and G. Landsberg, *Black hole production at future colliders*, in *Proc. of the APS/DF/DPB Summer Study on the Future of Particle Physics (Snowmass 2001)* ed. N. Graf, *eConf C010630* (2001) P321.
- [40] C.M. Harris, P. Richardson and B.R. Webber, *CHARYBDIS: a black hole event generator*, *JHEP* **08** (2003) 033 [[hep-ph/0307305](#)].
- [41] B. Koch, M. Bleicher and S. Hossenfelder, *Black hole remnants at the LHC*, *JHEP* **10** (2005) 053 [[hep-ph/0507138](#)].
- [42] J. D. Bjorken and S. J. Brodsky, *Statistical model for electron - positron annihilation into hadrons*, *Phys. Rev. D* **1** (1970) 1416.
- [43] S. Brandt, C. Peyrou, R. Sosnowski and A. Wroblewski, *The principal axis of jets. An attempt to analyze high-energy collisions as two-body processes*, *Phys. Lett.* **12** (1964) 57.
- [44] G.C. Fox and S. Wolfram, *Observables for the analysis of event shapes in e^+e^- annihilation and other processes*, *Phys. Rev. Lett.* **41** (1978) 1581;
G.C. Fox and S. Wolfram, *Event shapes in e^+e^- annihilation*, *Nucl. Phys. B* **149** (1979) 413 [*Erratum ibid.* **B 157** (1979) 543].