

Fig. 8. Distribution chart and chronostratigraphy of sporomorph species recovered from Site 959. Values are uncorrupted, absolute numbers for each taxa. Askerisks denote important biostratigraphic taxa discussed in the text (Sections 6.2.1 to 6.2.5). [^]Chronostratigraphic ages after Berggren et al. (1995). *Biozones are for planktonic foraminifera. ♦Shafik et al. (1998) utilized calcareous nannofossils to constrain core ages. See text for further discussion.




Fig. 8 Continued. Distribution chart and chronostratigraphy of sporomorph species recovered from Site 959. Values are uncorrupted, absolute numbers for each taxa. Asterisks denote important biostratigraphic taxa discussed in the text (Sections 6.2.1 to 6.2.5). [^]Chronostratigraphic ages after Berggren et al. (1995). *Biozones are for planktonic foraminifera. ♦Shafik et al. (1998) utilized calcareous nannofossils to constrain core ages. See text for further discussion.




Fig. 8 Continued. Distribution chart and chronostratigraphy of sporomorph species recovered from Site 959. Values are uncorrupted, absolute numbers for each taxa. Asterisks denote imporant biostratigraphic taxa discussed in Section 6.2. ^Chronostratigraphic ages after Berggren et al. (1995). *Biozones are for planktonic foraminifera. ◇ Shafik et al. (1998) utilized nannofossils to constrain core ages. See text for further discussion.